



## Use of NMR to Determine Compatible Solutes in Halophilic Bacteria Isolated from Highly Saline Areas

**Reda Hassan Amasha**

*King Abdulaziz University, Saudi Arabia*

### Abstract:

Ten halophilic bacteria (two gram-negative) belonging to the Halomonadaceae and (eight gram-positive), belonging to the Bacillaceae, were isolated from the Red Sea, Arabian Gulf and Dead Sea using a high salinity medium, followed by identification using 16S rRNA. Four of the isolates were designated on the basis of their tolerance to high salinity. The isolates respectively exhibited 97% homology to *Halomonas aquamarina*, 97% homology to *Sediminibacillus* sp., (Red Sea), 94% homology to *Halobacillus* sp., (Arabian Gulf) and 98% homology to *Halobacillus dabanensis* (Dead Sea). <sup>1</sup>H-NMR spectroscopy was used to determine the osmolytes accumulated by *H. aquamarina*, *Sediminibacillus* sp., *Halobacillus* sp. and *H. dabanensis* grown in a saline nutrient medium at varying concentrations of NaCl and a range of organic sources. In the case of *H. aquamarina*, betaine and ectoine concentrations increased at high salinities. In contrast, betaine was found when casein and peptone were used as nutrient sources, while ectoine was produced in the presence of peptone. In the case of *Sediminibacillus* sp., betaine was the only osmolyte produced at high salinities, while betaine and ectoine was produced when peptone and casein were used. In *Halobacillus* sp., betaine was the only osmolyte produced at high salinities, whereas betaine and ectoine were produced in the presence of peptone and casein. Finally, in the case of *H. dabanensis*, only betaine accumulated at high salinities and in the presence of all organic nutrient sources.

### Biography:

Reda Hassan Amasha received her PhD in Molecular Biology and Biotechnology at University of Sheffield, UK in 2012 and she has done Diploma of Professional Development in teaching and learning in Center for Teaching and Learning Development from King Abdul Aziz University, Saudi Arabia, 2016. Her areas of research interest are in the occurrence of extremophile microorganism in non-extreme environments; molecular microbial diversity in caves and desert varnish; halophilic microorganisms and their environments and accumulation of compatible solutes as a strategy for adapting to salinity stress by using nuclear magnetic resonance (NMR) spectroscopy. She has published several papers in reputed



journals.

### Publication of speakers:

1. **RH Amasha** et al; Use Of Nmr To Determine Compatible Solutes In Halophilic Bacteria Isolated From Highly Saline Areas
2. **RH Amasha** et al; Microbial diversity and phylogenetic studies of some microbes obtained from unexplored caves of Saudi Arabia
3. **RH Amasha** et al; Isolation, molecular characterization and extracellular enzymatic activity of culturable halophilic bacteria from hypersaline natural habitats
4. **RH Amasha** et al; Isolation and identification of some antimicrobial producing bacteria from caves in Summan plateau, Saudi Arabia
5. **RH Amasha** et al; Nematodes Rotifers Tardigrades and Diatoms as Vehicles for the Panspermic Transfer of Microbes
6. **RH Amasha** et al; Studies on the Environmental Microbiology and Biogeochemistry of Desert Surface Soils

5<sup>th</sup> World Congress and Expo on Applied Microbiology; November 12-13, 2018; Edinburgh, Scotland.

**Citation:** Reda Hassan Amasha; Use of NMR to Determine Compatible Solutes in Halophilic Bacteria Isolated from Highly Saline Areas; Applied Microbiology 2018; November 12-13, 2018; Edinburgh, Scotland