

A Study on the Correlation between Post-Traumatic Stress Disorder and Social Support for the General Public during the Covid-19 Epidemic in China

Yilian Huang*¹

¹Guangdong Pharmaceutical University, China

Abstract

Objective: To investigate the occurrence and influencing factors of post-traumatic stress disorder syndrome in ordinary people during the outbreak of new coronary pneumonia, and to explore the relationship between general information, social support and PTSD. **Methods:** From March to April 2020, 704 ordinary people were randomly selected and measured using the general data questionnaire, the posttraumatic stress disorder scale (PCL-C) and the social support assessment scale (SSRS). **Results:** The incidence rate of PTSD -positive in 704 ordinary people is 10.51%, which was mainly affected by education level and the frequency of going out during the epidemic ($P < 0.05$). The total PCL-C score was negatively correlated with the level of social support and its dimensions, and positively correlated with age and annual income ($r = -0.03 \sim -0.12$, $P < 0.01$ or $P < 0.05$). **Conclusion:** Relevant departments should pay attention to the PTSD of ordinary people. In particular, people with low education level, high frequency of going out during the epidemic, high annual income and high age should be given positive psychological intervention to increase social support. .

Background

In December 2019, many new coronavirus pneumonias (new coronary pneumonia) were discovered in Wuhan City, Hubei Province, and the World Health Organization named it "COVID-19" as a public health emergency that caused international- attention [1]. Because of its high infectivity, wide -spread, and high fatality rate, it -poses a huge threat to the health and safety of Chinese citizens, and the people bear a strong psychological burden. Post-traumatic stress disorder (PTSD) refers to the adverse reactions and threats to the safety and health of an individual's life, resulting in continuation and persistence of negative emotions, which often occur with catastrophic and sudden Event related [2]. Studies have shown [2] that active coping strategies can reduce the incidence of PTSD. A survey [3] shows that social support can improve the negative emotions of hospitalized patients, and after traumatic events, social support is an effective factor for the prevention of PTSD [4,5]. Therefore, this study aims to investigate the PTSD situation and influencing factors of ordinary people during the new coronary pneumonia, and to explore the relationship between social support and PTSD, in order to clarify the health benefits of social support during the epidemic, and to formulate specific targets for future major public emergencies for reference.

Objects and Methods

Respondents

Using the convenience sampling method, 722 ordinary people were selected as survey subjects from March to April 2020. Inclusion criteria: (1)11-65 years old; (2) WeChat or QQ

number; (3) Chinese citizens; (4) voluntarily participate. EThe exclusion criteria were the unfulfillment of the inclusion criteria.

Method

Investigation tools

General information questionnaire: A self-made general information questionnaire is used, which includes general demographic information such as gender, age, place of residence, marital status, education level, occupation, and annual income.

Post-traumatic stress checklist-civilian version (PCL-C): The -scale was developed by the American Institute of Post-Traumatic Stress Disorder, and after multiple verifications later, it can be applied to domestic PTSD screening tools. The PCL-C scale has a total of 17 items and is divided into three major symptom groups: re-experience symptom group (5 items), avoidance/numbness symptom group (7 items), highly alert symptom group (5 items), according to Likert 5 score: 1=not at all, 2=mild, 3=moderate, 4=severe, 5=very severe. Among them, the retest reliability of the scale is 0.96, and the internal consistency reliability is 0.94, which has good validity and reliability [6]. Accumulate the score of each item to get a total score (17-85), with a 38-point cut-off value as a symptom of PTSD, a higher score indicates a higher risk of PTSD.

Social support rating scale (SSRS): The scale was compiled by Xiao Shui yuan [7] according to the situation in China, and has a total of 10 items, including 3 dimensions: objective support

(items 2,6,7), subjective support (items 1,3,4,5), and Utilization of social support (items 8,9,10). The stability coefficient of this scale is 0.92, and the internal consistency of each item is between 0.89 and 0.94, which is suitable for domestic research. Most questions are scored on level 4, and the sum of the scores of 10 items is the total score of the scale (12-66). The higher the score, the higher the level of social support.

Investigation method: Use WeChat, QQ and other online social platforms, use unified guidance, and use the "questionnaire star" point-to-point method to distribute and collect the questionnaire. The questionnaire set that each mobile phone number/ computer user can only fill in one questionnaire, respect the principle of the volunteer's voluntary investigation, adopt anonymous survey methods, and protect the privacy of the respondent. After the questionnaires were collected, the two persons checked and extracted the data. The number of questionnaires recovered was 704, and the effective recovery rate was 97.51%.

Statistical methods: Export the data on the questionnaire star website, and the data analysis will be performed by two people

Discussion

Analysis of PTSD and SSRS status of ordinary people during the new coronary pneumonia

Analysis of PTSD status of ordinary people during the new coronary pneumonia: The positive rate of PTSD surveyed in this study -was 10.51% (74/704), which was consistent with the positive rate of PTSD (1% ? -14%) for ordinary people screened abroad [8], which shows that the PTSD of ordinary people during the epidemic The incidence is high, and it is a high-risk group with PTSD. This study also found that the general population had the highest incidence of positive relapse symptom clusters, and the lower incidence of positive avoidance / numbness symptom clusters and highly alert symptom clusters, which is similar to the research results of Lin Zhizhi [9] and Xia Zheyuan [10]. This indicates that re-experience symptoms are more common during the outbreak of new coronary pneumonia, which verifies that reexperience symptoms are the most clinical manifestations of PTSD. At the same time, it shows that the occurrence of new coronary pneumonia is a negative stimulus and has a negative impact on the psychology of ordinary people. Some studies [11] found that depression is the most common disease of PTSD, which suggests that ordinary people with PTSD the danger of depression.

using SPSS 20.0 software. Counting data such as general data of the surveyed object are described by descriptive statistics, and are expressed by frequency and percentage; measurement data are analyzed by analysis of variance, t-test for statistical analysis, and expressed by $x \pm s$; single -factor analysis of variance and multiple stepwise Regression analysis of influencing factors of PTSD; Pearson correlation coefficient analysis method was used to analyze the relationship between different variables and PCL-C. $P < 0.05$ means the difference is statistically significant.

Results

PCL-C scale score and PTSD detection

In this survey, 74 patients with PCL-C total score ≥ 38 points had a PTSD -positive rate of 10.5%. The total score of PCL-C is (24.89 ± 9.17) , and the average scores of re-experience symptom group, avoidance / numbness symptom group, and high alertness symptom group are (1.46 ± 0.62) , (1.45 ± 0.57) , (1.49 ± 0.69) . The dimension with the highest average score is the highly alert symptom group. See Table 1 for details.

Analysis of the status of SSRS of ordinary people during the new coronary pneumonia: In this study, the general public's total social support score (41.45 ± 8.73) was higher than that during the SARS period [12,13]. This is due to the rapid and effective prevention and control measures taken by the party and the country during the new coronary pneumonia epidemic. A sense of security is guaranteed, and strong social support gives ordinary people enough confidence to meet and overcome the epidemic. Studies have shown that [14,15] a supportive social environment can encourage ordinary people to actively face traumatic events and maintain a good psychological state. Therefore, relevant departments should attach great importance to the occurrence of ordinary people's PTSD, seek positive and effective measures to mitigate the negative impact of the epidemic, and increase social support to reduce the psychological burden of traumatic events and promote the physical and mental health of ordinary-people.

Analysis of influencing factors of PTSD of ordinary people during the new coronary pneumonia

Education: The single -factor analysis of this study showed that the difference in education level in PCL-C scores was statistically significant ($P < 0.05$), and entered the regression equation, indicating that education level is an independent influencing factor of PTSD. This result is similar to Yvon M[16]. Similarly, PCL-C scores are low for those with a high degree of education, which is mainly due to the high level of education, which is easy to accept and understand the new coronary pneumonia disease, and those with a low degree of education are prone to wrong cognitive concepts [17],

especially in emergencies Negative cognition is more likely to occur when it occurs, resulting in an increased incidence of PTSD. This shows that the people with high education level can deal with the occurrence and development of emergencies rationally. They have high self-regulation ability and can self-adjust the psychological pressure caused by the new coronary pneumonia epidemic. This also suggests that when public health emergencies occur, society should give more psychological and social support to people with low education level.

Frequency of going out during an epidemic: The data of this study showed that the difference between the outgoing frequency during the outbreak and the PCL-C score was statistically significant ($P < 0.05$). That is, the lower the outgoing frequency during the outbreak, the lower the PCL-C score. It is an independent influencing factor of PTSD. New coronary pneumonia is highly contagious, spreads and can even cause death, and the population is generally susceptible, coupled with -external environmental factors and online public opinion is easy to increase the psychological pressure of the people. Therefore, the National Health Organization recommends that citizens reduce their outings and avoid gatherings to reduce the infection of the new coronavirus. Studies have reported that [18,19] the time of exposure to trauma events is positively correlated with the detection rate of PTSD, that is, the longer people are exposed to trauma events, the higher the PTSD detection rate. Therefore, the low frequency of people going out can reduce the time of exposure to traumatic events and reduce the risk of PTSD.

Relationship between PTSD and general information and social support during the new coronary pneumonia epidemic

The results of this study showed that the total score of PCL-C was positively correlated with age, and negatively correlated with education level, frequency of going out during the epidemic, total score of SSRS and its various dimensions ($P < 0.05$), that is, the higher the age, the post-traumatic stress The more severe the symptoms of the disorder, the higher the education level, the frequency of going out during the epidemic, social support, objective support, subjective support, and utilization of social support. The higher the score, the less severe the symptoms of post-traumatic stress disorder. This is similar to the results of the Xiaobo Z [20] study. When the exposed person faces a traumatic event, good social support as a protective factor can reduce the negative impact of the traumatic event, have a protective effect on mental health, and can be better accepted The impact of the epidemic, especially

those with a high level of education, are easier to accept and understand the epidemic. The older the people, the easier it is to generate negative emotions for the epidemic. This is mainly due to the greater family and social responsibilities and the psychological burden of the elderly. In general, it shows that the ordinary people have high utilization of social support during the epidemic, including the help of the party and the state to ordinary people in work and life. Remind relevant departments to continue to increase investment in social support in multiple ways, aspects and angles, pay more attention to the elderly and those with low education level and give more active social support to reduce the risk of PTSD.

The survey found that the avoidance/numbness symptom group was positively correlated with age, education, and annual income, and negatively correlated with the frequency of going out during the epidemic, total SSRS score, objective support, and utilization of social support ($P < 0.05$). Refers to the intention to avoid the traumatic event and choose to forget the details related to the traumatic event. The higher the people's annual income, the greater the avoidance/numbness symptoms. This is inconsistent with the findings of Wang Shuai [21]. This may be because the high-income people have low economic pressure, they choose to avoid the face of the epidemic and have a blind and confident attitude towards the epidemic. People with good economic conditions are more likely to have a sense of submission to the epidemic, intentionally evade details related to the epidemic, and passively face the epidemic to meet their psychological needs. This shows that during the epidemic, ordinary people feel the support and respect of the society and still have a hopeful and optimistic attitude towards life. However, the high-income people face the traumatic impact of the new coronary pneumonia epidemic and cannot choose to avoid it properly. Remind the relevant departments to pay more attention to ordinary people with better economic conditions during the epidemic to reduce the people's sense of submission and avoidance in the epidemic, so as to avoid or reduce the occurrence of PTSD-related symptoms.

Summary

In summary, the New Coronary Pneumonia epidemic is a major public health emergency. The incidence of PTSD among ordinary people is relatively high, mainly affected by the education level and the frequency of going out during the epidemic. Correlation analysis shows that social support and its various dimensions show the negative correlation is positively correlated with age and annual income, that is, increasing the level of social support is conducive to reducing the risk of PTSD. This suggests that relevant departments should attach great importance to the occurrence of PTSD among ordinary

people, especially those with low education, high frequency of outings during the epidemic, high annual income, and high age, seeking active and effective measures to mitigate the negative impact of the epidemic and increase social Support the strength and do a good job of psychological pressure, which will help the epidemic to provide a reference for further optimizing the psychological intervention of ordinary people.

This study has certain limitations: the time period and sample size of the investigation are limited, and it is impossible to understand the occurrence and outcome of PTSD symptoms from multiple angles, especially the single-factor analysis of the impact of PTSD at different stages. More sample sizes can be included and longitudinal studies can be conducted to explore the impact of PTSD in order to provide a basis for the development of targeted strategies for major emergencies in the future.

References

- Wang C, Horby PW, Hayden FG, Gao GF (2020) A novel coronavirus outbreak of global health concern. *Lancet* 395:470-473.
- Reynolds K, Pietrzak RH, Mackenzie CS, Chou KL, Sareen J (2016) Post-Traumatic Stress Disorder Across the Adult Lifespan: Findings From a Nationally Representative Survey. *Am J Geriatr Psychiat* 24:81-93.
- Drageset S, Lindstrøm TC, Giske T, Underlid K (2015) Women's experiences of social support during the first year following primary breast cancer surgery. *Scan J Caring Sci* 30:340-348.
- Doley RM, Bell R, Watt BD (2016) An Investigation Into the Relationship Between Long-term Posttraumatic Stress Disorder Symptoms and Coping in Australian Volunteer Firefighters. *J Nerv Ment Dis* 204:530-536.
- Nickerson A, Creamer M, Forbes D, McFarlane AC, O'Donnell ML, et al. (2016) The longitudinal relationship between post-traumatic stress disorder and perceived social support in survivors of traumatic injury. *Psych Med* 47:115-126.
- Nickerson A, Creamer M, Forbes D, McFarlane AC, O'Donnell ML, et al. (2016) The longitudinal relationship between post-traumatic stress disorder and perceived social support in survivors of traumatic injury. *Psych Med* 47:115-126.
- Xiao S (1994) The theoretical basis and research application of "Social Support Rating Scale". *J Clinical Psychi* 98-100.
- Forbes D, Creamer M, Biddle D (2001) The validity of the PTSD checklist as a measure of symptomatic change in combat-related PTSD. *Behav Res Ther* 39:977-986.
- Lin Z, Hu R (2017) Research on the correlation between post-traumatic stress disorder and stress and psychological resilience of nursing undergraduates. *Chinese J Nursing Edu* 14:770-773.
- Xia Z, Kong Y, Yin T, Shi S, Huang R, et al. (2014) The effect of disability acceptance and psychological resilience on the occurrence of post-traumatic stress disorder in burn patients. *Chin J Nur* 49:1035-1039.
- Karam EG, Friedman MJ, Hill ED, Kessler RC, McLaughlin KA, et al. (2014) Cumulative traumas and risk thresholds: 12-month PTSD in the World Mental Health (WMH) surveys. *Depression and Anxiety* 31:130-142.
- Hu S, Yang L, Li S, Wang X, Deng Z (2004) Analysis of social support and influencing factors of isolated people. *J Nurs Chinese PLA* 05: 33-34.
- Tong H (2004) "SARS" stress response pattern and its characteristics. *J Psychol* 01:103-109.
- Getnet B, Medhin G, Alem A (2019) Symptoms of post-traumatic stress disorder and depression among Eritrean refugees in Ethiopia: identifying direct, mediating and moderating predictors from path analysis. *BMJ Open* 9:e021142-e021142.
- Henry A, Tourbah A, Camus G, Deschamps R, Mailhan L, et al.(2019) Anxiety and depression in patients with multiple sclerosis: The mediating effects of perceived social support. *Mult Scler Relat Dis* 27:46-51.
- Motreff Y, Baubet T, Pirard P, Rabet G, Petitclerc M, et al. (2020) Factors associated with PTSD and partial PTSD among first responders following the Paris terror attacks in November 2015. *J Psych Res* 121:143-150.
- Li W, Ma L, Li X (2015) Post-traumatic stress disorder and influencing factors of breast cancer inpatients analysis. *Nursing Res* 29:174-178.
- Juárez-Ramírez C, Théodore FL, Villalobos A, Jiménez-Corona A, Lerin S, et al. (2015) Social Support of Patients with Type 2 Diabetes in Marginalized Contexts in Mexico and Its Relation to Compliance with Treatment: A Sociocultural Approach. *PLoS ONE* 10:e0141766-e0141766.
- Sheynin J, Duval ER, Lokshina Y, Scott JC, Angstadt M, et al. (2020) Altered resting-state functional connectivity in adolescents is associated with PTSD symptoms and trauma exposure. *Neuroimage Clin* 26:102215-102215.

20.Zhou X (2020) Psychological crisis interventions in Sichuan Province during the 2019 novel coronavirus outbreak. *Psychiatry Res* 286:112895-112895.

Wang S, Chen Y, Fu T, Zhang L (2019) The effect of psychological flexibility and social support on post-traumatic growth of colorectal cancer patients. *J Nur Manag* 19:6-10.

Citation: Huang Y (2020) A Study on the Correlation between Post-Traumatic Stress Disorder and Social Support for the General Public during the Covid-19 Epidemic in China. *J Comm Pub Health Nursing* 6: 243.

Copyright: © 2020 Huang Y, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.