

## An Overview of Traditionally Used Herb, *Colocasia esculenta*, as a Phytomedicine

Harshal Ashok Pawar<sup>1\*</sup>, Pritam Dinesh Choudhary<sup>2</sup> and Swati Ramesh Kamat<sup>2</sup>

<sup>1</sup>Department of Quality Assurance, Dr. L. H. Hiranandani College of Pharmacy, Ulhasnagar, Maharashtra, India

<sup>2</sup>Dr. L. H. Hiranandani College of Pharmacy, Ulhasnagar, Maharashtra, India

\*Corresponding author: Harshal Ashok Pawar, Department of Quality Assurance, Dr. L. H. Hiranandani College of Pharmacy, Ulhasnagar, Maharashtra, India, Tel: +918097148638; E-mail: hapkmk@rediffmail.com

Received date: October 08, 2017; Accepted date: Aug 20, 2018; Published date: Aug 30, 2018

Copyright: © 2018 Pawar HA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

*Colocasia esculenta* (*C. esculenta*) is a widely cultivated plant for consumption of both leaves and tubers. Taro, a common name for the corms and tubers of several genera of the family Araceae, is a source of edible corms of *C. esculenta*. It is majorly cultivated in Southeast Asia by several common names like Arbi, Arvi and Eddoe. From this species, a wide range of chemical compounds have been isolated for example, flavanoids,  $\beta$ -sitosterol, steroids, etc. Various parts of *C. esculenta* are used traditionally to treat number of diseases. Extracts from this plant have been found to possess various pharmacological activities. The leaves of this plant have been studied to possess anti-helminthic, anti-diabetic and anti-inflammatory actions. There have also been reports proving hypoglycemic efficacy of *C. esculenta* due to the presence of cyanoglucoside. The present review tries to cover all the information available on research work of this plant in yesteryears in a nutshell with the intention to serve as a literature platform for further researches on this plant.

**Keywords:** Phytomedicine; *Colocasia esculenta*; Araceae; Medicinal uses; Taro

### Introduction

In order to meet the global healthcare needs, traditional medicine systems have played a very important role. Indian Systems of Medicine being one of the largest and oldest systems constitutes all the medicines which have their origin in India. Plants have played a very vital role in sustaining and improving quality of human life and have served humans well as important components of medicines, seasoning, beverages, cosmetics, and dyes. Due to this fact, the focus on plant research has increased all over the world. There are any herbs that are used to treat liver, cardiovascular, digestive, metabolic and central nervous system (CNS) disorders. Medicinal plants or herbal drugs and their extracts containing isolated compounds have showcased a wide spectrum of biological activities. One of such a plant with wide applications is *C. esculenta*.

Taro, a common name for the corms and tubers of numerous genera of the family Araceae, is a source of edible corms of *C. esculenta*. It is most extensively cultivated in Southeast Asia by several common names like Arbi, Arvi and Eddoe. The juice of the plant leaf is rubefacient, stimulant, styptic and is useful in internal haemorrhages, adenitis, otalgia and buboes. The corm juice is demulcent, laxative and anodyne. The leaves of the plant have been found to exhibit anti-helminthic, anti-diabetic and anti-inflammatory activity [1]. *Colocasia* species is an ancient crop used all over the world; Africa, Asia, the West Indies, and South America. It is grown throughout the humid tropics. Its edible corms and leaves are traditionally used for liver related ailments [2]. *Colocasia esculenta* (L) Schott of the family Araceae is an herbaceous perennial plant. The crop is cultivated as annuals. The green leaves of the plant are large in size and are often described as 'Elephant ear'. They can reach up to 1-2 m high during

growth period. The main edible parts of the crop are the starchy, tuberous roots; however, the leaves of the plant are also used as a leafy vegetable. The leaves of *C. esculenta* have been reported to be rich in nutrients including minerals and vitamins such as phosphorous, calcium, vitamin C, iron, riboflavin, thiamine and niacin [3]. Among various edible plants from Araceae family commercially cultivated in India, *C. esculenta* has a significant dietary importance for having multiple uses in various culinary preparations of its edible parts i.e., corm, stem and leaves. *C. esculenta* fresh edible leaves are rich sources of protein, dietary fibre, ascorbic acid and some nutritionally important minerals. Leaf juice of the plant also finds its application in scorpion sting, in snake bite, in food poisoning from plant origin, etc. The plant also finds its application in Ayurveda identified ailments viz. vata and pitta, constipation, alopecia, stomatitis, haemorrhoids and general weakness [4,5]. *Colocasia antiquorum* is found to possess hepatoprotective activity against liver injury in rats that were experimentally induced [2]. *C. esculenta* is reported to possess hypoglycemic efficacy due to the presence of cyanoglucoside [6]. The plant is also reported to possess hypolipidemic and antihyperlipidemic activity due to the presence of arabinogalactan [7] and mono and digalactoyl diacylglycerols. The plant gets its antifungal property due to presence of cystatin [8]. Antibacterial activity of *C. esculenta* has been reported by Ravikumar et al. [9].

The objective of this review article was to shed light on the vast applications of traditionally used herb, *Colocasia esculenta*. This contribution also provides a broad review of botanical features, chemical constituents, pharmacological profile and medicinal/pharmaceutical applications of *C. esculenta*.

### Plant Botany

*Colocasia esculenta* Linn. (Figure 1) is a tall herb having tuberous or a stout short caudex, leafing and flowering together. Leaves are simple

and have a stout petiole, ovate-cordate or sagittate-cordate, lamina peltate. Spadix is shorter than the petiole and much it is shorter than the spathe rather than slender. Appendix much shorter than the inflorescence, style very short; stigma discoid [10]. It is erect, elongate-conical or fusiform, subulate or abbreviate. Erect petiole is up to 1.2 m in length, with a dull and non-polished surface above, coloured or paler beneath. They are rarely glaucous. The leaf peduncle is shorter than the petiole. Spathe is pale yellow and measures 15 to 35 cm in length; tube greenish, oblong. The lamina is narrowly lanceolate, convolute, acuminate and curved slightly backwards in flower. Female inflorescence is short but male inflorescence is long, cylindrical and usually interposed neuters between the two. Seeds oblong, sulcate. Albumen copious; embryo axile. The plant stem is above ground and swollen slightly at the base of the leaf-sheaths, arising from a hard-tapering rhizome; stolon's and a tuberous rhizome suckers are sometimes present.




Figure 1: Picture of parts of *Colocasia esculenta* plant.

### Leaf microscopy

Histological features of *C. esculenta* leaf are summarised below [11].

**Epidermis:** Single layer of spherical to polygonal cells with cutinized upper surface are present in upper epidermis of the leaf. The shape of the cells are straight to slightly beaded anticlinal walls and wavy with presence of Chlorophyll in epidermal cells. Lower epidermis shows single layer of polygonal cells which have straight to slightly beaded anticlinal walls. The lower epidermis also shows presence of papillae and paracytic type of stomata.

**Mesophyll:** Mesophyll shows dorsiventral arrangement, differentiated into palisade and spongy parenchyma. Chlorophyll and phenolic compounds are present in palisade cells.

**Spongy parenchyma:** The leaf shows presence of vacuoles and is made up of parenchymatous cells which varies in sizes and shapes measuring about 7-9 cells in thickness. Interspersed vascular elements are present intermittently. A majority of cells are filled with compound-type starch grains which are simple, spherical with centric helium and less prominent striations.

**Conducting tissue system:** Layer of parenchymatous bundle. The larger vascular bundles are surrounded by sclerenchymatous bundle sheath which extends up to the upper or lower or both epidermis [11]. The vascular bundles present are simple and each one of them are surrounded by a single.

### Phytochemistry

Flavonoids and triterpenoids are the two pharmacologically active groups of compounds mainly present in the *Colocasia* leaf extracts. The flavanoids that are present in the *Colocasia* leaf extracts are vicenin-2, iso-vitexin, iso-vitexin 3'-O-glucoside, vitexin X''-O-glucoside, iso-orientin, orientin, orientin7-O-glucoside, leteolin 7-O-glucoside [12]. The leaves of the plant also contains fibres, calcium oxalate, minerals (Calcium phosphorus etc.), and starch, Vitamin A, B, C, etc. [13]. Phytochemical investigations on the *Colocasia* extracts have shown the presence of anthocyanins viz. cyanidin-3-rhamnoside, cyanidin-3-glucoside and pelargonidin-3-glucoside. These anthocyanins have antioxidant activities as evident from previous studies [14-16]. It is due to these anthocyanins that the *C. esculenta* plant leaves showcases hepatoprotective as well as anti-lipid peroxidative activity. The tubers of *C. esculenta* contains globulins that accounts for around 80% of the total tuber proteins present. The starch content present in the flour varies from 73-76% and the starch yields are in the range of 51-58% [17]. The starch contains 0.23-0.52% lipid and 0.017-0.025% phosphorus in the form of phosphate monoester derivatives [17]. The nitrogen content of the flours varies from 0.33-1.35%. Table 1 shows chemical constituents of different parts of *C. esculenta* plant.

| Plant part | Chemical constituents  |                            |
|------------|--|----------------------------|
| Leaves     | Calcium oxalate, minerals like calcium phosphorus, fibres, starch, vitamin A, B, C |                            |
|            | Apigenin   |                            |
|            | Luteolin   |                            |
|            | Anthocyanin  |                            |
|            | Flavanoids   | Orientin                   |
|            |  | Iso-orientin               |
|            |  | Iso-vitexin                |
|            |  | Vicenin-2                  |
|            |  | Orientin 7-O-glucoside     |
|            |  | Iso-vitexin 3'-O-glucoside |

|   |  |                            |
|---|--|----------------------------|
|   |  | Vitexin X" -O-glucoside    |
|   |  | Luteolin 7-O-glucoside     |
| Tubers                                    | Starch                                 | 73-76%                     |
|   | Natural polysaccharide                 | 56% Natural sugars         |
|   |  | 40% Anionic components     |
|   | Oxalate                                | Soluble-19-87 mg/100 g     |
|   |  | Insoluble-33-156 mg/ 100 g |
|   | Amino acids                            | 13 to 23%                  |
|   | Nitrogen content                       | 0.33 to 1.35%              |
|   | Lipid                                  | 0.23 to 0.52%              |
|   | Phosphate monoester derivatives        | 0.017 to 0.025%            |
|   | Dihydroxysterols                       |                            |
|   | β-sitosterol                           |                            |
|   | Stigmasterol                           |                            |
|   | Nonacosane                             |                            |
|   | Cyaniding 3-glucoside                  |                            |
|   | Aliphatic compounds                    | Tetracos-20-en-1, 18-diol  |
|   |  | 25-methyl triacont-10-one  |
| Octacos-10-en-1, 12-diol                  |  |                            |
| Pentatriacont-1, 7-dien-12-ol             |  |                            |
| 25-methyl-tritriacont-2-en-1, 9, 11-triol |  |                            |
| Octadecenoic acid                         |  |                            |
| Enzymes                                   | Lipoxygenase                           |                            |
|   | Lipid hydro peroxide-converting enzyme |                            |
| Petiole                                   | Anthocyanins                           | 3.29%                      |

**Table 1:** Chemical constituents of different parts of *Colocasia esculenta* plant [12,13,17-21].

## Pharmacological Activities

### Antimicrobial activity

The aqueous extract of *C. esculenta* has been reported to possess antimicrobial activity. Study was performed for isolates of bacteria namely *Escherichia coli*, *Aeromonas hydrophila*, *Flavobacterium sp.*, *Edwardsiella tarda*, *Klebsiella sp.*, *Salmonella sp.*, *Vibrio alginolyticus*, *V. parahaemolyticus*, *V. cholerae* and *Pseudomonas aeruginosa*. The plant extract showed maximum activity against *S. mutans* amongst all the selected strains of microbe. The extract of *C. esculenta* exhibited good antimicrobial activity against some of bacteria and fungus that were tested at low concentration [22].

### Antihepatotoxic activity

*C. esculenta* leaf juice demonstrated hepatoprotective and anti-hepatotoxic activity. The hepatoprotective and anti-hepatotoxic studies were conducted against two well-known hepatotoxins paracetamol and  $\text{CCl}_4$  using *in vitro* rat liver slice method.  $\text{CCl}_4$  and Paracetamol generate free radical that causes oxidative stress. It also damages various cell organelles thus injuring the hepatocytes. The hepatoprotective efficacy was measured using the leakage of marker enzymes of liver function such as AST, ALP and ALT in the incubation medium. In presence of  $\text{CCl}_4$  and paracetamol, there was an increase in the levels of marker enzymes. Insignificant changes were noted in the enzymes levels at one and two hour's interval. At four hours interval, the marked elevations of toxicity marker enzymes were seen. However, the leaf juice of *C. esculenta* abnormally declined the leakage of AST, ALP and ALT in the medium demonstrating integrity of hepatocyte [23].

### Antidiabetic activity

The antidiabetic activity of the ethanol extract of *C. esculenta* (EECE) leaves was carried out in rats using alloxan induced diabetes model. EECE (100, 200 and 400 mg/kg) and metformin (450 mg/kg) were administered orally in alloxan (120 mg/kg, i.p.) induced diabetic rats. At 4 h (96 mg/dl) the onset of reduction of blood glucose was observed, peak at 6 h (120 mg/dl) but anti-hyperglycaemic effect diminished at 24 h. Maximum reduction in blood glucose was observed (174.34 mg/dl) in subacute study, at the dose of 400 mg/kg on 14th day. These results suggested that EECE (400 mg/kg) exhibits anti-hyperglycaemic activity in alloxan induced diabetic rats [24,25].

### Anti-lipid peroxidative activity

*C. esculenta* whole leaf juice has been reported to possess the free radical scavenging property. The *in vitro* free radical scavenging effect was studied on liver cells by using rat liver slice model [26]. The liver slices were incubated in presence of cytotoxic concentrations of CCl<sub>4</sub> and acetaminophen. The marked elevations and prevention of depletion of total tissue glutathione were observed in the presence of *C. esculenta* whole leaf juice [26].

### Antimetastatic activity

Breast cancer mortality is primarily due to the occurrence of metastatic disease. The compound(s) that are derived from roots of the plant of *C. esculenta* has the capability to potentially and specifically inhibit tumor metastasis. It exhibited demonstrable activity in a preclinical model of metastatic breast cancer. Taro extract treatment also inhibited prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) synthesis and down regulated cyclooxygenase 1 and 2 mRNA expression. Taro extract modestly inhibits the proliferation of some, but not all, breast and prostate cancer cell lines and it completely blocks tumor cell migration [27].

### Antifungal activity

Yang et al. evaluated the antifungal activity of taro along with molecular cloning and recombinant gene expression studies. CeCPI, a

cysteine protease inhibitor (cystatin), was isolated from taro corm (*Colocasia esculenta*). The result confirmed that recombinant CeCPI protein exhibited strongly cysteine protease inhibitor activity. Thus, the investigation clearly discovered a toxic effect of the plant on the mycelium growth of phytopathogenic fungi [28].

### Anti-inflammatory activity

The ethanolic extracts of the leaves of *C. esculenta* possess anti-inflammatory activity. The study was conducted in Wistar rats using the carrageenan-induced left hind paw oedema, carrageenan-induced pleurisy, and cotton pellet induced granuloma model. The results indicated that the ethanolic extract exhibits significant anti-inflammatory activity when compared with the standard and untreated control [29].

### Uses / Applications

#### Medicinal uses

*C. esculenta* has many medicinal applications as discussed earlier. All the plant parts viz. leaves, stem and tubers shows different medicinal properties. Some of the properties includes antimicrobial, antihepatotoxic, anti-diabetic, anti-lipid peroxidative action, anti-metastatic, antifungal, anti-inflammatory and many more.

People of the Munda tribe, traditionally used corm of taro as a remedy for body ache. The juice extracted from corm of the plant is used in alopecia, as an expectorant, stimulant, appetizer and astringent. When cooked, the vegetable contains mucilage and found to be an effective nervine tonic. Table 2 summarizes chemical constituent and corresponding activity of *C. esculenta* plant parts.

| Plant part | Form                   | Responsible chemical constituent  | Activity  |
|------------|------------------------|---|---|
| Leaves     | Juice                  | β-carotenes, and cryptoxanthin along with vitamin A   | Maintaining healthy mucus membranes, skin and vision.   |
|            |                        |   | Helps to protect from lung and oral cavity cancers [30].  |
|            |                        |   | Good stimulant, expectorant, astringent, appetizer and otalgia.   |
|            | Ethanolic extract      |   | Anti-inflammatory [29].   |
|            | Hydroalcoholic extract |   | Anti-depressant, anxiolytic, sedative, and smooth muscle relaxant activity [31].                                    |
| Corm       | Juice                  | 9, 12, 13-trihydroxy-(E)-10-octadecenoic acid, lipoxygenase and lipid hydroperoxide-converting enzyme | External- alopecia, body ache.  |
|            |                        |   | Internal-laxative, demulcent, anodyne, galactagogue, piles, antidote to the stings of wasps and other insects [19]. |
|            |                        |   | Antifungal  |

|             |                    |  |  |
|-------------|--------------------|--|--|
| Leaf stalks | Juice              | Flavanoids and terpenoids may be responsible.                | Absorbent in cases of inflamed glands and buboes. Anti-diarrhoeal, increases body weight, prevents excessive secretion of sputum in asthmatic individuals. |
|             | Peel decoction     |  | Styptic, otorrhoea, arrests arterial haemorrhage, stimulant, rubefacient.  |
| Petiole     | Pressed juice      |  |  |
| Whole plant | Press extract      | Cyanoglucoside   | Hypoglycemic [32].   |
|             | Ethanollic extract | Monogalactosyl diacylglycerols, digalactosyl diacylglycerols | Hypolipidemic [7,8].   |

**Table 2:** Chemical constituents and activity of different parts of *Colocasia esculenta* plant.

### Pharmaceutical applications

The gum as well as starch obtained from the tuber of the *C. esculenta* plant can be used as an effective binder and mucoadhesive matrix forming agent [33].

Soumya et al. successfully prepared sustained release and evaluated matrix tablets of taro gum using metoprolol succinate as the model drug. The quantity of taro gum (X1) and polyvinylpyrrolidone (PVP) K30 (X2) were selected as independent variables. The time required for 90% of *in vitro* drug release was selected as the dependent variable. Tablets were prepared by direct compression and were evaluated for various post compression parameters such as tablet hardness, friability, weight variation, drug content and *in vitro* dissolution [34].

Another research paper by Arora et al. reported application of Taro gum in development of Mucoadhesive matrix tablets. Matrix tablets of Domperidone as a model drug was compressed using direct compression technique. This study showed concentration dependent mucoadhesive and release retardant potential of taro gum in the formulation of gastro retentive mucoadhesive matrix tablets [35].

Chukwu and Udeala studied the effectiveness of *C. esculenta* gum in poorly compressible drugs-paracetamol and metronidazole tablet formulations. The effectiveness of a polysaccharide gum obtained from the *C. esculenta* was evaluated comparatively with acacia and methylcellulose as binders in the formulation of poorly compressible drugs. At 4% w/w nominal concentration of Colocasia gum in metronidazole tablets and 6% w/w in paracetamol tablets showed very long disintegration time and prolonged release profile. The binders used for comparison yielded tablets that show better *in vitro* release characteristics [36].

*C. esculenta* polysaccharide can be used as a disintegrant in the formulation of orally disintegrating tablets. Its disintegrating property was reported to be comparable with that of the commercially available super-disintegrants.

### Patents filed/published

Table 3 summarises different patents filed /published/granted involving Colocasia plant.

| Title   | Patent Number | Publication date | Inventors           | Description   |
|---|---------------|------------------|---------------------|---|
| Colocasia plant named 'Mojito'                          | USPP21995 P2  | Jun 28, 2011     | Ty Richard Strode   | The present invention relates to a new and distinct cultivar of Colocasia plant, botanically known as <i>Colocasia esculenta</i> .  |
| Colocasia plant named 'Midnight'                        | USPP17887 P2  | Jul 31, 2007     | Richard Strode      |   |
| Colocasia plant named 'MAUI GOLD'                       | USPP24482 P2  | May 20, 2014     | John J. Cho         |   |
| Colocasia plant named 'Noble Gigante'                   | USPP25219 P2  | Jan 6, 2015      | Brian Paul Williams |   |
| Production process of <i>Colocasia esculenta</i> starch | CN103772511 A | May 7, 2014      | -                   | This invention discloses a production process of <i>C. esculenta</i> starch. Process includes peeling, smashing, adjusting pH of the slurry to 8.5-10, performing standing for 50-100 min. This is followed by squeezing, filtering, rinsing filtrate by water, collection of lotion, sieving, centrifugation and drying. |
| Preservation method of <i>Colocasia esculenta</i>       | CN103609665 A | Mar 5, 2014      | -                   | The invention discloses a preservation method of <i>Colocasia esculenta</i> , belonging to the technical field of storage of agricultural products. Carrying out jet impaction and blanching by overheated steam, carrying out jet impaction and micro-drying by gas, and packaging and                                   |

|   |                  |              |               |  |
|---|------------------|--------------|---------------|--|
|   |                  |              |               | preserving by using a freshness protection package. <i>Colocasia esculenta</i> can be preserved for 8 months under the low temperature condition (the temperature is 0-4°C and the relative humidity is 80-95 percent and can be preserved for above 2 months at a normal temperature; no chemical additives are introduced in a processing process, and thus the food safety is high. |
| Herbal formulations and cigarettes containing same useful in controlling body weight. | US20070042055 A1 | Feb 22, 2007 | Palpu et. al. | The present invention relates to herbal formulation comprising a combination of fiber, powder, and/or granules of aerial parts of plants, <i>Colocasia esculenta</i> being one of the components.  |

**Table 3:** List of Patents involving Colocasia plant.

## Conclusion

*C. esculenta* is a largely cultivated plant used from ages as food and medicine. Flavonoids and triterpenoid's are the two major therapeutically active groups of compounds found in the plant. Pharmacologically, the plant is antimicrobial, antihepatotoxic, antidiabetic, anti-lipid peroxidative, antimetastatic, antifungal and anti-inflammatory. There are also many pharmaceutical applications for the plant.

There is a lot of scope for usage of *C. esculenta* in pharmaceutical industries as well as in research work. Herbal formulations consisting of plant extracts from different parts of plants can be developed for many ailments as discussed in this paper. More research work can be done on structural investigation and applications of the gum obtained from *C. esculenta*. The starch and the gum obtained from the tubers can be commercially made use of in pharmaceutical industries in form of binder, matrix forming agent, thickening agent etc. Thus, there is need to explore this plant so as to make use of its medicinal and pharmaceutical properties to its fullest.

## Conflict of Interest

The author(s) declare that there is no conflict of interests regarding publication of this article.

## Acknowledgement

Authors are very much thankful to the Principal, Hyderabad (Sindh) National Collegiate Board's Dr. L. H. Hiranandani college of pharmacy, Ulhasnagar for his continuous support, guidance and encouragement.

## References

- Lin H, Huang AS (1993) Chemical composition and some physical properties of a water-soluble gum in taro (*Colocasia esculenta*). Food Chemistry 48: 403-409.
- Tuse TA, Harle UN, Bore VV (2009) Hepatoprotective activity of *Colocasia antiquorum* against experimentally induced liver injury in rat. Malaysian J Pharma Sci 2: 99-112.
- Lewu MN, Adebola PO, Afolayan AJ (2009) Effect of cooking on the proximate composition of the leaves of some accessions of *Colocasia esculenta* (L.) Schott in KwaZulu-Natal province of South Africa. African J Biotech 8: 1619-1622.
- Awasthi CP, Singh AB (2000) Nutritional quality evaluation of edible leaves of some promising Colocasia and Alocasia collections. Ind J Agric Res 34: 117-121.
- Devarkar VD, Marathe VR, Chavan DP (2011) Dietary and medicinal significance of wild vegetables from Osmanabad region, Maharashtra (India). Life Sciences Leaflets 11: 317-332.
- Phillip BA, Grindleya OF, Asemotaa HN, Errol Y, Morrisona A (2001) Carbohydrate digestion and intestinal ATPases in streptozotocin-induced diabetic rats fed extract of yam (*Dioscorea cayenensis*) or dasheen (*Colocasia esculenta*). Int J Food Sci Nutr 52: 429-433.
- Boban P, Nambisan B, Sudhakaran P (2006) Hypolipidaemic effect of chemically different mucilages in rats: A comparative study. British Journal of Nutrition 96: 1021-1029.
- Tanaka R, Sakano Y, Nagatsu A, Shibuya M, Ebizukab Y, et al. (2005) Synthesis of digalactosyl diacylglycerols and their structure-inhibitory activity on human lanosterol synthase. Bioorg Med Chem Lett 15: 159-162.
- Ravikumar S, Gracelin N, Anitha A, Selvan PG, Kalaiarasi A (2011) *In vitro* antibacterial activity of coastal medicinal plants against isolated bacterial fish pathogens. Int J Pharma Res Devel 3: 109-116.
- Kirtikar KR, Basu BD (2005) Indian Medicinal Plants. Dehradun: International Book Distributors 4: 2615.
- Prajapati P, Kalariya M, Umbarkar R, Parmar S, Sheth N (2011) *Colocasia esculenta*: A potent indigenous plant Int J Nutr Pharmacol Neurol Dis 1: 90-96.
- Iwashina T, Konishi T, Takayama A, Fukada M, Ootani S (1999) Isolation and identification of the flavanoids in the leaves of taro. Ann Tsukuba Botanical Garden 18: 71-74.
- Sheth AK (2005) The Herbs of Ayurveda. AK Sheth Publishers, Ahmedabad, India.
- Noda Y, Kaneyuki T, Mori A, Packer L (2002) Antioxidant activities of pomegranate fruitextract and its anthocyanidins: Delphinidin, cyanidin and pelargonidin. J Agric Food Chem 50: 166-171.
- Cambie RC, Ferguson LR (2003) Potential functional foods in the traditional Maori diet. Mutat Res 523: 109-117.
- Kowalczyk E, Kopff A, Fijalkowski P, Niedworok J, Blaszczyk J, et al. (2003) Effects of anthocyanins on selected biochemical parameters in rats exposed to cadmium. Acta Biochimica Polonica 50: 543-548.
- Anonymous (2005) The Wealth of India: A Dictionary of Indian Raw Materials and Industrial Products. NISCSIR, New Delhi, India pp: 1-225.
- Khare CP (2007) Indian Medicinal Plants. Springer, New Delhi, India.
- Anonymous (2001) The Wealth of India: A Dictionary of Indian Raw Materials and Industrial Products. NISCSIR New Delhi, India, PP: 2-157.
- Huang AS, Tanudjaja L (1992) Application of anion-exchange high-performance liquid chromatography in determining oxalates in taro (*Colocasia esculenta*) corms. J Agric Food Chem 40: 2123-2126.

21. Nakayama TO, Harvey TJ, Tsung HCK (2006) Anthocyanin composition of taro. J Food Sci 42: 19-21.
22. Singh B, Namrata LK, Dwivedi SC (2011) Antibacterial and antifungal activity of *Colocasia Esculenta* aqueous extract: An edible plant. Journal of Pharmacy Research 4: 1459-1460.
23. Bhagyashree RP, Hussein MA (2011) Antihepatotoxic activity of *Colocasia esculenta* leaf Juice. International Journal of Advanced Biotechnology and Research 2: 296-304.
24. Patel DK, Kumar R, Laloo D, Hemalatha S (2012) Diabetes mellitus: An overview on its pharmacological aspects and reported medicinal plants having antidiabetic activity. Asian Pac J Trop Biomed 2: 411-420. 25.
25. Kumawat NS, Chaudhari SP, Wani NS, Deshmukh TA, Patil VR (2010) Antidiabetic activity of ethanol extract of *Colocasia esculenta* leaves in alloxan induced diabetic rats. International Journal of Pharm Tech Research 2: 1246-1249.
26. Bhagyashree RP, Hussein M (2011) Anti-Lipid Peroxidative Activity of *Colocasia esculenta* leaf Juice against CCl<sub>4</sub> and Acetaminophen Mediated Cell Damage. International Journal of Pharmaceutical Applications 2: 141-149.
27. Kundu N, Campbell P, Hampton B, Lin CY, Ma X, et al. (2012) Antimetastatic activity isolated from *Colocasia esculenta* (taro). Anticancer Drugs 23: 200-211.
28. Yang AH, Yeh KW (2005) Molecular cloning, recombinant gene expression and antifungal activity of cystain from taro. Planta Med 22: 333-341.
29. Shah BN, Nayak BS, Bhatt SP, Jalalpure SS, Sheth AK (2007) The anti-inflammatory activity of the leaves of *Colocasia esculenta*. Sau Pharm J 15: 3-4.
30. Kirtikar KR, Basu BD (2005) Indian medicinal plants. Dehradun: International Book Distributors 4: 2615.
31. Kalariya M, Parmar S, Sheth N (2010) Neuropharmacological activity of hydroalcoholic extract of leaves of *Colocasia esculenta*. Pharm Biol 48: 1207-1212.
32. Grindley BA, Omoruyi F, Asemota HN, Morrison A (2002) Carbohydrate digestion and intestinal ATPases in streptozotocin-induced diabetic rats fed extract of yam (*Dioscorea cayenensis*) or dasheen (*Colocasia esculenta*). Nut Res 22: 333-41.
33. Pawar HA, Kamat SR, Choudhary PD (2015) An Overview of Natural Polysaccharides as Biological Macromolecules: Their Chemical Modifications and Pharmaceutical Applications. Biol Med (Aligarh) 6: 224.
34. Soumya M, Chowdary YA, Swapna VN, Prathyusha ND, Geethika R, et al. (2014) Preparation and optimization of sustained release matrix tablets of metoprolol succinate and taro gum using response surface methodology. Asian J Pharm 8: 51-57.
35. Gurpreet A, Karan M, Inderbir S (2011) Formulation and evaluation of mucoadhesive matrix tablets of Taro gum: Optimization using response surface methodology. Polimery Medycynie 41: 23-34.
36. Chukwu KI, Udeala OK (2000) Binding effectiveness of *Colocasia esculenta* gum in poorly compressible drugs-paracetamol and metronidazole tablet formulations. Boll Chim Farm 139: 89-97.