

Bilateral Impact between Dairy Cattle and Global Warming

Junichi Takahashi*

Department of Animal and Food Hygiene, Obihiro University of Agriculture and Veterinary Medicine, Obihiro, Japan

Although water vapour is the largest and essential greenhouse gas inclusive of ozone in the stratosphere, the mitigation of anthropogenic six greenhouse gases such as carbon dioxide, methane, nitrous oxide, sulphur hexafluoride, hydrofluorocarbons and perfluorocarbons have been established as legally binding commitments in The Kyoto Protocol [1]. Important greenhouse gases attributed to ruminant livestock are methane and nitrous oxide. When the greenhouse effect of carbon dioxide is set to 1, the impact of the greenhouse effect of methane and nitrous oxide is 21-fold and 310-fold higher than carbon dioxide respectively. Rumen fermentation of ruminant livestock like dairy cattle and anaerobic fermentation of organic waste including animal manures are major contributors of methane emission as anthropogenic sources [2]. The total contribution of belching methane derived from rumen fermentation and the manure to global methane emission might be accounted for nearly 18-20%. Thus, the development of mitigation methods of rumen methane is the most significant issue in the world ruminant livestock production. Although ionophores as feed additives such as monensin and salinomycin have a potential to abate rumen methane production, the efficacy may still continue in the manure, *i.e.*, uncontrolled antibiotics will be spread over the soil environment [3]. Therefore, manipulators to abate rumen methanogenesis must be safe to the environment, not to mention human and livestock. The prompt increase of atmospheric nitrous oxide since last century is closely related to abrupt expansion of human and animal population after an innovation of Haber-Bosch process. Severe environmental pollutions were caused at the same time though the reactive nitrogen withdrawn from atmosphere as stable paired nitrogen brought about prosperous food production. Anthropogenic increases in these greenhouse gases are not immune to continuous expansion of the chemical nitrogen fertilizer consumption [4]. Accurate assessment of the impact of inorganic and organic nitrogen

on greenhouse effect may be useful to make up an inventory and the mitigation strategies.

Recent climate change has amplified a risk to expose dairy cattle to hot environment [5]. In consequence, milk production and reproductive proficiency of dairy cows suffer from heat stress due to their strong stress sensitivity [6]. Heat stress might induce oxidative stress on the animals as an external inducer [7].

To alleviate these stresses on dairy cow the establishment of mitigation strategies of greenhouse gases is an urgent issue.

References

1. IPCC (1996) Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. IPCC-NGGIP.
2. Moss AR (1993) Methane: Global Warming and Production by Animals. Canterbury. Chalcombe Publications, England.105.
3. Mwenya B, Sar C, Pen B, Morikawa R, Takaura K, et al. (2006) Effect of feed additives on ruminal methanogenesis and anaerobic fermentation of manure in cows and steers. In: Soliva C, Takahashi J, Kreuzer M (eds), Greenhouse Gases on Animal Agriculture update. International Congress Series. Elsevier, Amsterdam. 1293: 209-212.
4. Takahashi J (2006) Sustainable manure management in the nitrogen recycling system FLRC/NZFMRA Conference on Implementing Sustainable Nutrient management Strategies in Agriculture, Palmerstone North, New Zealand.
5. Nidumolu U, Crimp S, Gobbett D, Laing A, Howden M, et al. (2013) Spatio-temporal modelling of heat stress and climate change implications for the Murray dairy region, Australia. Int J Biometeorol.
6. West JW (2003) Effects of heat-stress on production in dairy cattle. J Dairy Sci 86: 2131-2144.
7. Bernabucci U, Ronchi B, Lacetera N, Nardone A (2002) Markers of oxidative status in plasma and erythrocytes of transition dairy cows during hot season. J Dairy Sci 85: 2173-2179.

*Corresponding author: Junichi Takahashi, GGAA president, Professor emeritus, Department of Animal and Food Hygiene, Obihiro University of Agriculture and Veterinary Medicine, Obihiro, Japan, Tel: +81 9082742341; E-mail: junichi@obihiro.ac.jp, takjekf@gmail.com

Received November 26, 2013; Accepted November 28, 2013; Published November 30, 2013

Citation: Takahashi J (2013) Bilateral Impact between Dairy Cattle and Global Warming. Adv Dairy Res 2: e104. doi: [10.4172/2329-888X.1000e104](https://doi.org/10.4172/2329-888X.1000e104)

Copyright: © 2013 Takahashi J. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.