

Copper Oxidation State and Mycobacterial Infection

Marc Soliz^{*}

Laboratory of Biochemistry and Molecular Biology, Tomsk State University, Prospect Lenina, Tomsk, Russia

^{*}Corresponding author: Soliz M, Laboratory of Biochemistry and Molecular Biology, Tomsk State University, Prospect Lenina 36, 634050 Tomsk, Russia, Tel: 923 403 5002; E-mail: marc@soliz-scientific.ch

Rec date: March 28, 2016; Acc date: May 20, 2016; Pub date: May 27, 2016

Copyright: © 2016 Soliz M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Copper in aerobic environments prevails as cupric ions, Cu^{2+} , but inside cells, is usually present as cuprous ions, Cu^+ . Cu^+ is considerably more toxic to bacteria than Cu^{2+} . Phagosomes employ, among other mechanisms, Cu^+ to create a hostile environment for engulfed bacteria. Bacteria can reduce or oxidize copper by various mechanisms. How such reactions could affect bacterial survival in phagosomes is discussed, with a focus on *Mycobacteria*.

Keywords: Copper reduction; Multicopper oxidase; Copper oxidation; Mycobacterial infection

Introduction

Copper is an essential trace element for most living organisms due to its role as a cofactor in many enzymes. However, excess copper is toxic to cells. It causes oxidative stress by participating in Fenton-type reactions which lead to the generation of reactive oxygen species, or copper can damage [Fe-S]-cluster-proteins by displacing the iron [1]. Therefore, all cells have elaborated sophisticated systems for copper homeostasis.

Bacteria are particularly vulnerable to copper toxicity, with lethal copper concentrations ranging from μM to low mM, but strongly depending on the experimental setting [2]. Consequently, copper salts are widely used for pest control in agriculture. And as a corollary to this, copper ions participate in the killing of engulfed bacteria by phagosomes. The recent description of non-enzymatic copper reduction by *Lactococcus lactis* bears on several of these issues and moves the oxidation state of copper in the focus of bactericidal mechanisms [3].

Non-enzymatic copper reduction by bacteria

Under aerobic conditions, copper in solution is present as cupric ions, Cu^{2+} . Already forty years ago, it was recognized that in *E. coli* cultures, Cu^{2+} is reduced to Cu^+ and that the cuprous ions are considerably more toxic to cells than Cu^{2+} [4]. In aerobic cultures, Cu^+ is steadily re-oxidized by oxygen to the less toxic Cu^{2+} , whereas in anaerobic cultures, Cu^+ accumulates, creating a much more toxic environment for bacteria. More recently, the copper reducing activity of *E. coli* has been molecularly characterized. It was shown that approximately 70% of this activity was due to menaquinones, 10% to copper reduction by the NADH dehydrogenase, NDH^{-2} , and the remainder due to unknown sources [5-7].

The Gram-positive bacteria *Lactococcus lactis* have also been reported to exhibit extracellular Cu^{2+} -reductase activity [8,9]. It was shown that menaquinones in the bacterial membrane can catalyze copper reduction in a non-enzymatic reaction, presumably by direct interaction of copper with the membrane [3]. Lactic acid bacteria and

other fermenting organisms have an excess of reducing equivalents when growing fermentatively and will take advantage of any opportunity to dispose of these. In fermentatively growing *L. lactis*, sub-lethal concentrations of Cu^{2+} ions in the media can apparently serve as electron acceptors [3].

Short-chain menaquinones present in the membrane are reduced by NADH via membrane-bound NADH oxidases, NoxA or NoxB; the reduced menaquinones in turn reduce extracellular Cu^{2+} to Cu^+ (Figure 1A). Since electrons move from a standard redox potential of -315 mV for the NADH/NAD⁺ couple to one of +150 mV for the $\text{Cu}^{2+}/\text{Cu}^+$ couple, the reaction is thermodynamically very favorable. As expected, a mutant unable to synthesize menaquinones due a deficiency in O-succinylbenzoic acid CoA ligase (MenE) does not reduce extracellular copper and is therefore also less copper sensitive than the wild-type because it does not produce the more toxic Cu^+ species [3].

The copper reduction by *L. lactis* constitutes a type of metal-based respiration, which is a well-known phenomenon and has been studied extensively in many organisms [10]. In fact, iron respiration was proposed to be one of the earliest forms of respiration that evolved on the Earth [11]. Non-enzymatic copper reduction by *L. lactis* is a 'balancing act' for the bacteria: at copper concentrations which do not inhibit growth, Cu^{2+} reduction can stimulate growth by serving as an electron sink; however, this leads to the accumulation of more toxic Cu^+ , which could kill the organism.

Respiration competes with non-enzymatic copper reduction

L. lactis is classified as facultative anaerobe, meaning that it can tolerate oxygen, but cannot respire it. In 1970, Sijpesteijn [12] discovered that in growth media supplemented with heme, *L. lactis* consumes oxygen and grows to much higher cell density. It turned out that *L. lactis* possess functional genes for the two subunits of a terminal cytochrome *bd* oxidase [ubi (mena-) quinol: O₂ oxidoreductase], but cannot synthesize heme.

If heme is supplied to cultures, a functional oxidase is synthesized and respiration ensues. In addition, *L. lactis* is endowed with a ferrochelatase, which converts protoporphyrin IX to heme, so aerobic growth is also possible on protoporphyrinogen IX [13]. The ability of

heme to induce respiration in bacteria classified as "anaerobic" is most likely shared by many firmicutes [14].




Figure 1A: Electron flow in *L. lactis* under fermentative growth conditions. Under fermentative growth conditions in the presence of Cu^{2+} , the membrane-bound NADH oxidase, NoxA/B, convert oxidized menaquinones (MQ) to their reduced form (MQH_2), with NADH serving as reductant. The reduced menaquinones in turn reduce extracellular Cu^{2+} in a non-enzymatic reaction. The respective standard redox potentials, E° , are given in the figure and the preferred overall electron flow is indicated by the red arrow and that of partial reactions by black arrows.




Figure 1B: Electron flow in *L. lactis* under respiratory growth conditions. Under respiring growth conditions in the presence of heme, a terminal cytochrome *bd* oxidase is synthesized. Menaquinones reduced by NADH/NoxA/B are rapidly oxidized by the oxidase, using O_2 as an electron acceptor. The respective standard redox potentials, E° , are given in the figure and the preferred overall electron flow is indicated by the green arrow and that of partial reactions by black arrows.

Surprisingly, when *L. lactis* was supplied with heme, copper reduction was strongly reduced [3]. This could be explained by competition for electrons between the two redox reactions, copper reduction and oxygen reduction. The difference in standard redox potential between $\text{O}_2/\text{H}_2\text{O}$ and $\text{NADH}/\text{NADH}^+$ is $+815 \text{ mV} - (-315$

$\text{mV}) = +1130 \text{ mV}$, compared to that of the redox couple $\text{Cu}^{2+}/\text{Cu}^+$ and $\text{NADH}/\text{NADH}^+$, which is $+150 \text{ mV} - (-315 \text{ mV}) = +465 \text{ mV}$ (Figure 1B).

Oxygen reduction is thus the thermodynamically favored over copper reduction in respiring cells. Reduction of copper by menaquinones is most likely a fortuitous mechanism. First, copper is usually not present in sufficient concentrations in the natural habitats of *L. lactis* to make a significant contribution to metal respiration. Secondly, it produces toxic Cu^+ ions which can interfere with growth. When growing in milk, *L. lactis* substantially lowers the redox potential of this substrate from $+300 \text{ mV}$ to -220 mV , presumably as a way of disposing of excess reducing equivalents [15]. This could be of great importance in fermentative processes, but the electron acceptors in milk are unknown and the process remains largely unexplored.

Infection raises phagosomal copper

It has long been known that copper deficiency sensitizes mammalian hosts to microbial infections [16] and mechanistic insight into at least one such mechanism is beginning to emerge. The first line of defense against infection is often phagocytic immune cells. Upon activation of these cells, the phagosome adopts an anti-microbial environment by assuming an acidic pH and elevating reactive oxygen species, nitric oxide, hypochlorite (ClO^-), and proteases [17].

Following infection by *Mycobacteria* and other pathogens, iron and copper also accumulate in the phagosome [18,19]. The mechanism of phagosomal copper accumulation is quite well understood (for review of normal eukaryotic copper homeostasis, see [20]). Ctr1 is the universal copper importer in the plasma membrane of eukaryotic cells. White et al. [21] showed that in cultured macrophage-like RAW264.7 cells, killing of *E. coli* is stimulated by the addition of copper to the culture media, which leads to up regulation of Ctr1 and enhanced transport of copper into the cytoplasm (Figure 2).

There, copper is sequestered by the 7 kDa protein Atox1, which functions as a copper shuttle, or chaperone. Atox1 delivers copper to the copper transporting ATPase, ATP7A, which is located in the trans-Golgi network under basal conditions to deliver copper to secreted cuproenzymes like lysyl oxidase or ceruloplasmin. In activated macrophages, ATP7A traffics from the trans-Golgi network to phagosomes.

Apparently, Ctr1 and ATP7A fulfill cell defense functions by raising cytoplasmic copper levels and transporting copper into phagosomes that contain engulfed bacteria. Such a copper-up response may not be limited to macrophage, but occur in a range of cell types in response to internalized bacteria or other factors such as inflammation [20,22].

Mycobacterial defense against copper

Bacteria are well equipped to combat excessive environmental copper and respond by the upregulation of a number of genes which function in the defense against copper. *Mycobacteria*, including the human pathogen *Mycobacterium tuberculosis*, are unique among Gram-positive bacteria in featuring a complex cell wall that contains unusual lipids and thereby forms a kind of outer membrane [23].

This outer membrane functions as a barrier to chemotherapeutic agents and metal ions. A point of weakness are the porins that are required as entry point for essential nutrients, including copper, but may also allow excess copper to get access to the cytoplasmic membrane (Figure 2) [24].




Figure 2: Copper homeostasis in a macrophage with engulfed *M. tuberculosis*.

Cu homeostasis on the host side proceeds by the components indicated in blue: the high-affinity copper transporter of the plasma membrane, Ctr1, is induced by the activation of the macrophage and transports extra copper into the cytoplasm; Atox1, a cytosolic Cu⁺ chaperone, transfers the copper to ATP7A, which has trafficked from the trans-Golgi network to the phagosome and pumps copper into the phagosomal lumen. On the pathogen side, *M. tuberculosis* possesses several lines of defense against copper, indicated in red. Copper crosses the cell wall/outer membrane via porins. Excess cytoplasmic copper induces CtpV, a copper export ATPase which is under the control of the CsoR repressor. MctB functions in expelling copper, which has been pumped into the periplasmic space by CtpV, from the cell. Also induced under copper stress are a periplasmic multicopper oxidase, MmcO, which oxidizes Cu⁺ to Cu²⁺, and MymT, a metallothionein which sequesters excess cytoplasmic copper. These proteins are under the control of a second copper-inducible repressor, RicR. Cu⁺ is depicted as green dots and Cu²⁺ as blue dots. See text for additional explanations.

In macrophages, this will induce the *cso* operon, which is under the control of the copper-inducible CsoR repressor [25]. The operon encodes CtpV, which is an enzyme related to known bacterial copper ATPases and pumps cytoplasmic copper to the periplasmic space, and two proteins of unknown function, Rv0970 and Rv0968. MctB is required to transport the periplasmic copper to the cell exterior and is thus functionally, but not structurally, similar to the Cus-system of Gram-negative bacteria [26]. Mutation of the *metB* gene resulted in a dramatically reduced bacterial burden in lungs and lymph nodes in guinea pigs infected with *M. tuberculosis*, supporting the important role of copper in the phagosome [27]. RicR, a second copper-inducible repressor, controls a regulon of six monocistronic gene, namely RicR itself, MymT, a metallothionein which sequesters cytoplasmic copper, MmcO, a multicopper oxidase, and three proteins of unknown function, LpqS, Rv2963, and SocAB [28].

Of special interest is MmcO, a CueO-type multicopper oxidase that has been well characterized in *E. coli* and other bacteria [29]. CueO-type multicopper oxidases are common in Gram-negative bacteria and can oxidize Cu⁺, Fe²⁺, and various polycyclic aromatic hydrocarbons. MmcO has been shown to be required for copper resistance by *M. tuberculosis*, presumably by reducing Cu⁺ to less toxic Cu²⁺ [30]. A

CueO-type multicopper oxidase may not be of equal importance for all species; for *Brucella melitensis* it was found that the multicopper oxidase did not affect survival in macrophages [31], while mutation of *cueO* in *Salmonella Typhimurium* attenuated survival in liver and spleen, but not macrophages [32]. Clearly, the CueO-type multicopper oxidase contributes to bacterial copper resistance in culture, but it may not be a key factor in all species for survival in phagosomes [33]. This may have multiple underlying reasons. For one, the phagosomal interior may be sufficiently oxidizing for spontaneous copper oxidation to occur. Secondly, the primary role of CueO-like enzymes *in vivo* may not always be the oxidation of copper, but oxidation of iron or organic substrates. Yet, two general observations underline the importance of bacterial copper resistance for survival in macrophages: bacterial mutants which are copper-sensitive in culture are hyper sensitized to macrophage killing and, conversely, chelation of macrophage copper results in increased survival of engulfed organisms [34].

Redox state of copper in phagosomes and Mycobacterial infections

Copper transported into phagosomes by ATP7A is in the reduced Cu⁺ form, which is considerably more toxic to bacteria than Cu²⁺ [3]. Reactive oxygen species in the phagosomal compartment could lead to the generation of significant Cu²⁺ by spontaneous oxidation. This Cu²⁺ could, at least in principle, be re-reduced by non-enzymatic copper reduction via menaquinones in the membrane of *Mycobacteria*, restoring the more toxic Cu⁺ pool. However, this reaction is unlikely to occur because *Mycobacteria* harbor two terminal oxidases, a cytochrome *aa₃* oxidase and a cytochrome *bd* oxidase [35], which would effectively compete with non-enzymatic copper reduction.

But another mechanism may be very significant: compared to wild-type, bacteria deficient in CueO-type multicopper oxidase exhibit much poorer survival in various eukaryotic cell types. This suggests that the oxidation of phagosomal Cu⁺ to less toxic Cu²⁺ by multicopper oxidase can partially detoxify the phagosomal lumen and aid the survival of the pathogen. MmcO of *M. tuberculosis* could catalyze such a detoxification reaction. Purified CueO multicopper oxidases of *E. coli* and *Desulfosporosinus* sp. OT were shown to be inhibited by Ag⁺ with an I₅₀ of 1 μM ([36] and unpublished observations). This contrasts with the inhibition of bacterial growth, which requires 10- to 1000-fold higher silver concentrations, depending on the species and the media [37]. Low silver ion concentrations could thus potentiate the antibacterial milieu of phagosomes by preventing the oxidation of Cu⁺ to less toxic Cu²⁺ by bacterial multicopper oxidases. Silver in various formulations has been used for centuries to treat venereal diseases and other bacterial infections and to dress open wounds and burns [38]. The latter applications continue to be in use up to this day. How silver acts in these applications remains unknown. One possibility is the inhibition of bacterial CueO-type multicopper oxidases by silver, thereby compromising survival in phagosomes. Such a mechanism would certainly deserve further investigation in model systems.

Recent progress in the understanding of both, eukaryotic and prokaryotic copper homeostasis, have greatly advanced our insight into copper's role in phagosomal killing of bacteria. The much higher bactericidal activity of Cu⁺ versus Cu²⁺ and thus copper redox reactions appear to play an important role in the process, but addressing this question remains difficult. Although intracellular copper indicators specific for Cu⁺ or Cu²⁺ have become available in recent years, their targeting to specific cell compartments is still a

challenge. Clearly, the role of copper redox reactions in phagosomes requires further investigation, but its comprehension may eventually open new avenues to drug development.

Acknowledgment

The author is supported by the Russian Federation Ministry of Education and Science, contract number 11.519.11.2004.

References

- Macomber L, Imlay JA (2009) The iron-sulfur clusters of dehydratases are primary intracellular targets of copper toxicity. *Proc Natl Acad Sci USA* 106: 8344-8349.
- Magnani D, Solioz M (2007) How bacteria handle copper. *Microbiology Monographs* 6: 259-285.
- Abicht HK, Gonskikh Y, Gerber SD, Solioz M (2013) Non-enzymic copper reduction by menaquinone enhances copper toxicity in *Lactococcus lactis* IL1403. *Microbiology* 159: 1190-1197.
- Beswick PH, Hall GH, Hook AJ, Little K, McBrien DC, et al. (1976) Copper toxicity: evidence for the conversion of cupric to cuprous copper in vivo under anaerobic conditions. *Chem Biol Interact* 14: 347-356.
- Volentini SI, Fariás RN, Rodríguez-Montelongo L, Rapisarda VA (2011) Cu(II)-reduction by *Escherichia coli* cells is dependent on respiratory chain components. *Biometals* 24: 827-835.
- Rapisarda VA, Chehín RN, De Las Rivas J, Rodríguez-Montelongo L, Fariás RN, et al. (2002) Evidence for Cu(I)-thiolate ligation and prediction of a putative copper-binding site in the *Escherichia coli* NADH dehydrogenase-2. *Arch Biochem Biophys* 405: 87-94.
- Rapisarda VA, Montelongo LR, Fariás RN, Massa EM (1999) Characterization of an NADH-linked cupric reductase activity from the *Escherichia coli* respiratory chain. *Arch Biochem Biophys* 370: 143-150.
- Wunderli-Ye H, Solioz M (1999) Effects of promoter mutations on the in vivo regulation of the cop operon of *Enterococcus hirae* by copper(I) and copper(II). *Biochem Biophys Res Commun* 259: 443-449.
- Rezaiki L, Lamberet G, Derre A, Gruss A, Gaudu P (2008) *Lactococcus lactis* produces short-chain quinones that cross-feed Group B *Streptococcus* to activate respiration growth. *Mol Microbiol* 67: 947-957.
- Lovley DR (2002) Dissimilatory metal reduction: from early life to bioremediation. *ASM* 68: 231-237.
- Vargas M, Kashefi K, Blunt-Harris EL, Lovley DR (1998) Microbiological evidence for Fe(III) reduction on early Earth. *Nature* 395: 65-67.
- Sijpesteijn AK (1970) Induction of cytochrome formation and stimulation of oxidative dissimilation by hemin in *Streptococcus lactis* and *Leuconostoc mesenteroides*. *Antonie Van Leeuwenhoek* 36: 335-348.
- Duwat P, Sourice S, Cesselin B, Lamberet G, Vido K, et al. (2001) Respiration capacity of the fermenting bacterium *Lactococcus lactis* and its positive effects on growth and survival. *J Bacteriol* 183: 4509-4516.
- Gaudu P, Vido K, Cesselin B, Kulakauskas S, Tremblay J, et al. (2002) Respiration capacity and consequences in *Lactococcus lactis*. *Antonie Van Leeuwenhoek* 82: 263-269.
- Tachon S, Brandsma JB, Yvon M (2010) NoxE NADH oxidase and the electron transport chain are responsible for the ability of *Lactococcus lactis* to decrease the redox potential of milk. *Appl Environ Microbiol* 76: 1311-1319.
- Muñoz C, Rios E, Olivos J, Brunser O, Olivares M (2007) Iron, copper and immunocompetence. *Br J Nutr* 98 Suppl 1: S24-28.
- Festa RA, Thiele DJ (2012) Copper at the front line of the host-pathogen battle. *PLoS Pathog* 8: e1002887.
- Wagner D, Maser J, Moric I, Vogt S, Kern WV, et al. (2006) Elemental analysis of the *Mycobacterium avium* phagosome in Balb/c mouse macrophages. *Biochem Biophys Res Commun* 344: 1346-1351.
- Hodgkinson V, Petris MJ (2012) Copper homeostasis at the host-pathogen interface. *J Biol Chem* 287: 13549-13555.
- Wang Y, Hodgkinson V, Zhu S, Weisman GA, Petris MJ (2011) Advances in the understanding of mammalian copper transporters. *Adv Nutr* 2: 129-137.
- White C, Lee J, Kambe T, Fritsche K, Petris MJ (2009) A role for the ATP7A copper-transporting ATPase in macrophage bactericidal activity. *J Biol Chem* 284: 33949-33956.
- Hasan NM, Lutsenko S (2012) Regulation of copper transporters in human cells. *Curr Top Membr* 69: 137-161.
- Bansal-Mutalik R, Nikaido H (2014) Mycobacterial outer membrane is a lipid bilayer and the inner membrane is unusually rich in diacyl phosphatidylinositol dimannosides. *Proc Natl Acad Sci USA* 111: 4958-4963.
- Speer A, Rowland JL, Haeili M, Niederweis M, Wolschendorf F (2013) Porins increase copper susceptibility of *Mycobacterium tuberculosis*. *J Bacteriol* 195: 5133-5140.
- Ward SK, Hoyer EA, Talaat AM (2008) The global responses of *Mycobacterium tuberculosis* to physiological levels of copper. *J Bacteriol* 190: 2939-2946.
- Nies DH (2003) Efflux-mediated heavy metal resistance in prokaryotes. *FEMS Microbiol Rev* 27: 313-339.
- Wolschendorf F, Ackart D, Shrestha TB, Hascall-Dove L, Nolan S, et al. (2011) Copper resistance is essential for virulence of *Mycobacterium tuberculosis*. *Proc Natl Acad Sci USA* 108: 1621-1626.
- Darwin KH (2015) *Mycobacterium tuberculosis* and Copper: A Newly Appreciated Defense against an Old Foe? *J Biol Chem* 290: 18962-18966.
- Singh SK, Grass G, Rensing C, Montfort WR (2004) Cuprous oxidase activity of CueO from *Escherichia coli*. *J Bacteriol* 186: 7815-7817.
- Rowland JL, Niederweis M (2013) A multicopper oxidase is required for copper resistance in *Mycobacterium tuberculosis*. *J Bacteriol* 195: 3724-3733.
- Wu T, Wang S, Wang Z, Peng X, Lu Y, et al. (2015) A multicopper oxidase contributes to the copper tolerance of *Brucella melitensis* 16M. *FEMS Microbiol Lett* 362: fmv078.
- Achard ME, Tree JJ, Holden JA, Simpfendorfer KR, Wijburg OL, et al. (2010) The multi-copper oxidase CueO of *Salmonella enterica* serovar Typhimurium is required for systemic virulence. *Infect Immun* 78: 2312-2319.
- Osman D, Cavet JS (2011) Metal sensing in *Salmonella*: implications for pathogenesis. *Adv Microb Physiol* 58: 175-232.
- Achard ME, Stafford SL, Bokil NJ, Chartres J, Bernhardt PV, et al. (2012) Copper redistribution in murine macrophages in response to *Salmonella* infection. *Biochem J* 444: 51-57.
- Cook GM, Hards K, Vilchère C, Hartman T, Berney M (2014) Energetics of Respiration and Oxidative Phosphorylation in *Mycobacteria*. *Microbiol Spectr* 2.
- Singh SK, Roberts SA, McDevitt SF, Weichsel A, Wildner GF, et al. (2011) Crystal structures of multicopper oxidase CueO bound to copper(I) and silver(I): functional role of a methionine-rich sequence. *J Biol Chem* 286: 37849-37857.
- Silvestry-Rodríguez N, Sicairos-Ruelas EE, Gerba CP, Bright KR (2007) Silver as a disinfectant. *Rev Environ Contam Toxicol* 191: 23-45.
- Castellano JJ, Shafii SM, Ko F, Donate G, Wright TE, et al. (2007) Comparative evaluation of silver-containing antimicrobial dressings and drugs. *Int Wound J* 4: 114-122.