

Cytokines Network and Influenza Virus Infection

Chengcai Lai, Xiliang Wang and Penghui Yang*

302 Military Hospital, Beijing Institute of Microbiology and Epidemiology, Beijing, China

Abstract

Influenza virus is the common trigger for cytokines. Excessive cytokine response can lead to different extent pulmonary injury while defending pathogens, which depends on the pathogenicity of the virus and the strength of host response. Insight of the host reaction and the kinetics of cytokines network response can guide the clinical diagnosis and treatment, which can help reduce the morbidity and the mortality.

Keywords: Cytokine; Influenza virus; Lung injury

Cytokines play an important role in innate immune system recognition and rapid clearance of pathogens in mammalian at the cost of causing tissues injury, which also can regulate the innate immunity and adapt immunity [1]. Adequate cytokine activation enhances the ability of defending pathogens. However, excessive activation of the cytokine responses may destroy the homeostasis and lead to pathological inflammatory consequences [2].

An inflammatory response flaring out of control is known as "cytokine storm". The term of "cytokine storm" was first coined by Ferrara JL to describe their observations in graft-versus-host disease (GVHD) [3], after which it began to appear more frequently in the scientific researches about acute lung injury induced by the influenza A viruses. Previous researches show that influenza A viruses were the most common pathogens during respiratory infection which can induce different cytokines profiles in lung tissues and have a significant influence on patients' prognosis [4]. Different subtypes of influenza A viruses can raise various network of cytokines, even in the same species.

The cytokines can act as pro-inflammatory or anti-inflammatory in the host response, and the advent of pro-inflammatory cytokines usually accompanied with consistent presence of anti-inflammatory cytokines [5]. Once the balance between anti- and pro-inflammatory response was broken, it would have irreparable consequences.

The cytokines including IL-1 β , IL-1Ra, IL-4, IL-5, IL-6, IL-7, IL-8, IL-10, IL-13, IL-17, G-CSF, GM-CSF, IFN- α , IFN- β , IFN- γ , TNF- α , IP-10, MCP-1, MIP-1 α , MIP-1 β , RANTES, keratinocyte-derived chemokine (KC) [6-9], which were reported to aroused to some extent in the serum samples of patients infected with influenza A virus, and most of which can recruit macrophages, neutrophils and other polymorphonuclear cells to inflammatory sites [2,10]. Previous studies showed that pro-inflammatory cytokines TNF- α , IL-6, IP-10 are excessively elevated in infections caused by highly virulent subtypes such as H5N1,2009 pandemic influenza A virus (H1N1) [11,12], which are consistent with findings in SARS [8,13], compared to the seasonal influenza. On 31 March 2013, an explosive human infection with a novel reassortant avian influenza H7N9 virus has recently been reported in China. Researchers also grasped the significant increased levels of IL-6, IP-10, TNF- α and MCP-1 in the serum gathered from H7N9 patients [14]. Similarly, significant increase of IL-6 and TNF- α was probed in the H1N1 with bacterial co-infection group [15], indicating that TNF- α , IL-6, IP-10 may be the key contributors in virus mediated severe respiratory diseases, which would lead to Acute Lung Injury (ALI) or Acute Respiratory Distress Syndrome (ARDS) and add the morbidity and mortality. On the other hand, anti-inflammatory cytokine appears somewhat later as the body attempts to control the

acute systemic inflammatory response. For example, IL-10 is a cytokine with pleiotropic effects in immunoregulation and inflammation, which can downregulate the expression of Th1 cytokine and can recruit fibrocytes into the lung [16,17].

The expression of cytokine is associated with the condition of the patients. The IL-6 is lower in small-for-gestational-age (SGA) neonates / appropriate-for-gestational-age (AGA) neonates/children than in the adults [18,19]. Additionally, the pessimistic emotions were correlated with lower Th1 cytokine (IL-2, IFN- γ) expression during virus stimulation, whereas optimism contributes to greater cytokine responses. It appeared to be more common than those between mood and the Th2 cytokine response [19]. Furthermore, compared to non-asthmatic patients, asthmatic patients also show higher level of IL-5, histamine, protease during virus infection, which are allergic [20,21].

The amplified situation of cytokine expression was proved by multiple researches, but the mechanisms remain unclear. It is known that influenza virus may escape the cytokines' antiviral effects and consistent replicate [22], the increased viral load may mount host cytokine response, such as IL-1Ra, IL-6, MCP-1, TNF- α and IP-10 [6, 23], and cause hypercytokinemia. Moreover, as the cytokines secreted, it appears to have secondary stimulation. For example, when treated with TNF- α , differentiated astrocytes can express IL-6 as well as other cytokine; it is not the case in turn [24]. IFN- γ can stimulate the delivery of IP-10 and MIG, and they can modulate the Th1 cellular immune response in influenza infection [25,26]. Likewise, IFN- α , IFN- γ , IL-1 β can affect the secretion of IL-6 and IL-12 to various degrees as well [27-29].

There are many complete researches about the relationship between cytokines, and some cytokine regulators are already licensed for human use, such as IL-2 and interferon (IFNs) [30]. Nevertheless, it is limited in guidance on the control of the cytokine storm, and the threaten caused by influenza A virus still exist. Base on that, scientists turn to study the effect of different cell types on initiation and amplification of the cytokine storm that follow virulent influenza infection. It shows

*Corresponding author: Penghui Yang, Beijing Institute of Microbiology and Epidemiology, 302 Military Hospital, Beijing, 100039, China, Tel: +86-010-66933336; E-mail: ypenghuiamms@hotmail.com

Received April 05, 2014; Accepted April 28, 2014; Published May 02, 2014

Citation: Lai C, Wang X, Yang P (2014) Cytokines Network and Influenza Virus Infection. Clin Microbiol 3: 147. doi:10.4172/2327-5073.1000147

Copyright: © 2014 Lai C, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

that diversified types of cells have capacities for secreting cytokines, including macrophages, neutrophils, though fibrocytes and endothelial cells [5]. Previous studies show that the sphingosine-1-phosphate (S1P) signaling system acted as immune response modulator is related to cytokine amplification during influenza virus infection [31]. Infected mice treated with the AAL-R, a promiscuous S1P receptor agonist [32], or the S1P₁-receptor specific agonist, CYM-5442 show suppression of early inflammatory cells recruitment through dampen cytokine production by lung endothelial cells [33]. The achievement provides insight into immune cell trafficking and immune responses points to a new shield for cytokine storm.

In summary, monitoring the change of those key cytokines as biomarkers, such as IL-6, TNF- α , while improving the efficiency of virus detection goes a long way towards evaluating the disease severity of flu patients and guiding physicians to institute immunomodulatory treatment accompanied by antiviral treatment. In the process of research on the network and the kinetics of cytokines responses, comprehending the critical part of host reaction should be attached to more importance.

References

- Lacy P, Stow JL (2011) Cytokine release from innate immune cells: association with diverse membrane trafficking pathways. *Blood* 118: 9-18.
- Tisoncik JR, Korth MJ, Simmons CP, Farrar J, Martin TR, et al. (2012) Into the eye of the cytokine storm. *Microbiol Mol Biol Rev* 76: 16-32.
- Ferrara JL, Abhyankar S, Gilliland DG (1993) Cytokine storm of graft-versus-host disease: a critical effector role for interleukin-1. *Transplant Proc* 25: 1216-1217.
- La Gruta NL, Kedzierska K, Stambas J, Doherty PC (2007) A question of self-preservation: immunopathology in influenza virus infection. *Immunol Cell Biol* 85: 85-92.
- D'Elia RV, Harrison K, Oyston PC, Lukaszewski RA, Clark GC (2013) Targeting the "cytokine storm" for therapeutic benefit. *Clin Vaccine Immunol* 20: 319-327.
- Gao R1, Bhatnagar J, Blau DM, Greer P, Rollin DC, et al. (2013) Cytokine and chemokine profiles in lung tissues from fatal cases of 2009 pandemic influenza A (H1N1): role of the host immune response in pathogenesis. *Am J Pathol* 183: 1258-1268.
- Li C, Yang P, Sun Y, Li T, Wang C, et al. (2012) IL-17 response mediates acute lung injury induced by the 2009 pandemic influenza A (H1N1) virus. *Cell Res* 22: 528-538.
- Ichikawa A, Kuba K, Morita M, Chida S, Tezuka H, et al. (2013) CXCL10-CXCR3 enhances the development of neutrophil-mediated fulminant lung injury of viral and nonviral origin. *Am J Respir Crit Care Med* 187: 65-77.
- Takano T, Tajiri H, Kashiwagi Y, Kimura S, Kawashima H (2011) Cytokine and chemokine response in children with the 2009 pandemic influenza A (H1N1) virus infection. *Eur J Clin Microbiol Infect Dis* 30: 117-120.
- Lee N, Wong CK, Chan PK, Chan MC, Wong RY, et al. (2011) Cytokine response patterns in severe pandemic 2009 H1N1 and seasonal influenza among hospitalized adults. *PLoS One* 6: e26050.
- Wang S, Le TQ, Kurihara N, Chida J, Cisse Y, et al. (2010) Influenza virus-cytokine-protease cycle in the pathogenesis of vascular hyperpermeability in severe influenza. *J Infect Dis* 202: 991-1001.
- Cheng XW, Lu J, Wu CL, Yi LN, Xie X, et al. (2011) Three fatal cases of pandemic 2009 influenza A virus infection in Shenzhen are associated with cytokine storm. *Respir Physiol Neurobiol* 175: 185-187.
- Wong CK, Lam CW, Wu AK, Ip WK, Lee NL, et al. (2004) Plasma inflammatory cytokines and chemokines in severe acute respiratory syndrome. *Clin Exp Immunol* 136: 95-103.
- Chi Y, Zhu Y, Wen T, Cui L, Ge Y, et al. (2013) Cytokine and chemokine levels in patients infected with the novel avian influenza A (H7N9) virus in China. *J Infect Dis* 208: 1962-1967.
- Jiangfang Z, Dayan W, Rongbao G (2013) Biological features of novel avian influenza A (H7N9) virus. *Nature*, doi:10.1038/nature12379
- Sun L, Louie MC, Vannella KM, Wilke CA, LeVine AM, et al. (2011) New concepts of IL-10-induced lung fibrosis: fibrocyte recruitment and M2 activation in a CCL2/CCR2 axis. *Am J Physiol Lung Cell Mol Physiol* 300: L341-L353.
- Dann SM, Le C, Choudhury BK, Liu H, Saldarriaga O, et al. (2014) Attenuation of Intestinal Inflammation in Interleukin-10-Deficient Mice Infected with *Citrobacter rodentium*. *Infect Immun* 82: 1949-1958.
- Li J, Li H, Mao H, Yu M, Yang F, et al. (2013) Impaired NK cell antiviral cytokine response against influenza virus in small-for-gestational-age neonates. *Cell Mol Immunol* 10: 437-443.
- Costanzo ES, Lutgendorf SK, Kohut ML, Nisly N, Rozeboom K, et al. (2004) Mood and cytokine response to influenza virus in older adults. *J Gerontol A Biol Sci Med Sci* 59: 1328-1333.
- Yamaya M (2012) Virus infection-induced bronchial asthma exacerbation. *Pulm Med* 2012: 834826.
- Giuffrida MJ, Valero N, Mosquera J, Alvarez de Mon M, Chacín B, et al. (2014) Increased cytokine/chemokines in serum from asthmatic and non-asthmatic patients with viral respiratory infection. *Influenza Other Respir Viruses* 8: 116-122.
- Seo SH, Goloubeva O, Webby R, Webster RG (2001) Characterization of a porcine lung epithelial cell line suitable for influenza virus studies. *J Virol* 75: 9517-9525.
- Seo SH, Hoffmann E, Webster RG (2002) Lethal H5N1 influenza viruses escape host anti-viral cytokine responses. *Nat Med* 8: 950-954.
- Ng YP, Lee SM, Cheung TK, Nicholls JM, Peiris JS, et al. (2010) Avian influenza H5N1 virus induces cytopathy and proinflammatory cytokine responses in human astrocytic and neuronal cell lines. *Neuroscience* 168: 613-623.
- Bermejo-Martin JF, Martin-Loeches I, Rello J, Antón A, Almansa R, et al. (2010) Host adaptive immunity deficiency in severe pandemic influenza. *Crit Care* 14: R167.
- Agrati C, Gioia C, Lalle E, Cimini E, Castilletti C, et al. (2010) Association of profoundly impaired immune competence in H1N1v-infected patients with a severe or fatal clinical course. *J Infect Dis* 202: 681-689.
- Mussá T, Ballester M, Silva-Campa E, Baratelli M, Busquets N, et al. (2013) Swine, human or avian influenza viruses differentially activates porcine dendritic cells cytokine profile. *Vet Immunol Immunopathol* 154: 25-35.
- de Jong MD, Simmons CP, Thanh TT, Hien VM, Smith GJ, et al. (2006) Fatal outcome of human influenza A (H5N1) is associated with high viral load and hypercytokinemia. *Nat Med* 12: 1203-1207.
- Abraham E, et al. (1997) p55 Tumor necrosis factor receptor fusion protein in the treatment of patients with severe sepsis and septic shock. A randomized controlled multicenter trial. *Ro 45-2081 Study Group. JAMA* 277: 1531-1538
- Masihi KN (2003) Progress on novel immunomodulatory agents for HIV-1 infection and other infectious diseases *Expert Opin. Ther. Pat.* 13: 867-882
- Rivera J, Proia RL, Olivera A (2008) The alliance of sphingosine-1-phosphate and its receptors in immunity. *Nat Rev Immunol* 8: 753-763.
- Marsolais D, Hahm B, Walsh KB, Edelmann KH, McGavern D, et al. (2009) A critical role for the sphingosine analog AAL-R in dampening the cytokine response during influenza virus infection. *Proc Natl Acad Sci U S A* 106: 1560-1565.
- Teijaro JR, Walsh KB, Cahalan S, Fremgen DM, Roberts E, et al. (2011) Endothelial cells are central orchestrators of cytokine amplification during influenza virus infection. *Cell* 146: 980-991.