

Clinical Image

Open Access

Hyperkeratotic Psoriasis

Abhishek Maiti^{1*} and Avash Das²

¹Department of Internal Medicine, The University of Texas Health Science Center at Houston, USA

²Department of Cardiology, Beth Israel Deaconess Medical Center, Boston, USA



Figure 1a: Hyperkeratotic plaques over the dorsum of the hand dorsum of hand.



Figure 1b: Hyperkeratotic plaques over knees and shins.

A 49 year old woman was brought to the emergency department with altered mental status. Physical examination revealed tachycardia, hypotension. Skin examination showed hyperkeratotic plaques over the dorsum of the hand [Figure 1a], and extensive thickening, scaling, desquamation of the skin, with hyperkeratotic plaques over knees and shins bilaterally [Figure 1b]. She was found to be in septic shock and was admitted to the intensive care unit. After improvement of sepsis, a skin biopsy of the lesions revealed hypogranulosis, confluent parakeratosis, tortuous blood vessels in papillary dermis, suprapapillary plate thinning, Monro's microabscess, and spongiform pustule of Kogoj which confirmed the diagnosis of hyperkeratotic psoriasis.

Hyperkeratotic psoriasis also referred to as palmoplantar psoriasis is a variant of chronic plaque psoriasis. This has been reported to be a common variant of psoriasis among children, but is very rare in adults [1,2]. Well demarcated lesions, in the absence of other clinical signs of psoriasis, often make the diagnosis of hyperkeratotic psoriasis difficult. Treatment of this subtype has been matter of speculation with emollients being the first line of treatment for dry scales. In case of severe hyperkeratotic palmoplantar psoriasis, either oral methotrexate alone or combination of etanercept and alitretinoin have been demonstrated to be well tolerated and effective treatment options [3-5]. Untreated or refractory patients can present with such extensive lesions.

References

1. Amode R, Hadj-Rabia S, Bursztejn AC (2015) Palmoplantar psoriasis, a frequent and severe clinical type of psoriasis in children. *J Eur Acad Dermatol Venereol*.
2. Rotchford JP, Hayman AB (1961) Extreme hyperkeratotic psoriasis in a mongoloid -A case report. *Arch Dermatol* 83: 973-976.
3. Meyer V, Goerge T, Luger TA, Beissert S (2011) Successful treatment of palmoplantar hyperkeratotic psoriasis with a combination of etanercept and alitretinoin. *J Clin Aesthet Dermatol* 4: 45-46.
4. Wald JM, Klufas DM, Strober BE (2015) The use of methotrexate, alone or in combination with other therapies, for the treatment of palmoplantar psoriasis. *J Drugs Dermatol* 14: 888-892.
5. Mrowietz U, van de Kerkhof PC (2011) Management of palmoplantar pustulosis: Do we need to change? *Br J Dermatol* 164: 942-946.

*Corresponding author: Abhishek Maiti, Department of Internal Medicine, The University of Texas Health Science Center at Houston, 2950 Old Spanish Trail, Apt 379, Houston, TX 77054, Tel:832-696-8407; E-mail: abhishek.maiti@uth.tmc.edu

Received March 17, 2016; Accepted March 22, 2016; Published March 24, 2016

Citation: Maiti A, Das A (2016) Hyperkeratotic Psoriasis. *Gen Med (Los Angel)* 4: i114. doi:10.4172/2327-5146.1000i114

Copyright: © 2016 Maiti A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.