

Metallic Content of One Hundred Medicinal Plants

Waseem Hassan^{1*}, Shakilla Rehman¹, Hamsa Noreen¹, Shehnaz Gul¹, Syeda Nida Zainab Kazmi^{1,2}, Maryam Jan¹, Bakht Zaman¹, Ata Ur Rehman¹, Ziarat Shah¹, Ali Riaz² and Imdad Ullah Mohammadzai¹

¹Institute of Chemical Sciences, University of Peshawar, Peshawar -25120, Khyber Pakhtunkhwa, Pakistan

²Department of Chemistry, Sarhad University of Science and Information Technology, Peshawar -25120, Khyber Pakhtunkhwa, Pakistan

Abstract

Heavy metals are necessary for the normal functioning of cells and the survival of organisms. These are required for physiological and biochemical functioning of the body, but at higher concentration they causes the oxidative damage. In fact non-essential metals like (lead (Pb), mercury (Hg) and cadmium (Cd)) are more noxious even at very low concentration. It's important to note that heavy metals are present in soil, air and water can easily enter into the organisms. Literature highlighted the role of metals like chromium (Cr), iron (Fe), nickel (Ni), copper (Cu), cadmium (Cd), lead (Pb) and arsenic (As) in oxidative damage. Fenton chemistry/Haber-Weiss reaction are most extensively studied mechanisms by which heavy metals produces reactive oxygen and nitrogen species (ROS and RNS) and ultimately causes oxidative stress. Both redox-active and inactive metals can diminish cellular antioxidant system specifically thiol- containing enzymes and antioxidants and may lead to cellular death. This review will contribute in providing valuable information on metallic contents of selected plants. For the purpose, data has been collected from the literature regarding ten mineral contents (sodium (Na), potassium (K), calcium (Ca), magnesium (Mg), zinc (Zn), iron (Fe), copper (Cu), chromium (Cr), nickel (Ni) and manganese (Mn)) of 100 medicinal plants grown in Pakistan. The metallic contents shows that majority of the plants contain optimum values as compared with daily mineral intake standards. However, several factors may account for variations in mineral concentration, which include soil composition, different geographic areas and environmental changes.

Keywords: Plants; Ethnobotanical uses; Metallic content

Ethnobotanical Uses of 100 Medicinal Plants

Medicinal plants are considered as key sources of drugs for the treatment of various ailments worldwide. The flora of Pakistan due to its highly favorable diverse climatic and soil conditions, many topographical regions and ecological zones offer a great variety of medicinal plants [1]. According to a general survey, approximately 6000 flowering plants exist in Pakistan, out of which 400-600 are utilized for medicinal purposes [2,3]. Literature reported that about 80% of the rural population of Pakistan relies primarily on traditional medicines while 90% of the country's medicinal herbs are imported [4,5].

Furthermore, all the selected plants are utilized by the native people for cure of various diseases such as diarrhea, dysentery, stomach disorders, pain, wounds, skin diseases, scrofula, ulcers, urinary tract infections, bronchitis, asthma, whooping cough, loose motion, emmenagogues, toothaches, rheumatism, sensory and nervous system dysfunctions, sexual disorders, kidney troubles, piles, jaundice, hepatitis, typhoid, malarial fevers, heart diseases and immune deficiency diseases [1-5].

Metallic Content of 100 Medicinal Plants

Mineral elements are assumed of immense value as each of these elements show a distinctive individual role in the structural and functional integrity of the organization of living systems. Although minerals comprise only 4-6% of the human body and do not take part efficiently in fulfilling the energy requirements, yet their importance can be established from their involvement in a large number of physicochemical processes of utmost importance which takes place continuously in living cells and organisms [6,7]. Minerals may be generally classified as macro (sodium (Na), calcium (Ca), phosphorus (P) and chlorine (Cl)) or micro (manganese (Mn), zinc (Zn), iron (Fe), potassium (K), magnesium (Mg), copper (Cu), cobalt (Co) and chromium (Cr)) elements which mostly depend on their requirements to the body and it has been found that approximately more than 100 mg/dl of the macronutrients and less than 100 mg/dl of the micronutrients

are necessary on daily basis [6,8]. Deficiencies of the mineral elements are the foremost public health problem in many under developed countries with women and infants at high risk [9]. Studies have shown that severe cases of anaemia most likely due to mineral deficiencies are a direct cause of child and maternal mortality [10].

The medicinal plants serve as a good source of these vital mineral elements which can be consumed both for dietary and medicinal purposes. However the lack of knowledge regarding the mineral quantities in these plants often produce hazardous effects on consumers health as various plants contain toxic metals in sufficient higher amounts. Thus metallic screening of medicinal plants is of immense importance [11]. This review will contribute in providing valuable information on metallic contents of selected plants and both the beneficial and harmful effects of the minerals on human health. For the purpose, data has been collected from literature regarding ten mineral contents (Na, K, Ca, Mg, Zn, Fe, Cu, Cr, Ni and Mn) of 100 medicinal plants grown in Pakistan.

The data concerning sodium (Na) content of the selected medicinal plants have been obtained from the literature and presented in (Table 1). The highest amounts of Na are present in *S.cordifolia* (3.649%) *Z.mauritiana* (34.16 mg/g) and *E.ribes* (12500 ppm). Plants like *D.sisso* (0.07 ppm), *P.granatum* (1.279 ± 0.003 mg/L), *R.communis* (1.64 ppm) and *T.terrestris* (1.70 mg/kg) contain lower amounts of Na, while the

***Corresponding author:** Waseem Hassan, Institute of chemical sciences, University of Peshawar, Peshawar -25120, Khyber Pakhtunkhwa, Pakistan, Tel: 0092-91-9216652; E-mail: waseem_anw@yahoo.com

Received November 05, 2015; **Accepted** November 28, 2015; **Published** December 03, 2015

Citation: Hassan W, Rehman S, Noreen H, Gul S, Kazmi SNZ, et al. (2015) Metallic Content of One Hundred (100) Medicinal Plants. J Nutr Disorders Ther 5: 177. doi:10.4172/2161-0509.1000177

Copyright: © 2015 Hassan W, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Sr. No.	Plant name	Na	K	Ca	Mg	Zn	Fe	Cr	Cu	Ni	Mn
1.	<i>Terminaliaarjuna</i> (ppm) [51]		5900	3500	12100	76	750			10	86
2.	<i>Bauhinia variegata</i> (mg/kg) [51]			560 ± 40.9	545 ± 50.0	37 ± 2.2	53 ± 1.6				18 ± 0.6
3.	<i>Cuminumcuminum</i> (ppm) [52]	0.27	1.81	0.96	0.48	33	229		21		35
4.	<i>Violaodorata</i> (ppm) [53]	528	100	78	11.2	2.17	19.4				
5.	<i>Capsellabursa pectoris</i> (ppm) [54]					11.4	212	1	4.3	3.1	29
6.	<i>Allium sativum</i> (mg/100g) [55]	4.10 ± 0.14	54.00 ± 1.40	26.30 ± 0.14	4.10 ± 0.14	0.34 ± 0.17	5.29 ± 0.08		.001 ± 0.0		0.001 ± 0.0
7.	<i>Minthaspicata</i> (ppm)[56]	147.7 ± 4.68	15.56 ± 0.51	255.95 ± 9.01	3.90 ± 0.14	0.79 ± 0.025	2.03 ± 0.052	-	0.88 ± 0.045	-	-
8.	<i>Trigonellafoenum</i> (ppm) [54]	270	50	88	11	2.24	14.2				
9.	<i>Euphorbia tirucalli</i>	-	-	-	-	-	-	-	-	-	-
10.	<i>Cyperusrotundus</i>	-	-	-	-	-	-	-	-	-	-
11.	<i>Saussureaelappa</i>	-	-	-	-	-	-	-	-	-	-
12.	<i>Fumariaindica</i>	-	-	-	-	-	-	-	-	-	-
13.	<i>Solanumnigrum</i> (mg/kg) [57]	810.77 ± 57.69			185.92 ± 6.12	8.53 ± 0.29	311.88 ± 5.02		17.471 ± .84		98.43 ± 3.67
14.	<i>Zizyphusjujuba</i> (mg/100g)[58]	6.21 ± 0.24	375 ± 6.3	45.6 ± 2.1		0.47 ± 0.06	6.42 ± 0.12		0.42 ± 0.02		51.2 ± 2.8
15.	<i>Eugenia aromatic</i> (ug/g) [59]	2409 ± 81.60	1618 ± 135.6			4.94 ± 1.228	60.03 ± 0.785	0.96 ± 0.109	3.24 ± 0.261	0.7 ± 0.082	57.3 ± 6.826
16.	<i>Glycyrrhizaglabra</i> (mg/100g) [60]	63.20 ± 0.56	180.48 ± 0.12			00.96 ± 0.01	08.08 ± 0.02				
17.	<i>Piper longum</i>	-	-	-	-	-	-	-	-	-	-
18.	<i>Crocus sativus</i> (mg/100g)[61]	25.75 ± 0.01	542.13 ± 0.01	486.25 ± 0.12	2.93 ± 0.02	1.80 ± 0.13	17.99 ± 0.09		0.87 ± 0.008		
19.	<i>Pipernigrum</i>	-	-	-	-	-	-	-	-	-	-
20.	<i>Tagetesminuta</i> (ppm) [53]	208	80	109	11.5	2.33	14.3	ND	ND	ND	ND
21.	<i>Thymuslinearis</i>	-	-	-	-	-	-	-	-	-	-
22.	<i>Carumcarvi</i> (mg/kg) [53]	0.1	44.5	11.8	3.1	22.3	74.7	0.3	5.2	59.8	ND
23.	<i>Menthapiperata</i> (mg/kg) [62]			12150	3602	12.64	531.5	5.41	11.5		70.8
24.	<i>Acaciamedesta</i> (ppm) [63]	1.8	19000			10.2			3.4		
25.	<i>Nigella sativa</i> (mg/kg) [64]	0.35 ± 0.02	0.83 ± 0.01%	9.13 ± 0.16	10.20 ± 0.09	0.05 ± 0.00	0.26 ± 0.02		0.03 ± 0.00		103.9 ± 3.69
26.	<i>Acacia nilotica</i> (ppm) [65]	87.25	549	2.08	2.075						
27.	<i>Adhatodavasica</i> [66]		2000 ppm	20900 ppm	20500 ppm		1250				67
28.	<i>Psidiumguajava</i> (mg/100g) [67]	750.9	895	172	158.7	1.84	11.71		1.38		1.38
29.	<i>Cassia fistula</i> (%) [68]	118.4	838	925	947.4	70.67	179.5				37.2
30.	<i>Cannabis sativa</i> [69]	-	-	-	-	2.25 ± 0.01 mg/l	135.20 ± 0.12 mg/l	-	26.6 ± 0.01 mg/l	-	40.30 ± 0.05
31.	<i>Allium cepa</i> (mg/100g) [70]	5.4	140	22	20	2	0.2	-	-	-	-
32.	<i>Phoenix dactylifera</i> (mg/100 g) [71]	3.49 ± 0.12	624 ± 5	81.9 ± 1.9	60.9 ± 1.2	0.45 ± 0.02	0.60 ± 0.11	0.01 ± 0.02	0.77 ± 0.03		0.30 ± 0.04
33.	<i>chenopodium album</i> (ppm) [72]	375	10455	4242	45460	0.2	102.8	1.8	4.3	0.3	
34.	<i>Abutilon indicum</i> (mg/100g) [73]		261.01 ± 0.01	237.32 ± 0.03	176.50 ± 0.052	1.94 ± 0.03	3.13 ± 0.10				
35.	<i>Zanthoxylumarmatum</i> (m/100g) [74]	5.84 ± 2.09	87.84 ± 8.34	96.78 ± 6.86	9.80 ± 2.61	0.032 ± 0.012	0.425 ± 0.02	ND	6.34 ± 2.69		0.043 ± 0.017
36.	<i>Citrus limon</i> (mg/100g) [75]	755.50 ± 0.058	8600.00 ± 0.028	8452.50 ± 0.050	1429.50 ± 0.008	13.94 ± 0.007	147.65 ± 0.068		4.94 ± 0.012		
37.	<i>Sidacordifolia</i> (%) [76]	3.649	10.2	13.1		5.163	1.135	1.31	8.9		1.86
38.	<i>Corriandrumsativum</i> (ppm) [77]					59.4	150.4	<0.003	41.4	<0.006	
39.	<i>Ricinuscommunis</i> (ppm) [78]	1.64p	0.79	3.21	0.19	0.49	4.1	3.41	5.2		0.92
40.	<i>Lepidiumsativum</i> (mg/100g) [79]	12.4	85.3	8.2	31.8	1	1.1		0.1		0.3
41.	<i>Sesamumindicum</i> (mg/kg) [80]	122.50 ± 4.21	851.35 ± 3.44	415.38 ± 3.14	579.53 ± 0.42						

42.	<i>Curcuma longa</i> (mg/kg) [81]	210	7940	0.12	0.28	57.06			5.71		429
43.	<i>Grewiaoptiza</i> (%) [82]	0.15	0.64	3.35							
44.	<i>Valerianawallichii</i> (ppm) [83]					11.64 ± 0.11	140.8 ± 0.17	266 ± 0.12	11.28 ± 0.06		6 ± 0.05
45.	<i>Tribulusterrestris</i> (mg/kg) [84]	1.7	55	28	32						
46.	<i>Origanumvulgare</i> (mg/100g) [85]	47.2	1970	164	80.67	1.5	17.26	0.26	0.58	0	0.77
47.	<i>Daturastromium</i> (mg/g) [75]	2.5	2.5	295	307.5	0.51	5.63	2.85	8.07		8.49
48.	<i>Swertiachirata</i> (ppm) [86]					45.4			9.8		
49.	<i>Punicagranatum</i> (mg/L) [87]	1.279 ± 0.003	55.19 ± 0.021	1.650 ± 0.00	3.721 ± 0.001	0.430 ± 0.001	0.466 ± 0.003		0.231 ± 0.003		0.033 ± 0.002
50.	<i>Cedrusdeodara</i> (mg/100g) [88]		0.06	2.6	0.017						
51.	<i>Hippophaërhamnoides</i> (ppm) [54]	-	-	-	-	8.1	87	0.4	4.5	4	122
52.	<i>Menthalongifolia</i> (µg/g)/ mg/l)±SD	373.1 ± 0.40 [89]	8372 ± 16 [89]	2810 ± 10 [89]	3350 ± 13 [89]	0.84 [69]	0.12 [69]	0.51 [69]	0.58 [69]		0.25 [69]
53.	<i>Juglansregia</i> (mg/100 g) [90]	0.25 ± 0.02 [91]	3.02	0.79	0.51	3.19 ± 0.85 [91]	5.88 ± 1.10 [91]	0.85 ± 0.44 [91]	3.12 ± 1.48 [91]	*	10.45 ± 5.12 [91]
54.	<i>Plantago ovata</i> (ug/g) [72]	1200 ± 60	1000 ± 60	1600 ± 80	63.5 ± 10.5	99.4 ± 10.8	21.7 ± 4.8	11.754 ± 0.01	59 ± 7.3	0.0213 ± 0.0001	6.0 ± 1.2
55.	<i>Berberislyceum</i> [92]	14.5 ± 0.11 mg/100 g	161.42 ± 0.41 mg/100 g	18.272 ± 0.212 mg/100 g	0.541% [93]	27.50 mg/kg [93]	2649.99 mg/kg [93]		33.50 mg/kg [93]		58.67 mg/kg [93]
56.	<i>Withaniasomnifera</i> (%) [94]	0.54	2.2	0.16	0.184	44.1 ppm	349.5 ppm	4 ppm [95]	33 ppm	7 ppm [95]	59.0 ppm
57.	<i>Dalbergiasisso</i> [13]	0.07 ± 0.02	2.4 ± 0.70	64.5 ± 1.60	7.4 ± 0.50	*	1.5 ± 0.10	*	*	*	0.03 ± 0.01
58.	<i>Pyruspyrifolia</i> (mg/100 g) [96]	4	96	9	7	0.09	0.29	-	0.05	-	0.03
59.	<i>Vernoniaamygdalina</i> (mg/100g) [97]	-	-	-	0.43 ± 0.00	0.04 ± 0.01	0.14 ± 0.01	0.04 ± 0.01	0.10 ± 0.00	-	0.07 ± 0.03
60.	<i>Trachyspermumammi</i> (ug/g) [98]	-	-	-	-	80.6 ± 24.1	2792 ± 304	-	145 ± 27	-	771 ± 11
61.	<i>Caricapapaya</i> (mg/100 g) [99]	4	223	24.9	23.54	0.056	2.56	-	0	-	0.01
62.	<i>Citrulluscolocynthis</i> (mg/100g) [100]	1.75b ± 0.01	2.85c ± 0.05	1.90b ± 0.01	1.10 b ± 0.02	0.31a ± 0.02	0.10b ± 0.01	*	0.10a ± 0.01	*	
63.	<i>AzadirachtaindicaA. Juss</i> (mg/100 g) [101]	-	-	-	0.69 ± 0.01 %	0.06 ± 0.01	0.14 ± 0.01	0.58 ± 0.00*	0.06 ± 0.01*	0	0.06 ± 0.03
64.	<i>Raphanussativus</i> (mg/100 g) [102]	10.7 a**	209 a**	21.0 c**	10.2 b**	0.09 b**	0.73 b**	0.01 c**	0.07 a*	0.07 a**	0.09 b**
65.	<i>Tamarindusindica</i> (mg/100 g) [103]	40.20a ± 0.10	38.93a ± 0.14	454.74c ± 0.13	16.54b ± 0.10	29.60b ± 0.17	27.36a ± 0.60		1.45b ± 0.04		0.20c ± 0.01
66.	<i>Pisum sativum L</i> (mg/100 g) [104]	262 a	756	61	84.06 [105]	3.97 [105]	8				
67.	<i>Foeniculumvulgare</i> (µg/g) [106]	69.61 ± 0.532 [59]	9038 ± 571.0 [59]			212.56cd	451.02d	0.53a	37.1 ± 1.220 [17]	9.00b	41.2 ± 1.264 [59]
68.	<i>Morus alba</i> (ppm) [107]	160.65 ± 1.45	0.94 ± 0.02	0.02 ± 0.001	0.04 ± 0.001	20.08 ± 0.52	47.90 ± 2.23	<0.006 ± 0.001	1.61 ± 0.01	0.34 ± 0.01	3.95 ± 0.07
69.	<i>Moringa oleifera</i> (ppm) [108]		732 ± 164	602 ± 122	4BDL	BDL	BDL	BDL			
70.	<i>Ipomoea batatas</i> (mg/100g) [109]	4.23 ± 0.02	4.50 ± 0.01	28.44 ± 0.03	340.00 ± 0.17	0.08 ± 0.00		16.00 ± 0.04	0.00 ± 0.00		4.64 ± 0.10
71.	<i>Rumexsagittatus</i> (mg/100g) [110]	0.177	3	2.31	0.703	39	441		14		80
72.	<i>Taraxacumofficinale</i> (mg/100g) [111]					88 [112]	20.06 ± 0.009	0.08 ± 0.005	1.35 ± 0.02	0.21 ± 0.004	1.79 ± 0.03

73.	<i>Oscimum sanctum</i> (mg/100g) [113]	156	180	45	0.48 to 0.18 [114]	74.7	36.6		3	10.2	
74.	<i>Oscimum basilicum</i> (ppm) [115]	280 mg/kg [116]	28	17460 mg/kg [116]	34	2.00 to 0.01 [117]	15.15 to 0.20 [117]				2.01 to 0.80 [117]
75.	<i>Rubus ellipticus</i> (mg/100g) [118]	-	1.82 ± 0.25	-	5.60 ± 0.15	-	-	-	-	-	-
76.	<i>Cressacretica</i> [119]					55.3-70.2 mg/kg	125.2-151.1 mg/kg		12.2-14.3 mg/kg		24.6-28.9 mg/kg
77.	<i>Acorus calamus</i> (mg/kg) [120]	-	-	-	-	40.8	-	11.5	26.4	9.6	-
78.	<i>Citrus sinensis</i> [121]	-	-	-	-	-	-	-	-	0.08 mg/kg	-
79.	<i>Mangifera indica</i> (mg/kg) [122]	-	-	-	-	12.22 ± 3.20	-	0.66 ± 0.12	5.74 ± 0.70	0.06 ± 0.06	-
80.	<i>Gymnema sylvestre</i> (mg/kg) [123]	-	-	-	-	0.0092 ± 0.008	0.0110 ± 0.003	0.2108 ± 0.016	32.8 mg/kg [126]	-	0.0559 ± 0.02 mg/kg
81.	<i>Eclipta prostrata</i> (mg/kg) [124]	-	-	-	-	1.61	23.9	0.47	1.18	-	0 6.97
82.	<i>Peganum harmala</i> (ppm) [125]	148.88 ± 8.16	16176.93 ± 2686.17			32.52 ± 4.28	318.72 ± 36.27	16.36 ± 1.29	18.72 ± 3.24	9.10 ± 2.93	74.60 ± 5.25
83.	<i>Embeliaribes</i> % [17]	1.25%	####			0.05%	0.23%	####	####		####
84.	<i>Silybum marianum</i> [126]	4 mg/kg [127]	2 mg/kg [127]			99.50 mg/kg	360 mg/kg	48.80 mg/kg	27.25 mg/kg	33.75 mg/kg	30.00 mg/kg
85.	<i>Musa paradisiaca</i> [128]	84.53 ppm	3.96 ppm			1.01 ppm	27.83 ppm		1.89 ppm	6.10 mg/kg [129]	18.82 ppm
86.	<i>Woodfordia fruticosa</i> (mg/100g) [130]	500 ± 8.54	150 ± 12.52			1.589 ± 0.14	55.14 ± 0.48		0.640 ± 0.011		1.06 ± 0.11
87.	<i>Vitex negundo</i> (mg/ml) [131]		1618 mg/ml			2.8 mg/ml	53.3 mg/ml		0.2 mg/ml	0.07 mg/ml	
88.	<i>Broussonetia papyrifera</i> [132]		9098.7 ± 14.18 mg/kg			29.8 ± 101 mg/kg	414.8 ± 8.23 mg/kg	3.48 ± 0.11 mg/kg	19.75 ± 0.43 mg/kg		44.56 ± 0.76 µg/kg
89.	<i>Urtica dioica</i> (mg/kg) [133]					113 mg/kg	432 mg/kg	6.6 mg/kg	14.0 mg/kg	9.1 mg/kg	291 mg/kg
90.	<i>Verbascum thapsus</i> (µg/g) [13]	1624 ± 3.46	6725 ± 1.15			24 ± 1.10	394 ± 2.45	7 ± 0.00	7 ± 1.15	9 ± 1.45	15 ± 0.00
91.	<i>Malus domestica</i> [134]		102.35 ± 0.32 mg [135]			3.14 mg/kg		0.144b mg/kg	3.30 mg/kg	0.181b mg/kg	
92.	<i>Mucuna pruriens</i> (ppm) [136]					32.48 ± 0.04	33.21 ± 0.03	0.27 ± 0.01	8.43 ± 0.07	3.18 ± 0.02	14.70 ± 0.01
93.	<i>Tinospora cordifolia</i> (ppm) [75]					3.4	37.6		ND	ND ND	
94.	<i>Stevia rebaudiana</i> (mg/100g) [137]	14.93 ± 0.05	21.15a ± 0.35			1.26f ± 0.72	5.89d ± 0.34	1.45 ± 0.01 ppm [138]	0.73g ± 0.002		2.89e ± 0.58 \
95.	<i>Zingiber officinale</i> (mg/100g) [139]					12.23 ± 0.16	80.00 ± 2.89		12.50 ± 0.29	BDL	7.33 ± 0.22
96.	<i>Euphorbia hirta</i> [140]	-	-	-	-	90.35 ± 70.54	-	-	-	-	-
97.	<i>Mimosa pudica</i> (mg/kg) [141]	-	-	-	-	63.14 mg/kg	-	-	102.31 mg/kg	-	-
98.	<i>Ziziphus mauritiana</i> (mg/g) [142]	34.16 mg/g	60.21 mg/g	-	-	0.23 mg/g	7.21 mg/g	0.13 mg/g	0.004 mg/g	0.002 mg/g	0.0234 mg/g
99.	<i>Chenopodium ambrosioides</i>	-	-	-	-	-	-	-	-	-	-
100.	<i>Cynodon dactylon</i> [143]	3.41 - 0.0159 (g kg ⁻¹ DM) [144]	31.6 ± 2.2 mg/kg [145]			259.91 mg/kg	9035.16 mg/kg		190.12 mg/kg	2.58 ± 0.15 mg/kg [146]	659.49 mg/kg

Table 1: Metallic Contents of 100 Medicinal Plants.

mean Na value for all the other plants is projected as ± 1400 mg/kg. The recommended daily minimum intake of Na is estimated as 2400 mg [12]. Na exists as the primary cation in extracellular fluids and is known for regulation of membrane potentials, cell permeability, muscle irritability, conduction of nerve impulses and keeping a constant

osmotic pressure [8,13,14]. Hyponatremia results due to excess serum concentration of Na and this has been suggested as a consequence of various factors such as, administration of several hormones, excessive release of adrenocorticotropin in Cushion's disease and diabetes insipidus whereas hyponatremia is reported in diarrhea, vomiting,

severe burns, nephrosis, intestinal obstruction and Addison's disease [7].

The potassium (K) content of the plants ranged from 0.0024 ppm in *D.sisso* to (1618 mg/mL) in *V. negundo*, whereas the average K content of the other selected plants is found as ± 6000 mg/kg. *D.stromium* (295.10 mg/g), *E.ribes* (21.5%), *S.cordifolia* (10.21%), *C.limon* (8600.00 \pm 0.028 mg/100g), *W.somnifera* (2.2%), and *A.modesta* (19000 ppm) are also considered as good sources of K as their K values are far more higher than the minimum K daily intake value which is 3500 mg [12]. The importance of K is speculated from its participation in large number of biological processes such as acid base balance, movement of muscles, nerve impulse conduction and regulation of osmotic pressure. K is the major cation present in the intracellular fluid, its reduced concentration has been associated with paralysis, muscular weakness, mental confusions [7,8,14], cardiac arrhythmias, defects in renal functions and carbohydrate tolerance [15,16].

The data obtained from literature (Table 1) revealed the presence of calcium (Ca) in highest concentration in plants like *S.cordifolia* (13.09%) followed by *C.limon* (8452.50 \pm 0.050 mg/100 g) and *A.vasica* (20900 ppm), while lowest Ca contents are estimated in *M. alba* (0.02 ppm), *C. longa* (0.12 ppm) and *C.cuminum* (0.96 ppm). The recommended daily Ca intake required for normal biochemical activities of the body is 1500 mg. Ca is known as an indispensable macro mineral for the composition of bones and teeth. Ca deficiency implements rickets and osteomalacia in children and adults, respectively. This may also contribute in the pathogenesis of osteoporosis resulting in weak and porous bones and considerable high risks of fracture [8,11]. Ca plays significant role in membrane permeability and regulates the transmission of nerve impulses, any decrease in the extracellular blood Ca may results in tetany and convulsions, due to impetuous discharges of nerve impulses [17,7,8]. Ca converts prothrombin to thrombin efficiently and thus helps in blood coagulation; it increases the absorption of dietary vitamin D, clotting of milk and activation of enzymes such as lipase, ATPase succinic dehydrogenase. Ca ions normally aids in the strength of cardiac muscles, however excess absorption of Ca may cause Ca toxicity and leads to cardiac failure [18].

Medicinal plants such as *D.stromium*, *C.album* and *A.vasica* contain maximum magnesium (Mg) contents of 307.49 mg/g, 45460 ppm and 20900 ppm respectively. The minimum amount of Mg were found in *D.sisso* (0.0074 ppm), *C.deodra* (0.017 mg/100g) and *R. communis* (0.19 ppm), while the mean Mg quantity present in all the plants is projected as ± 350 , which is fairly in comparison with the predicted daily Mg value (400 ppm). Mg is required for teeth, bones, several enzyme systems [8] and for maintenance of osmotic pressure in extracellular fluids and plasma. Mg depletion results in chronic vomiting, diarrhea, vasodilation with erythema and hyperemia. More serious situations have been reported in long term deficiency of Mg such as neuromuscular hyperirritability and cardiac arrhythmia [18,19].

Data obtained from literature concerning the quantity of zinc (Zn) in medicinal plants revealed the presence of considerable higher amount of Zn in *S.cordifolia* (5.163%), *V.negundo* (2.8 mg/ml), *E.ribes* (0.0450%) and *C.limon* (13.94 \pm 0.007 mg/100 g), while *G.sylvestre*, *C.album*, *Z.armatum*, *C.cuminum*, *P.granatum*, and *R.communis* contain lesser amount of Zn as 0.0092 \pm 0.008 ppm, 0.2 ppm, 0.032 \pm 0.012 mg/100g, 0.401 ppm, 0.430 \pm 0.001 mg/L and 0.49 ppm respectively. The essential Zn value needed on daily basis is 11 mg which is slightly less than the projected Zn content in the mentioned plants i. e. ± 20 . The importance of Zn is elucidated from its role as a

co-factor for majority of enzymes, performing remarkable functions in living systems such as cell replication and metabolism of nucleic acids and other macronutrients [20-22]. In addition, Zn is considered as an integral component of insulin, a key factor in protein synthesis, wound healing and repairment of tissues [8,23,24]. Decreased amount of Zn is associated with the metabolic disorders, growth retardation, skin abrasion and coronary heart pathologies [18,25].

The occurrence of iron (Fe) in medicinal plants is ranged from a lowest value in *G.sylvestre* (0.0110 \pm 0.003 mg/kg) to a quantity as much as high in *V.negundo* (53.3 mg/ml). The suggested mean value for Fe in all the plants is ± 400 , while the daily dietary requirements of Fe are 15mg. Fe exists primarily as hemoglobin and transferrin in the erythrocytes and plasma fractions of the blood, respectively. Fe is considered vital in composition of cytochromes, myoglobin, hemoglobin and succinate dehydrogenase as well as a co-factor for enzymes which participate in the synthesis of neurotransmitters [26]. The deficient concentrations of Fe intake have been associated with impairments in brain development and functioning, protein synthesis, restless legs syndrome and organogenesis [27]. Exceeded levels of Fe in various organs and tissues including brain can cause haemosiderosis, type-1 neurodegeneration, Alzheimer's disease and Parkinson's disease [7,8]. The role of Fe in the body is clearly allied with the transfer of oxygen from lungs to the tissue cells and hemoglobin [28] moreover; its lack is the most rampant nutritional deficit in humans and is usually caused by deficient dietary intake [29]. Furthermore, *S.cordifolia* having (1.309%) of chromium (Cr) is considered to be the rich Cr containing medicinal plant, while the minimum Cr concentration was found in *C.sativum* (<0.003 ppm) of all the selected plants for this review. The FAO/WHO estimated permissible value of Cr should not be more than 2 ppm [30] which is in close proximity with the mean Cr content of all the plants, which is ± 4 . Cr is an efficient mediator of carbohydrate metabolism and an imperative component of glucose tolerance factor (GTF), which enhances the action of insulin and sustains normal levels of glucose tolerance [8,31]. Cr deficiency leads to hyperglycemia, cataracts and atherosclerosis [32]. The role of Cr in maintenance of the configuration of RNA molecule, protein and lipid metabolism is also known from various animal and human studies [31,33]. Cr toxicity due to its increased concentration in body tissues results in impairment of liver, kidney, blood and nervous system functions and ultimately death [34,35].

The maximum amount of copper (Cu) is present in plants like *S.cordifolia* (8.899%), *W.fruiticosa* (6400 mg/100 kg) and *D.stromium* (8.07 mg/g), whilst *A.sativum* (0.001 \pm 0.0 ppm), *N.sativa* (0.03 \pm 0.00 mg/kg), and *C.papaya* (0.001 mg/100 g) are considered as lowest Cu containing plants. The daily permissible value of Cu in diet is recommended as 900 mg for children and adults. Cu is an important redox-active element and an indispensable component of various enzyme systems including ceruloplasmin, cytochrome oxidase, cytosolic superoxide dismutase and lactase. Cu facilitates the absorption and incorporation of Fe in hemoglobin, thus regulates hemoglobin synthesis and its deficiency may cause anemia [8,36,37]. Cu is also necessary for the smooth functioning of neurologic and hematologic systems [27]. However, excess amount of dietary Cu accumulates in liver resulting in a decline in blood hematologic concentration and Cu poisoning, which adversely affects the functioning of liver and as a consequence cause jaundice due to erythrocyte haemolysis [18].

Nickel (Ni) is quite low in all the selected medicinal plants ranged from <0.006 ppm in *C.sativum* to 0.07 mg/mL in *V.negundo*. The mean Ni content for all the plants is projected as ± 2 which is comparable to

the WHO recommended value of Ni (1.63ppm) in edible plants [38], also higher amount of Ni is noxious for health [39].

In addition, the absorbance of Ni by the body is very low [40], yet it has been suggested that Ni contribute in the metabolism of nucleic acids, preservation of membrane structure, control of prolactin and act as a key component of enzymes such as plant ureases and several hydrogenases. Also, experimental findings reported the malfunctioning of chicks and rats' liver when fed on Ni deficient diets [41]. Ni is present mostly in the pancreas and helps in regulation of insulin while its deficiency results in liver disorders [42].

Similarly, Manganese (Mn) rich plants include *S.cordifolia* (1.863%), *D.stromium* (8.49 mg/g) and *T.ammi* (771 ± 11 ppm), while *A.sativum* (0.001 ± 0.0 mg/100 g), *P.granatum* (0.033 ± 0.002 mg/L) and *D.sisso* (0.03 ± 0.01 ppm) contain lowest amounts of Mn. Mn is required for several primary biochemical processes as it has been known as an active component of a number of enzymes including decarboxylases, transferases and hydrolyses [8,43]. Literature reported the pharmacological inference of Mn in anticipation and cure of diabetes mellitus [44]. Moreover, the existence of Mn in mitochondrial superoxide dismutase provides an evidence for the involvement of Mn in oxidative phosphorylation reactions. Optimum Mn levels maintain the normal bone structures and regulate functioning of central nervous system (CNS), while reduced amount lead to reproductive failure in both sexes [45], in addition Mn overexposure accounts for drastic effects on CNS and mood [27].

Conclusion

The human body requires a number of minerals in order to sustain good health [48]. Macro and microelements play a significant part in the metabolism, composition of certain proteins (hemoglobin and haemoprotein) and some mineral elements stay chelated with organic ligands and make them bioavailable to the body system [47-49]. The excessive amount of elements or specifically metals can lead to the generation of free radicals and consequently oxidative damage. The current study was aimed to review and compile ethnobotanical uses of 100 medicinal plants (which belongs to 53 families). The selected plants are utilized by the native people for cure of various diseases. The metallic contents of all the plants are presented here, shows that majority of the plants contain optimum values as compared with daily mineral intake standards. However, several factors may account for variations in mineral concentration of a particular plant, which include soil composition, different geographic areas and environmental changes.

Reference

- Hussain K, Shahzad A, Zia-ul-Hussnain S (2008) An ethnobotanical survey of important wild medicinal plants of Hattar district Haripur, Pakistan. Ethnobotanical Leaflets 12: 29-35.
- Abbasi AM, Khan, MA, Ahmed M, Zafar M (2010) Herbal medicines used to cure various ailments by the inhabitants of Abbottabad district, North West Frontier Province, Pakistan. Indian J Trad Know 9: 175-183.
- Hamayun M (2005) Ethnobotanical profile of Utror and Gabral valleys, district Swat, Pakistan. Ethnobotanical Leaflets 2005: 9.
- Saqib Z, Sultan A (2005) Ethnobotany of Palas valley, Pakistan. Ethnobotanical Leaflets 2005: 28.
- Trivedi PC (2002) Ethnobotany: An overview. Ethnobotany. Jaipur: Aavishkar publisher 1.
- Eruvbetine D (2003) Canine Nutrition and Health. A paper presented at the seminar organized by Kensington Pharmaceuticals Nig. Ltd., Lagos.
- Wadhwa N (2015) Textbook of Pediatric Gastroenterology, Hepatology and Nutrition: Anupam Sibal, Sarath Gopalan (eds), Akshay Kapoor, Vidyut Bhatia (co-eds) : Published by Jaypee Brothers Medical Publishers (P) Ltd, New Delhi, India; First Edition: 2015; ISBN: 978-93-5152-740-4. Indian J Pediatr .
- Murray RK, Granner DK, Mayes PA, Rodwell VW (2000) Harper's Biochemistry, (25th edtn), McGraw-Hill, Health Profession Division, USA.
- Batra J, Seth PK (2002) Effect of iron deficiency on developing rat brain. Indian J Clin Biochem 17: 108-114.
- Chakravarty I, Ghosh K (2000) Micronutrient malnutrition--present status and future remedies. J Indian Med Assoc 98: 539-542.
- Saiki M, Vasconcellos MB, Sertié JA (1990) Determination of inorganic components in Brazilian medicinal plants by neutron activation analysis. Biol Trace Elem Res 26-27: 743-50.
- Baysal A (2002) Fundamentals of Nutrition. Hatipoglu Press, Ankara (in Turkish).
- Lokhande R, Singare P, Andhale M (2010) Study on Mineral content of Some Ayurvedic Indian Medicinal Plants by Instrumental Neutron Activation Analysis and AAS Techniques. Health Sci J 4: 157-168.
- Hays VW, Swenson MJ (1985) Minerals and Bones. In: Dukes' Physiology of Domestic Animals. (10th edtn) 449-466.
- Martin JDW, Mayers PA, Rodwell VW, Granner DK (1985) Harper's Review of Biochemistry. (20th edtn) Lange Medical Publications, California: 651-660.
- Streeten Dh, Williams EMV (1952) Loss of cellular potassium as a cause of intestinal paralysis in dogs. J Physiol 118: 149-170.
- Indrayan AK, Sharma S, Durgapal D, Kumar N, Kumar M (2005) Determination of nutritive value and analysis of mineral elements for some medicinally valued plants from Uttaranchal. Curr Sci 89: 1252-1255.
- Soetan KO, Olaiya CO, Oyewole OE (2010) The importance of mineral elements for humans, domestic animals and plants: A review. Afr J Food Sci 4: 200-222.
- Chatterjee MN, Shinde R (1995) Text book of medical biochemistry. (2nd edtn), Jaypee Brother Medical Pub Ltd, New Delhi, India, 811-846.
- Hambiadge M (2000) Human Zinc deficiency. Tour of Nutrition Denver 130: 1344-1349.
- Atukorala TMS, Waidyanatha US (1987) Zinc and copper content of some common foods. J Nat Sci Coun Sri Lanka 15: 61-69.
- Arinola OG (2008) Essential trace elements and metal binding proteins in Nigerian consumers of alcoholic beverages. Pak J Nutr 7: 763-765.
- Malhotra VK (1998) Biochemistry for Students (10th edtn) Jaypee Brothers Medical Publishers (P) Ltd, New Delhi, India.
- Gebremedhin S, Enquselassie F, Umeta M (2011) Prevalence of prenatal zinc deficiency and its association with socio-demographic, dietary and health care related factors in rural Sidama, Southern Ethiopia: a cross-sectional study. BMC Public Health 11: 898.
- Prasad AS (1981) Symposia from the XII International Congress on Nutrition. Prog Clin Biol Res 77: 172-177.
- Larkin EC, Rao CA (1990) Importance of fetal and neonatal iron; adequacy for normal development of central nervous system. Brain behaviour and iron in the infant diet 43-57.
- Tan JC, Burns DL, Jones HR (2006) Severe ataxia, myelopathy, and peripheral neuropathy due to acquired copper deficiency in a patient with history of gastrectomy. JPEN J Parenter Enteral Nutr 30: 446-450.
- Sigel H (1978) Metals in Biological Systems. Marcel Dekker, New York 7: 241-275.
- Reddy MB, Chidambaram MV, Bates GW (1987) Iron Transport in Microbes, Plants and Animals. VCH New York.
- FAO/WHO (1984) Contaminants. In codex Alimentarius. (1st edtn) FAO/WHO codex Commissopn, Rome 17: 1-33.
- Kaplan LA, Pesce AJ, Kazmierczak SC (2003) Clinical Chemistry, Theory, Analysis, Correlation. (4th edtn) Mosby, London 503-509.
- Saner G, yüksel T, Gürson CT (1980) Effect of chromium on insulin secretion and glucose removal rate in the newborn. Am J Clin Nutr 33: 232-235.
- Frieden E (1984) Biochemistry of the essential ultratrace elements. Plenum press, New York 1-15.

34. Zayed AM, Terry N (2003) Chromium in the environment: factors affecting biological remediation. *Plant Soil* 249: 139-156.
35. Angard S (1980) Chromium, in *Metals in the Environment*. Academic Press Inc, London 111-132.
36. Cobanoglu U, Demir H, Sayir F, Duran M, Mergan D (2010) Some mineral, trace element and heavy metal concentrations in lung cancer. *Asian Pac J Cancer Prev* 11: 1383-1388.
37. Chandra RK (1990) Micronutrients and immune functions. An overview. *Ann N Y Acad Sci* 587: 9-16.
38. WHO (2005) *Quality Control Methods for Medicinal Plant Materials*. Revised Geneva Wong MK Tan P Wee.
39. McGrath SP, Smith S (1990) Chromium and Nickel in heavy metals in soils. In Alloway, B. J. Blackie, Glasgow 40-80.
40. Onianwa PC, et al. (2000) Cadmium and Nickel composition of Nigerian foods. *J Food Compos Anal* 13: 961-969.
41. Huheey JE, Keiter EA, Keiter RL (1993) *Inorganic Chemistry: Principles of Structure and Reactivity*, Harper Collins College Publishers, New York: 943-949.
42. Pendias AK, Pendias H (1992) *Trace Elements in Soils and Plants*. (2nd edn), CRC Press, Boca Raton, FL, USA 365.
43. Weber G, Konieczny A, ski P (2003) Speciation of Mg, Mn and Zn in extracts of medicinal plants. *Anal Bioanal Chem* 375: 1067-1073.
44. Debrah SK, Ayiyor JE, Denutsui D, Buah A, Nuviadenu C, et al. (2011) Elemental evaluation of some herbal plants used in Ghana using INAA. *Der Pharma Chemica* 3: 202-207.
45. Lokhande R, Singare P, Andhale M (2010) Study of mineral content of some Ayurvedic Indian medicinal plants by Neutron Activation Analysis and AAS techniques. *Health Sci J* 4: 163.
46. Ajasa A, Bello MO, Ibrahim AO, Ogunwande IA, Olawore NO (2004) Heavy trace metals and macronutrients status in herbal plants of Nigeria. *Food Chem* 85: 67-71.
47. Kolasani A, Xu H, Millikan M (2011) Evaluation of mineral content of Chinese medicinal herbs used to improve kidney function with chemometrics. *Food Chem* 127: 1465-1471.
48. Choudhury RP, Garg AN (2007) Variation in essential, trace and toxic elemental contents in *Murraya koenigii*—A spice and medicinal herb from different Indian states. *Food Chem* 104: 1454-1463.
49. Başgel S, Erdemoğlu SB (2006) Determination of mineral and trace elements in some medicinal herbs and their infusions consumed in Turkey. *Sci Total Environ* 359: 82-89.
50. Bachheti RK, Rai I, Joshi A, Pandey DP, Sharma A (2012) Physico-chemical and elemental analysis of ash of some medicinal plants from garhwal region, uttarakhand, india by atomic absorption spectrophotometer (aas). *Int J Pharm Pharm Sci* 4: 359-362.
51. Khosravinasab Z, Safarnejad A (2011) Comparison of Biochemical Characterization in *Cuminum cyminum* and *Cuminum setifolium* Seeds. *IJSN* 2: 874-877.
52. Ata S, Farooq F, Javed S (2011) Elemental profile of 24 common medicinal plants of Pakistan and its direct link with traditional uses. *J Med Plant Res* 5: 6164-6168.
53. Sandu DS, Gergen I, Trasca TI, Harmanescu M, Stef L, et al. (2010) Screening of 33 Medicinal Plants for the Microelements Content. *Scientific Papers: Animal Science and Biotechnologies* 43: 127-132.
54. Otunola GA, Oloyede OB, Oladiji AT, Afolayan AJ (2010) Comparative analysis of the chemical composition of three spices—*Allium sativum* L. *Zingiber officinale* Rosc. and *Capsicum frutescens* L. commonly consumed in Nigeria. *Afr J Biotechnol* 9: 6927-6931.
55. Padmini E, Valarmathi A, Rani MU (2010) Comparative analysis of chemical composition and antibacterial activities of and *Menthaspicata* *Camellia sinensis* *Asian J Exp Biol Sci* 1: 772-781.
56. Subramanian R, Gayathri S, Rathnav C, Raj V (2012) Analysis of mineral and heavy metals in some medicinal plants collected from local market. *Asian Pac J Trop Biomed* 2: 74-78.
57. Li JW, Fan LP, Ding SD, Ding XL (2007) Nutritional composition of five cultivars of Chinese jujube. *Food chem* 103: 454-460.
58. Kirmani MZ (2011) Determination of some toxic and essential trace metals in some medicinal and edible plants of Karachi city. *J Basic Appl Sci* 7: 89-95.
59. Dayanand B, Raghavan A, Khanum K, Singh B (2010) In vitro antioxidant and free radical scavenging activity of *Glycyrrhiza glabra* root extracts. *J herb med toxicol* 4: 97-102.
60. Fahima NK, Janati SSF, Feizy J (2012) Chemical Composition of Agriproduct Saffron (*Crocus sativus* L.) Petals and its considerations as animal feed gida 37: 197-201.
61. Kizil S, Hasimi N, Tolan V, Kilinc E, Yuksel U (2010) Mineral content, essential oil components and biological activity of two mentha species (*M. piperita* L., *M. spicata* L.). *Turk J Field Crops* 15: 148-153.
62. Akhtar S (1992) Chemical and nutritional evaluation of Genus *Acacia* of the family Leguminosae of pakistan.
63. Iqbal MS, Ghafoor A, Abbasi FM, Qureshi AS, Ahmad H (2013) Study of nutritional characteristics, mineral nutrients and agro-biodiversity in black cumin (*Nigella sativa* L.) genotypes from Pakistan. *Afr J Biotechnol* 10: 14757-14766.
64. Bello AG, Abdu I (2011) Nutrient and mineral elements levels in four indigenous tree seeds in Sokoto State, Nigeria. *J Plant Breed Crop Sci* 3: 396-400.
65. El-Safy FS, Salem RH, Abd El-Ghany ME (2012) Chemical and nutritional evaluation of different seed flours as novel sources of protein. *World J Dairy Food Sci* 7: 59-65.
66. Akinyede AI, Amoo IA (2009) Chemical and functional properties of full fat and defatted *Cassia fistula* seed flours. *Pak J Nutr* 8: 765-769.
67. Ghani A, Saeed S, Ali Z, Ahmad I, Ishtiaq M (2012) Heavy metals and nutritional composition of some selected herbal plants of Soon Valley, Khushab, Punjab, Pakistan. *Afr J Biotechnol* 11: 14064-14068.
68. Shehu K, Aliero AA (2010) Effects of Purple Blotch Infection on the Proximate and Mineral Contents of Onion Leaf. *Int J Pharma Sci Res* 1: 131-133.
69. Khan FA, Hussain I, Zahoor M, Rehman M (2011) Inorganic Constituent of Blue and White Flowering *Silybum Marianum* from Different Districts of Khyber pakhtoonkhawa, Pakistan. *Middle East J Sci Res* 7: 990-993.
70. Ullah Z, Baloch MK, Khader JA, Baloch IB, Ullah R, et al. (2013) Proximate and nutrient analysis of selected medicinal plants of Tank and South Waziristan area of Pakistan. *Afr J Pharm Pharmacol* 7: 179-184.
71. Kashmiri MA, Yasmin S, Ahmad M, Mohy-ud-Din A (2009) Characterization, compositional studies, antioxidant and antibacterial activities of seeds of *Abutilon indicum* and *Abutilon muticum* grown wild in Pakistan. *Acta Chim Slov* 56: 345-352.
72. Shad AA, Shah HU, Bakht J (2013) Ethnobotanical Assessment and Nutritive Potential of Wild Food Plants. *J Anim Plant Sci* 23: 92-97.
73. Oseni OA, Olarinoye CO, Amoo IA (2011) Studies on chemical compositions and functional properties of thorn apple (*Daturastramonium* L) Solanaceae. *Afr J Food Sci* 5: 40-44.
74. Rout OP, Acharya R, Gupta R, Mishra SK (2013) Analysis Of Some Essential Elements In Medicinal Plants Used In Ayurveda. *IJABPT* 4: 335-338.
75. Javid H, Khan AL, Rehman N, Khan F, Shinwari ZK, et al. (2009) Proximate and nutrient investigations of selected medicinal plants species of Pakistan. *Pak J Nutr* 8: 620-624.
76. Obumselu FO, Okerulu IO, Onwukeme VI, Onuegbu TU, Eze RC (2011) Phytochemical And Antibacterial Analysis Of The Leaf Extracts Of *Ricinus Communis*. *J Basic Phyl Res* 2: 68-72.
77. Khalil E, Esoh R, Rababah T, Almajwal AM, Alu'datt, MH (2012) Minerals, proximate composition and their correlations of medicinal plants from Jordan. *J Med Plants Res* 6: 5757-5762.
78. Nzikou JM, Matos L, Bouanga-Kalou G, Ndangui CB, Pambou-Tobi NPG, et al. (2009) Chemical composition on the seeds and oil of sesame (*Sesamum indicum* L.) grown in Congo-Brazzaville. *Adv J Food Sci Technol* 1: 6-11.
79. Bhowmik S, Chowdhury SD, Kabir MH, Ali MA (2008) Chemical composition of some medicinal plant products of indigenous origin. *Bangladesh Vet* 25: 32-39.
80. Shakira G, Latif A, Mirza IH, Nadeem MA (2011) Macro-Minerals Concentrations

- of Major Fodder Tree Leaves and Shrubs of District Chakwal, Pakistan. Pak J Nut 10: 480-484.
81. Khuda F, Iqbal Z, Zakiullah, Khan A, Nasir F, et al. (2012) Metal analysis, phytotoxic, insecticidal and cytotoxic activities of selected medicinal plants of Khyber Pakhtunkhwa. Pak J Pharm Sci 25: 51-58.
82. Hussain I, Khan FA, Khan N, Khan MA, Khan H (2011) Analysis of inorganic profile of *Tribulus terrestris*. Int J Phy Sci 6: 6147-6149.
83. Marcela F, Oana CT, Silvica A, Neculai M, Ioan B (2008) A new variety of *origanum vulgare* L.-denis, created at vrds bacau in ecologic agriculture condition, certified in 2007 year. J Plant Develop 15: 125-131.
84. Hussain J, Khan AL, Rehman N, Hamayun M, Shinwari ZK, Ullah W (2009) Assessment of herbal products and their composite medicinal plants through proximate and micronutrients analyses. J Med Plants Res 3: 1072-1077.
85. Haq F, Rehman S, Ahmad H, Iqbal Z, Ullah R (2012) Elemental analysis of *Paonia emodi* and *Punica granatum* by atomic absorption spectroscopy. Am J Biochem 2: 47-50.
86. Tewari LM, Tewari G, Nailwal T, Pangtey YPS (2009) Bark factors affecting the distribution of epiphytic ferns communities. Nat Sci 7: 76-80.
87. Jena VK, Gupta S, Patel KS, Patel SC (2013) Evaluating heavy metals Contents in medicinal plant *Mentha longifolia*. J Mater Environ Sci 4: 384-389.
88. Ashrafi EN, Vahdati K, Ebrahimzadeh H, Mirmasoumi M (2010) Analysis of in-vitro explants mineral contents to modify medium mineral composition for enhancing growth of Persian walnut (*Juglans regia* L.). J Food Agr Environ 8: 325-329.
89. Cosmulescu S, Botu M, Trandafir I (2010) Mineral Composition and Physical Characteristics of Walnut (*Juglans regia* L.) Cultivars Originating in Romania. Selcuk J Agr Food Sci /Selcuk Tarim ve Gida Bilimleri Dergisi 24: 33-37.
90. Shabbir A, Shahzad M, Arfat Y, Ali L, Aziz RS, Murtaza G, Waqar SA (2012) *Berberis lycium* Royle: A review of its traditional uses, phytochemistry and pharmacology. Afr J Pharm Pharmacol 6: 2346-2353.
91. Rahimi-Madiseh M, Heidarian E, Rafieian-kopaei M (2014) Biochemical components of *Berberis lycium* fruit and its effects on lipid profile in diabetic rats. J HerbMed Pharmacol 3: 15-19.
92. Krishnamurthy SR, Sarala P (2010) Proximate Nutritive Values and Mineral Components of *Withania Somnifera* (Linn.) Dunal. J Chem 7: 985-996.
93. Sah SP, Jha PK, Lamersdorf N (2002) Nutrient status of natural and healthy sissou forest and declining plantation sissou forest (*Dalbergiasissou*, Roxb.) in Nepal. J For Sci 48: 459-466.
94. Atangwho J, Ebong PE, Eyong, EU Williams IO, Eteng MU, et al. (2009) Comparative chemical composition of leaves of some antidiabetic medicinal plants: *Azadirachta indica*, *Vernonia amygdalina* and *Gongronemalati folium*. Afr J Biotech 8: 4685-4689.
95. Xie D, Zhong HY, Mo J, Li ZH, Cui T, Yi CP (2007) Nutritional and medicinal quality of pear juice: next hotspot? Food 1: 41-48.
96. Igwenyi IO, Eze CA, Azoro BN, Offor CE, Nwuke CP (2011) Proximate, Mineral and Amino Acid Compositions of *Irvignia gabonensis* and *Citrullus colocynthis* Used as Soup Thickener in South Easter Nigeria. Int J Biotechn Biochem 7: 493-499.
97. Oloyede OI (2005) Chemical profile of Unripe pulp of *Carica papaya*. Pak J Nutr 4: 379-381.
98. Akpambang VOE, Amoo IA, Izuagie AA (2008) Comparative compositional analysis on two varieties of melon (*ColocynthiscitrullusandCucumeropsis edulis*) and a variety of almond (*Prunusamygdalus*). Res J Agr Biol Sci 4: 639-642.
99. Khattak MI, Khattak MI (2011) Study of Heavy Trace Metals In Some Medicinal–Herbal Plants Of Pakistan. Pak J Bot 43: 2003-2009.
100. Kaymak HC, Guvenc I, Gurol A (2010) Elemental analysis of different radish (*Raphanus sativus* L.) Cultivars by using wavelength-dispersive-ray fluorescence spectrometry (WDXRF). Bulg J Agric Sci 16: 769-774.
101. Amoo IA, Atasié VN (2012) Nutritional and functional properties of *Tamarindusindicapulp* and *Zizyphus spina-christifruit* and seed. J Food Agri Environ 10: 16-19.
102. Jabeem T, Iqbal P, Khalil IA (1998) Amino Acid and Mineral Composition of Pea Cultivars Grpwn in Peshawar Pakistan. J Agri Res 9: 171-175.
103. Harmankaya M, Özcan MM, Karadas S, Ceyhan E (2010) Protein and mineral contents of pea (*Pisum sativum* L.) Genotypes grown in central Anatolian region of Turkey. South-west J Hortic Biol Environ 1: 159-165.
104. Hina B, Rizwani GH, Naseem S (2011) Determination of toxic metals in some herbal drugs through atomic absorption spectroscopy. Pak J Pharm Sci 24: 353-358.
105. Ameyaw Y, Barku VYA, Ayivor J, Forson A (2012) Phytochemical screening of some indigenous medicinal plant species used in the management of diabetes mellitus in Ghana. J Med Plants Res 6: 4573-4581.
106. Kawo AH, Abdullahi BA, Gaiya ZA, Halilu A, Dabai M, Dakare MA (2009) Preliminary Phytochemical Screening, Proximate And Elemental Composition Of *Moringa Oleifera* Lam Seed Powder Bajopas. Bayero J Pure appl Sci 2: 96-100.
107. Antia BS, Akpan EJ, Umoren OPA (2006) Nutritive and Anti-Nutritive Evaluation of Sweet Potatoes (*Ipomoea batatas*) Leaves. Pak J Nutr 5: 166-168.
108. Ghaly AE, Mahmoud N, Dave D (2012) Nutrient Composition of Dandelions and Its Potential as Human Food. Am J Biochem Biotech 8: 118-127.
109. Tiwari MR, Tamrakar NL, Shrestha HR, Shrestha BK, Jha RK (2006) Nutritive Value of Different Non-conventional Feeding Resources Found in Nepal. Green Field 4: 38-50.
110. Hussain I, Khan L (2010) Comparative Study on Heavy Metal Contents in *Taraxacum Officinale*. Int J Pharm Phytochem Res 2: 15-18.
111. Czarnowska K, Milewska A (2000) The Content of Heavy Metals in an Indicator Plant (*TaraxacumOfficinale*) in Warsaw Polish. J Environ Stud 9: 125-128.
112. Pachkore GL, Dhale DA (2012) Phytochemicals, Vitamins and Minerals Content of Three *Ocimum* Species. IJSSID 2: 201-207.
113. Ullah S, Khurram M, Ullah A, Rehman K, Khan FA (2013) Comparative Analyses of *OcimumsantumStem* and Leaves for Phytochemicals and Inorganic Constituents. Middle-East J Sci Res 13: 236-240.
114. Daniel, VN, Daniang IE, Nimyel ND (2011) Phytochemical Analysis And Mineral Elements Composition Of *Ocimum Basilicum* Obtained In Jos Metropolis, Plateau State, Nigeria. Int J Eng Tech 11: 161-165.
115. Saklani S, Chandra S, Badoni PP, Dogra S (2012) Antimicrobial activity, nutritional profile and phytochemical screening of wild edible fruit Of *Rubusellipticus*. Int J Med Arom Plants 2: 269-274.
116. Pirzada AJ, Shaikh W, Ghani KU, Laghari KA (2009) Study of Anti-Fungal Activity and Some Basic Elements of Medicinal Plant *Cressa Cretica* Linn Against Fungi Causing Skin Diseases. Res Jour (Sci Ser) 41: 15-20.
117. Skorbilowicz E (2009) Aquatic Plants as Bioindicators of Contamination of Upper Narew River And Some of Its Tributaries With Heavy Metals. Environ Prot Eng 35: 66-77.
118. Orisakwe OE, Nduka JK, Amadi CN, Dike DO, Bede O (2012) Heavy metals health risk assessment for population via consumption of food crops and fruits in Owerri, South Eastern, Nigeria. Chem Cent J 6: 77.
119. Ang LH, Ng LT (2000) Trace element concentration in mango (*Mangifera indica* L.), seedless guava (*Psidium guajava* L.) and papaya (*Carica papaya* L.) grown on agricultural and ex-mining lands of Bidor, Perak. Pertanika J Trop Agric Sci 23: 15-22.
120. Fatima N, Maqsood ZT, Khan B (2005) Study of some micronutrients in selected medicinal plants. Sci Iran 12: 269-273.
121. Jothivel S, Pillai SGP, Suburayan A (2011) Copper and Zinc Accumulation in few medicinal plants and Their Commercial Product in Tiruchirappalli Gandhi markwt, Tamilnadu, india. Afr J Basic Appl Sci 3: 223-227.
122. Prakash MMS, Kinthada PVS, Muralidhar NP (2011) Biologically estimation of heavy / toxic metals present in traditional medicinal plant – *Eclipta alba*. Int J Pharm Biomed Sci 2: 99-102.
123. Ghani A, Ali Z, Ishtiaq M, Maqboo M, Parveen S (2012) Estimation of macro and micro nutrients in some important medicinal plants of Soon Valley, District Khushab, Pakistan. Afr J Biotech 11: 14386-14391.
124. Chishti KA, Khan FA, Shah SMH (2011) Estimation Of Heavy Metals In The

- Seeds Of Blue And White Capitulum's Of *Silybum Marianum* Grown In Various Districts Of Pakistan. J Basic Appl Sci 7: 45-49.
125. Khan FA, Hussain I, Zahoor M, Rehman M (2011) Inorganic Constituent of Blue and White Flowering *Silybum Marianum* from Different Districts of Khyber Pakhtoonkhawa, Pakistan. Middle-East J Sci Res 7: 990-993.
126. Ighodaro OM (2012) Evaluation study on Nigerian species of *Musa paradisiaca* Peels. Researcher 4: 17-20.
127. Kalita M, Devi A (2012) Uptake of Metals by Four Commonly Available Plant Species Collected From Crude Oil Contaminated Sites of Lakowa Oil Field. Int J Agr Sci 2: 121-134.
128. Mahadkar S, Valvi S, Rathod V (2012) Nutritional assessment of some selected wild edible plants as a good source of mineral. Asian J Plant Sci Res 2: 468-472.
129. Navneet S, Tiwari MM, Chauhan A (2010) Elementals Profile Of Traditional Some Important Medicinal Plants of Uttarakhand State, India. Report and Opinion 2: 34-36.
130. Sun J, Liu SF, Zhang CS, Yu LN, Bi J, et al. (2012) Chemical composition and antioxidant activities of *Broussonetia papyrifera* fruits. PLoS One 7: e32021.
131. Tack FM, Verloo MG (1996) Metal contents in stinging nettle (*Urtica dioica* L.) as affected by soil characteristics. Sci Total Environ 192: 31-39.
132. Pinamonti F, Stringari G, Gasperi F, Zorzi G (1997) The use of compost: Its effect on heavy metals leaves in soil and plants. Resour Conserv Recy 21: 129-143.
133. Bamidele J, Akinyele OC, Akinkunmi KD (2012) Effect Of Electromagnetic Field On The Spoilage Fungi Of Some Selected Edible Fruits In Southwestern, Nigeria. J Microbiol Biotech Food Sci 2: 701-712.
134. Khan SA, Khan L, Hussain I (2008) Profile Of Heavy Metals In Selected Medicinal plants. Pak J Weed Sci Res 14: 101-110.
135. EsmatAbou-Arab A, AzzaAbou-Arab A, Abu-Salem MF (2010) Physico-chemical assessment of natural sweeteners steviosides produced from *Stevia rebaudiana bertonii* plant. Afr J Food Sci 4: 269-281.
136. Thakur S, Bawara B, Dubey A, Nandini D, Chauhan NS, et al. (2009) Effect of *Carum carvi* and *Curcuma longa* on hormonal and reproductive parameter of female rats. Int J Phytomed 1: 31-38.
137. Shahid M, Hussain F (2012) Chemical Composition and Mineral Contents of *Zingiber officinale* and *Alpinia allughas* (*Zingiberaceae*) Rhizomes. Int J Chem Biochem Sci 2: 101-104.
138. Ray JG, George J (2010) Zn in Tolerant Roadside plants in Relation to the metal in the soil in south India. American-Eurasian J Agric Environ Sci 9: 548-559.
139. Ashraf MA, Maah MJ (2011) Heavy metals accumulation in plants growing in ex tin mining, catchment. Int J Environ Sci Tech 8: 40-416.
140. Niamat R, Khan MA, Khan KY, Ahmad M, Ali B, et al. (2012) Element Content of Some Ethnomedicinal *Ziziphus* Linn. Species Using Atomic Absorption Spectroscopy Technique. J Appl Pharm Sci 2: 96-100.
141. Sekabira K, Oryem-Origa H, Mutumba G, Kakudidi E, Basamba TA (2011) Heavy metal phytoremediation by *Commelinabenghalensis* (L) and *Cynodon dactylon* (L) growing in Urban stream sediments. Int J Plant Physiol Biochem 3: 133-142.
142. Rathore M (2009) Nutrient content of important fruit trees from arid zone of Rajasthan. J Hortic For 1: 103-108.
143. Ameh GI, Nwammoh AO (2010) Phytochemical and Ethnobotanical Evaluation of Garlic Bulb (*Allium sativum* L.). Bio Res 8: 618-621.
144. Madejón P, Murillo JM, Marañón T, Cabrera F (2006) Bioaccumulation of trace elements in a wild grass three years after the Aznalcóllar mine spill (South Spain). Environ Monit Assess 114: 169-189.