

## Perceptions Regarding Hearing Impairment by Audiometry as Cognitive Domain Batteries?

Paul J. Regal<sup>1\*</sup>, Peter Lange<sup>2</sup>

<sup>1</sup>Department of Geriatric Medicine, University of Newcastle, Lake Haven, New South Wales, Australia; <sup>2</sup>Department of Geriatric Medicine, Geriatrician University of Melbourne, Royal Melbourne Hospital, Melbourne, Australia

### ABSTRACT

This article is about hearing acuity in a memory clinic. This memory clinic is typical of almost all clinics in that background noise is low and there are no alarms or signals from monitors as in emergency departments. We performed pure tone audiometry with the Siemen's hear check Screener (Siemens AG, Munich) at three intensities and two frequencies (1000 and 3000 Htz). We analysed the best ear for hearing acuity.

**Keywords:** Pure tone audiometry; Hearing; Dementia; Delirium

### DESCRIPTION

This article [1] is about hearing acuity in a memory clinic. This memory clinic is typical of almost all clinics in that background noise is low and there are no alarms or signals from monitors as in emergency departments. We performed pure tone audiometry with the Siemen's Hear Check Screener (Siemens AG, Munich) at three intensities and two frequencies (1000 and 3000 Htz). We analysed the best ear for hearing acuity. The best possible score is 6/6 tones and the worst is 0/6. Hearing loss was trichotomized severe as acuity 0/6-2/6, intermediate as 3/6 to 5/6 and normal as 6/6. Global cognition was assessed by two tests: Mini-Mental State Examination (MMSE) and Montreal Cognitive Assessment (MoCA). Cognitive domains were assessed using multiple tests for attention, executive function, language, memory, and visual-spatial function. Each domain had an average of four tests. We recruited 124 subjects whose median age was 82. The 31 subjects with severe hearing loss were on average 4 years older than the intermediate hearing group and 11.4 years older than those with normal hearing. The severe group also had twice the frequency of low instrumental activities of daily living (Nottingham IADL < 6/22), twice the prevalence of dementia as the intermediate group and four-fold more dementia than the normal group. Portable amplifiers with headphones were used on 66%, 15% and 0% of the three groups. We used Bonferroni adjustment of P values from  $P < 0.05$  to  $P < 0.0014$ . Best hearing

acuity was highly correlated with 13 of 36 scores including 11 of 31 (35%) cognitive measures. Yet cognitive domains scores were better correlated with other cognitive domains than were hearing acuity scores. We compared our results to ten other studies [2-12]. We did not use Z scores to combine cognitive tests but this was similar to other studies: none of the ten studies used Z scores to combine cognitive tests.

One weakness of the study is that portable amplifiers were not used in 100% of the severely impaired group. Another weakness was the convenience sample from a single centre. A strength of the study is exclusion of delirium by the Delirium Index. Strength is the Bonferroni adjustment of P values for multiple comparisons. An additional strength is use of open-access cognitive tests which can be used by all clinicians and not just registered psychologists.

In conclusion, this article supports portable pure tone audiometry in older outpatients with cognitive complaints.

### REFERENCES

1. Regal P, Lange P. Is hearing impairment by audiometry as much a cognitive score as cognitive domain batteries. *European Geriatric Medicine*. 2020.
2. Lin FR, Yaffe K, Xia J, Qian LX, Luigi F, Tamara BH, et al. Hearing loss and cognitive decline in older adults. *JAMA Intern Med*. 2013; 293:299.

**Correspondence to:** Paul J Regal, Department of Geriatric Medicine, University of Newcastle, Lake Haven, New South Wales, E-mail: regalelderlymedicine@gmail.com

**Received:** March 1, 2021, **Accepted:** March 14, 2021, **Published:** March 22, 2021

**Citation:** Regal PJ, Peter L (2021) Perceptions Regarding Hearing Impairment by Audiometry as Cognitive Domain Batteries? *J Phonet Audiol*. 6: 157

**Copyright:** © 2021 Regal PJ, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

3. Na W, Kim G, Kim G, Han W, Kim J. Effects of hearing loss on speech recognition under distracting conditions and working memory in the elderly. *Clin Interv Aging*. 2017; 1175-1181.
4. Amieva H, Ouvrard C, Giulioli C, Meillon C, Rullier L, Dartigues JF. Self-reported hearing loss, hearing aids and cognitive decline in elderly adults: A 25-year study. *J Am Geriatr Soc*. 2015; 2099-2104.
5. MacDonald AA, Joyson A, Lee R, Seymour DG, Soiza RL. The effect of hearing augmentation on cognitive assessment scales at admission to hospital. *Am J Geriatr Psychiatry*. 2012; 355-361.
6. Lin FR, Metter EJ, O'Brien RJ, Resnick SM, Zonderman AB, Ferrucci L. Hearing loss and incident dementia. *Arch Neurology*. 2011; 214-220.
7. Davies HR, Cadar D, Herbert A, Orrell M, Steptoe A. Hearing impairment and incident dementia: Findings from the English Longitudinal Study of Aging. *J Am Geriatr Soc*. 2017; 2074-2081.
8. Reed NS, Betz J, Kendig N, Korczak M, Lin FR. Personal sound amplification products vs. a conventional hearing aid for speech understanding in noise. *JAMA*. 2017; 89-90.
9. Volter C, Gotze L, Dazert S, Falkenstein M, Thomas JP. Can cochlear implantation improve neurocognition in the aging population? *Clin Interv Aging*. 2018; 701-712.
10. Strawbridge WJ, Walhagen MI. Simple tests compare well with a hand-held audiometer for hearing loss screening in primary care. *J Am Geriatr Soc*. 2017; 2282-2284.
11. Kiely KM, Gopinath B, Mitchell P, Luszcz M, Anstey KJ. Cognitive, health, and sociodemographic predictors of longitudinal decline in hearing acuity among older adults. *J Gerontol Med Sciences*. 2012; 997-1003.
12. Lim MY, Loo JH. Screening an elderly hearing impaired population for mild cognitive impairment using Mini-Mental State Examination (MMSE) and Montreal Cognitive Assessment (MoCA). *Int J Geriatr Psychiatry*. 2018; 972-979.