

Specific IgE Antibodies in Infant with Cow's Milk Protein Allergy

Hatem M El-Sebay¹, Eman AE Badr^{1*}, Yasser El-Ghobashi¹, Marwa MI Khalil¹ and Ghada M El-Mashad²

¹Medical Biochemistry Department, Faculty of Medicine- Menofia University, Egypt

²Pediatric Department, Faculty of Medicine- Menofia University, Egypt

Abstract

Background: Allergy to cow's milk is the most common food allergy in infant and young children. Symptoms of a milk allergy reaction can be mild like rash or severe, such as anaphylaxis. The aim of this study was to assess the value lactalbumin and lactoglobulin specific IgE in diagnosis of cow's milk protein allergy.

Design and Methods: This study was carried on 70 subjects classified into the following groups: Group 1: Include 50 patients with suspected cow milk protein allergy. Diagnosed by presence of chronic diarrhea with history of recent introduction of cow milk and positive elimination test. Group 2: Include 20 ages and sex matched apparently healthy subjects, their ages were ranged between (8-18) months. All individuals included in this study were subjected to full history taking, clinical examination, complete blood count and determination of serum total IgE, lactoglobulin and lactalbumin specific IgE which were carried out by ELISA technique.

Results: The diagnostic accuracy of lactoglobulin IgE in diagnosis of protein allergy was (84%), with sensitivity (78%), specificity (100%), positive predictive value (100%) and negative predictive value (65%) at cutoff point of 0.345 IU/ml. While, the diagnostic accuracy of lactalbumin IgE in diagnosis of protein allergy was (83%), with sensitivity of (84%), specificity (80%), positive predictive value (91%) and negative predictive value (67%) at cutoff point of 0.335 IU/ml.

Conclusions: Lactalbumin and lactoglobulin specific IgE assay are important in diagnosis of cow milk protein allergy and their combination may give better diagnostic accuracy than total IgE assay.

Keywords: Milk allergy; Lactoalbumin specific IgE; Lactoglobulin specific IgE

Introduction

Food allergy is defined as an adverse health effect arising from a specific immune response that occurs reproducibly following exposure to a given food [1].

Allergy to cow's milk protein (CMP) is an immunologically mediated reaction to one or more of the milk proteins. These proteins include caseins and whey proteins [2].

The immunological mechanisms that lead to the development of cow's milk protein allergy is not still clarified. There are different mechanisms that contribute to the pathogenesis and the main two described mechanisms at the basis of this disease refer to immediate or delayed response [3].

Cow milk allergy can be further split into IgE and non-IgE (mostly cellular) mediated. While IgE-mediated reactions are well recognized with validated diagnostic tests, the non IgE-mediated immune reactions are not so well defined and more difficult to recognize [4].

IgE-mediated allergy is associated with atopic manifestations such as urticaria, angioedema, vomiting, diarrhea, eczema, rhinitis, and anaphylaxis. Non-IgE-mediated allergy is associated with symptoms including gastro-esophageal reflux, vomiting, constipation, hemosiderosis, malabsorption, villous atrophy, eosinophilic proctocolitis, enterocolitis, and eosinophilic esophagitis [5].

The prevalence of food allergy (FA) varies from 6% to 8% in children, and it is currently increasing in many countries. Among all food allergens, cow's milk is one of the most common and often the first food introduced in the infant diet, even during breastfeeding. Cow's milk allergy (CMA) affects approximately 2.5% of children and may occur early in life, even during the neonatal period [6].

There is no one symptom pathognomonic of CMPA; it can present with an array of symptoms affecting different organ systems typically the skin, respiratory, and gastrointestinal tracts with many infants developing symptoms in more than one organ system [7].

There are a number of confirmatory tests which can add value when diagnosing CMPA [8].

Specific IgE testing helps to confirm diagnosis in IgE-mediated allergy, and prick tests can be used to add value to the diagnosis [5].

The aim of this study was to assess the value of serum levels of total IgE, lactalbumin specific IgE and lactoglobulin specific IgE in patient suspected to have milk protein allergy.

Subjects and Methods

Subjects

This study was carried on 70 subjects, 50 patients with milk protein intolerance and 20 apparently health persons. They were 31 females and 39 males with age ranging from (8-18) month. The patients were attendants of out-patients clinic and inpatient of pediatrics Department,

***Corresponding author:** Eman AE Badr, Medical Biochemistry Department, Faculty of Medicine- Menofia University, ShebinElkom, Menoufia, Egypt, Tel: 01090365171; E-mail: ebadr2014@gmail.com

Received January 28, 2015; **Accepted** February 24, 2015; **Published** March 03, 2015

Citation: El-Sebay HM, Badr EAE, El-Ghobashi Y, Khalil MM, El-Mashad GM (2015) Specific IgE Antibodies in Infant with Cow's Milk Protein Allergy. J Nutr Food Sci 5: 350. doi:10.4172/2155-9600.1000350

Copyright: © 2015 El-Sebay HM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Menofia University Hospital during the period from April 2013 to January 2014. They were classified into the following groups:

Group 1: Include 50 patients with suspected cow milk protein intolerance. They were 21 females and 29 males with mean age 11.36 ± 3.46 months. Diagnosed by presence of chronic diarrhea with history of recent introduction of cow milk and positive elimination test.

Group 2: Include 20 ages and sex matched apparently healthy subjects. They were 10 females and 10 males with mean age 10.60 ± 3.56 months.

After approval of the local ethical committee and informed consent from each one, patients who were selected scheduled to undergo a sheet was taken to all patients subjected.

Methods

All patients were subjected to the following:

- **History taking including: History of the disease:** Onset, duration, presence of prior episodes of diarrhea and its association with introduction of cow milk and history of breast feeding. Family history of milk allergy or any other type of food allergies.
- **Complete clinical examination:** Measurement of the weight and height of the infant to know if the infant growth is retarded or not. Search for signs of dehydration (as sunken eyes, thirst and delay in return of abdominal skin fold) to assess the severity of diarrhea and vomiting. Examination of the skin for any urticarial rash. Auscultation of the chest wheezes to know if the respiratory tract is involved or not.
- Laboratory investigations were measured for both patients and controls including: Complete blood picture, Serum total IgE, lactalbumin specific IgE, lactoglobulin specific IgE, total proteins and albumin levels.

Samples collection

Five milliliters (ml) of venous blood were taken from each subject and divided as follows: 2 ml were put immediately in an EDTA tube for complete blood count (CBC), while the remaining 3 ml were put in a plain tube, left to clot for 30 minutes at room temperature then subjected to centrifugation for 10 minutes at 4000 rotation per minute (rpm) and the serum obtained was put in several aliquots, stored at -80 c until the time of assay.

Assay methods: Complete blood picture was measured with Pentra-80 automated blood counter (ABX- France -Rue du Caducee-Paris Euromedecine-BP-7290.34184 Montpellier-Cedex 4.)

- Serum total IgE was determined using solid phase enzyme-linked immunosorbent assay [9]. The kits provided by Chemux Bioscience, USA.
- Lactalbumin and lactoglobulin specific IgE was determined using a cellulose disc-based enzyme allergosorbent test (EAST) [10]. The kit provided by RIDASCREEN Germany.
- Colometric determination of total plasma proteins [11] and albumin level [12] Diamond Egypt.

Statistical analysis

The data collected were tabulated & analyzed by SPSS (statistical package for the social science software) statistical package version

20 on IBM compatible computer. Quantitative data were expressed as mean ± standard deviation (X ± SD) and analyzed by applying t test for comparison between two groups of normally distributed variables, while for comparison between two groups of not normally distributed variables Mann-Whitney Test was applied. Qualitative data were expressed as number and percentage (No & %) and analyzed by applying chi-square test and for 2x2 table and one cell has expected number less than 5 fisher's exact test was applied. Spearman correlation was used for no normally distributed quantitative variables or when one of the variables is qualitative. ROC curve was used to determine cutoff points, sensitivity and specificity for quantitative variables of interest.

Results

The result of the present study are represented in Tables 1-6.

The results show no significant statistical difference among the two studied groups as regards age and gender distribution (not shown).

There was a significant statistical difference between the studied groups regarding family history, immediate, respiratory and skin symptoms. While non-significant difference as regarding breast feeding (Table 1).

There was a significant increase of WBCs count, eosinophil % total IgE, lactalbumin specific IgE and lactoglobulinspecific IgE in group I when compared with group II. Also it shows significant decrease of hemoglobin levels and serum total proteins in group I when compared with group II and non significant statistical difference regarding other parameters (Table 2).

The diagnostic accuracy of total IgE in diagnosis of protein allergy was (59%), with sensitivity of (44%), specificity (95%), positive predictive value (96%) and negative predictive value (40%) at cutoff point of 50.05 IU/L. The diagnostic accuracy of lactoglobulin IgE in diagnosis of protein allergy was (84%), with sensitivity (78%), specificity (100%), positive predictive value (100%) and negative predictive value (65%) at cutoff point of 0.345 IU/ml. While, the diagnostic accuracy of lacto albumin IgE in diagnosis of protein allergy was (83%), with sensitivity of (84%), specificity (80%), positive predictive value (91%)

History	Studied Groups				Fisher's Exact test	P value
	Cases (n=50)		Control (n=20)			
	NO.	%	NO.	%		
Family history:						
Present	20	40	0	0	9.33	<0.01
Absent	30	60	20	100		
Breast feeding :						
Yes	26	52	11	55	0.05*	>0.05
No	24	48	9	45		
Immediate symptoms:						
Present	24	48	0	0	12.56	<0.001
Absent	26	52	20	100		
Respiratory symptoms:						
Present	18	36	0	0	7.9	<0.01
Absent	32	64	20	100		
Skin symptoms:						
Present	27	54	0	0	15.38	<0.001
Absent	23	46	20	100		

* χ² test

Table 1: Statistical comparison of history and clinical data suggestive of CMA among the studied groups.

Laboratory parameters	Studied groups		Test of significance	P value
	Cases (n=50)	Controls (n=20)		
Hb%(gm/dl):	13.10 ± 1.18	11.72 ± 1.28	t=4.13	<0.001
RBCs count(×10 ⁶ /L):	4.88 ± 0.32	4.72 ± 0.34	t=1.80	>0.05
Platelets count(×10 ³ /L):	344.70 ± 72.41	326.40 ± 84.79	t=0.84	>0.05
WBCs count(×10 ³ /L):	5.74 ± 1.20	7.05 ± 2.37	t=3.03	<0.01
Neutrophils%	63.30 ± 2.07	62.64 ± 3.09	t=0.88	>0.05
Lymphocytes%	28.06 ± 3.89	29.15 ± 2.25	t=1.17	>0.05
Monocytes%	3.44 ± 1.0751	3.05 ± 0.76	t=1.71	>0.05
Eosinophil%	4.98 ± 4.05	1.60 ± 1.05	U=3.96	<0.001
Basophils%	0.50 ± 0.51	0.65 ± 0.49	U=1.13	>0.05
Total protein(gm/dl)	7.29 ± 1.31	8.41 ± 1.88	t=2.43	<0.05
Albumin(gm/dl):	3.71 ± 0.66	3.66 ± 0.41	t=0.37	>0.05
Total IgE(IU/ml):	86.53 ± 139.66	19.07 ± 16.75	U=2.62	<0.01
Lactoglobulin IgE (IU/ml):	1.04 ± 1.23	0.29 ± 0.05	U=4.91	<0.001
Lactalbumin IgE (IU/ml):	0.65 ± 0.40	0.28 ± 0.06	U=5.11	<0.001

t: t test, U: Mann-Whitney Test

Table 2: Statistical comparison of laboratory parameters among studied groups.

Cutoff point	Sensitivity	Specificity	Positive predictive value	Negative predictive value	Diagnostic accuracy
Total IgE 50.05 IU/ml	44%	95%	96%	40%	59%
Lactoglobulin specific IgE 0.345 IU/ml	78%	100%	100%	65%	84%
Lactalbumin specific IgE 0.335 IU/ml	84%	80%	91%	67%	83%

Table 3: Diagnostic validity of total IgE, lactoglobulin and lacto albumin specific IgE (IU/ml) in diagnosis of protein allergy cases.

Combinations of specific IgE antibodies	Sensitivity	Specificity	Positive predictive value	Negative predictive value	Diagnostic accuracy
Combined total IgE and lactoglobulin	88%	50%	81%	63%	77%
Combined total IgE and lactalbumin	92%	35%	78%	64%	76%
Combined lactoglobulin and lactalbumin	90%	80%	92%	76%	87%
Combined total IgE, lactoglobulin and lactalbumin	92%	35%	78%	64%	76%

Table 4: Diagnostic validity of combinations of specific IgE antibodies in diagnosis of protein allergy cases.

and negative predictive value (67%) at cutoff point of 0.335 IU/ml (Table 3).

The diagnostic accuracy of combined total IgE and lactoglobulin was 77% with sensitivity of 88%, specificity of 50%, positive predictive value of 81% and negative predictive value 63%, the diagnostic accuracy of combined total IgE and lactalbumin was 76% with sensitivity of 92%, specificity of 35%, positive predictive value of 78% and negative predictive value 64%, the diagnostic accuracy of combined lactoglobulin IgE and lactalbumin was 87% with sensitivity of 90%, specificity of 80%, positive predictive value of 92% and negative predictive value 76%. While, the diagnostic accuracy of combined total IgE, lactoglobulin IgE and lactalbumin was 76% with sensitivity of 92%, specificity of 35%, positive predictive value of 78% and negative predictive value 64% (Table 4).

There was a significant statistical difference between positive and negative cases of total IgE and lactoglobulin specific IgE in infant suggestive of cow milk allergy according to family history. While, There was a significant statistical difference between positive and negative cases of total IgE in infant suggestive of cow milk allergy according to immediate symptom and non-significant statistical difference according to other parameters (Table 5).

There was a significant statistical difference between positive and negative cases of total IgE in infant suggestive of cow milk allergy according to WBCs count, eosinophil %, lactoglobulin IgE and lactalbumin IgE and a significant statistical difference between positive and negative cases of lactoglobulin IgE in infant suggestive of cow milk allergy according to eosinophil % and total IgE, RBCs count and lactalbumin IgE. Also, there was significant statistical difference between positive and negative cases of lactalbumin IgE in infant suggestive of cow milk allergy according to eosinophil %, total IgE and lactoglobulin IgE while, other parameters show a non-significant statistical difference (Table 6).

Discussion

The prevalence of CMA in children living in the developed world is approximately 2 to 3%, making it the most common cause of food allergy in the pediatric population [13].

Specific IgE testing helps to confirm diagnosis in IgE-mediated allergy, and prick tests can be used to add value to the diagnosis. Vandenplas et al. [7], but a combination of the 2 tests is not necessary for the diagnostic workup.

This study assesses the value of lactoglobulin and lacto albumin specific IgE in diagnosis of CMPA.

In the present study neither the age nor the gender was significant. As the age ranges from 6 month to 18 month with mean of 11.36 month and male to female ratio was 1:3:1. This is agree with the studies of Castro et al., [6], Topal et al., [14], Van den Hogen et al., [15], Robert et al., [16].

In the present study the positive family history of allergic diseases was present in 40% of cases. It is also significant in relation to lactoglobulin specific IgE and total IgE.

This agrees with the study of Sirasuda et al., [17] who found that, fifty-two percent of parents had atopic diseases. While the study of Mowszet et al., [18] stated positive family history of allergy in only 11%.

A genetic basis for atopic disease is supported by twin studies which show that allergies such as asthma, eczema, and hay fever correlate more highly in monozygotic than dizygotic twins irrespective of whether the monozygotic twins were raised together or apart [5].

CMPA can induce a diverse range of symptoms of variable intensity in infants. It is helpful to differentiate between the "immediate" (early) reactions and "delayed" (late) reactions. Immediate reactions occur from minutes up to 2 hours after allergen ingestion and are more likely to be IgE mediated, whereas delayed reactions manifest up to 48 hours or even 1 week following ingestion. Combinations of immediate and delayed reactions to the same allergen may occur in the same patient [19].

In the current study the presence of respiratory symptoms present in 36% of cases. The immediate symptoms and the skin symptoms were present in 48% and 54% of cases respectively.

This is in accordance with the studies of Skripak et al., [20], Castro

History	Total IgE				LactoglobulinIgE				LactalbuminIgE				P value
	Positive (n=22)		Negative (n=28)		Positive (n=39)		Negative (n=11)		Positive (n=42)		Negative (n=8)		
	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	
Family history:													P1<0.001
Present	15	68.2	5	17.9	19	48.7	1	9.1	19	45.2	1	12.5	P2<0.05
Absent	7	31.8	23	82.1	20	51.3	10	90.9	23	54.8	7	87.5	P3>0.05
Breast feeding :													P1>0.05
Yes	11	50	26	53.6	20	51.3	6	54.5	22	52.4	4	50	P2>0.05
No	11	50	21	46.4	19	48.7	5	45.5	20	47.6	4	50	P3>0.05
Immediate symptoms:													P1< 0.01
Present	15	68.2	9	32.1	20	51.3	4	36.4	22	52.4	2	25	P2>0.05
Absent	7	31.8	19	67.9	19	48.7	7	63.6	20	47.6	6	75	P3>0.05
Respiratory symptoms:													P1>0.05
Present	10	45.5	8	28.6	14	35.9	4	36.4	15	35.7	3	37.5	P2>0.05
Absent	12	54.5	20	71.4	25	64.1	7	63.6	27	64.3	5	62.5	P3>0.05
Skin lesions:													P1>0.05
Present	14	63.6	13	46.4	22	56.4	5	45.5	25	59.5	2	25	P2>0.05
Absent	8	36.4	15	53.6	17	43.6	6	54.5	17	40.5	6	75	P3>0.05

P1 between total IgE positive and negative cases

P2 between lactoglobulinspecific IgE positive and negative cases

P3 between lactoalbuminspecific IgE positive and negative cases

Table 5: Comparison of family history and history of symptoms suggestive of CMA in infants with total IgE-positive and negative cases.

Laboratory parameters	Total IgE		Lactoglobulin IgE		Lactalbumin IgE		P value
	Positive	Negative	Positive	Negative	Positive	Negative	
	(n=22)	(n=28)	(n=39)	(n=11)	(n=42)	(n=8)	
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	
Hb% (gm/dl)	11.47 ± 1.16	11.91 ± 1.36	11.69 ± 1.16	11.81 ± 1.70	11.56 ± 1.21	12.07 ± 1.66	P1>0.05 P2>0.05 P3>0.05
RBCs count (109/L)	4.68 ± 0.33	4.75 ± 0.35	4.79 ± 0.35	4.52 ± 0.21	4.76 ± 0.33	4.54 ± 0.37	P1>0.05 P2<0.05 P3>0.05
WBCs count (×10 ⁹ /L)	8.78 ± 2.36	5.69 ± 1.24	7.33 ± 2.48	6.06 ± 1.68	7.22 ± 2.36	6.16 ± 2.39	P1<0.001 P2>0.05 P3>0.05
Platelets count (×10 ⁹ /L)	339.18 ± 71.00	316.35 ± 94.28	324.64 ± 83.46	332.64 ± 93.29	331.69 ± 83.95	298.63 ± 89.41	P1>0.05 P2>0.05 P3>0.05
Eosinophil%	8.68 ± 3.32	2.07 ± 1.15	5.87 ± 4.16	1.81 ± 0.87	5.33 ± 3.94	3.12 ± 4.42	P1<0.001 P2<0.001 P3<0.05
Total protein (gm/dl)	7.65 ± 1.49	7.00 ± 1.09	7.35 ± 1.38	7.09 ± 1.03	7.35 ± 1.38	6.98 ± 0.81	P1>0.05 P2>0.05 P3>0.05
Albumin (gm/dl)	3.62 ± 0.59	3.77 ± 0.72	3.72 ± 0.67	3.66 ± 0.70	3.68 ± 0.66	3.87 ± 0.72	P1>0.05 P2>0.05 P3>0.05
Lactoglobulin IgE(IU/ml)	1.56 ± 1.58	0.62 ± 0.62	-----	-----	1.15 ± 1.30	0.47 ± 0.44	P1<0.001 ----- P3<0.01
Lactalbumin IgE(IU/ml)	0.85 ± 0.44	0.49 ± 0.28	0.69 ± 0.36	0.51 ± 0.50	-----	-----	P1<0.001 P2<0.01 -----
Total IgE	-----	-----	106.75 ± 152.24	14.83 ± 16.07	93.93 ± 144.32	47.63 ± 111.74	----- P2<0.001 P3<0.01

Table 6: Comparison of laboratory investigations in infants with total IgE, lactoglobulin and lacto albumin specific IgE positive and negative cases.

et al., [5], Sirasuda et al., [17], Robert et al., [16] and Van den Hogen et al., [15] who stated that, the most common presenting symptoms of milk allergy were skin-related reaction. In contrast to the present study Merras et al., [21] found that skin symptoms were non-significant this may be because the study focused mainly on non IgE mediated milk allergy.

In the current study the history of breast feeding was 52% with non-significant relation to total IgE, lactoglobulin specific IgE and lactalbumin specific IgE. This is in accordance with the study of Skripak et al., [20]. In contrast the study of Sirasuda et al., [17] found that the mean age of ceasing breast feeding only was 1.9 month (0-10), while that of starting CM formula was 1.5 month (0-10). So, exclusive breastfeeding has been shown to be the best method to prevent allergy [5].

In the current study the WBCs and Eosinophil % were significantly increased in patients group in comparing to controls. WBCs count significantly increased in cases with positive total IgE. The Eosinophil % significantly increased in cases with positive total IgE, lactalbumin specific IgE and lactoglobulin specific IgE. This agrees with studies of Sirasuda et al., [17] and Ömeret al., [22].

In the current study the total IgE level was significantly increase in patients group in compare to controls with mean of 86.53+1.23 IU/ml. The cutoff value was 50.05 with sensitivity of 44% and specificity of 95%, positive predictive value of 96% and negative predictive value of 40% and diagnostic accuracy of 59%.

In contrast the study of Ahren et al., [23] stated that the mean of total IgE in baseline diagnosis was 436.9 ± 924.2 IU/ml this may be because their study population was 52 children with CMA who had at least two consecutive food challenge tests.

A positive test for specific IgE at the time of diagnosis predicts a longer period of intolerance as compared with those children who have negative tests [24].

In the present study the cutoff point of lactoglobulin specific IgE was 0.345 IU/ml with sensitivity of 78% and specificity of 100%, the positive predictive value is 100%, the negative predictive value is 65% and the diagnostic accuracy is 84%. The cutoff point of lactalbumin specific IgE is 0.335 IU/ml with sensitivity of 84% and specificity of 80%, the positive predictive value is 91%, the negative predictive value is 67% and the diagnostic accuracy is 83%.

This agrees with the study of Skripak et al., [20], Ahren et al., [23], Corinne et al., [25] and Sirasuda et al., [17] in which IgE-mediated disease was defined as having a skin prick test with a wheal diameter 3 mm and/or a cm (cow milk) IgE 0.35 kU/L.

In contrast with the study of Lisa et al., (26) Sensitivity, specificity, and PPV, for α -lactoalbumin and β -lactoglobulin were poor. While the NPV for β - lactoglobulins IgE at 0.35 kU/L was 84.2% with AUC>90%

In contrast the study of Castro et al., [26] stated that the best specific IgE concentrations found were: 3.06 kUI/l for whole milk, 2.08 kUI/l for lactalbumin, 1.85 kUI/l for lactoglobulin and 1.47 kUI/l for casein this difference may be because this study included 123 children with confirmed CMA and the present study included only 50 children with suspected CMA.

Also, in contrast to the study of Federica et al. [27] in which the determination of cow's milk specific IgE was performed and values greater than 0.10 kUa/L were considered as positive.

In the present study the diagnostic validity of combined levels of total IgE and specific IgE showed that the combination of lactalbumin and lacto globulin specific IgE was the best as sensitivity was 90%, specificity was 80%, positive predictive value was 92%, negative predictive value was 76% and diagnostic accuracy was 87%.

It can be concluded that lactalbumin and lactoglobulin specific IgE are important in diagnosis of cow milk protein allergy and their combination may give better diagnostic accuracy. Total IgE has lesser diagnostic use in milk protein allergy. The level of lactoglobulin and lactalbumin specific IgE are related to family history, immediate symptoms and eosinophil %.

References

1. Koletzko, Niggemann, Arato, Dias, Heuschkel, et al. (2012) Diagnostic Approach and Management of Cow's-Milk Protein Allergy in Infants and Children: ESPGHAN GI Committee Practical Guidelines JPGN 55: 221-229.
2. Karabus SJ, Du Toit G (2012) IgE-mediated cow's milk protein allergy Current Allergy & Clinical Immunology 25: 4-9.
3. Vitaliti G, Cimino C, Coco A, Praticò AD, Lionetti E (2012) The immunopathogenesis of cow's milk protein allergy (CMPA). Ital J Pediatr 38: 35.
4. Benhamou AH, Schäppi Tempia MG, Belli DC, Eigenmann PA (2009) An overview of cow's milk allergy in children. Swiss Med Wkly 139: 300-307.
5. Vandenplas Y, Abuabat A, Al-Hammadi S, Aly GS, Miqdady MS, et al. (2014) Middle East Consensus Statement on the Prevention, Diagnosis, and Management of Cow's Milk Protein Allergy. Pediatr Gastroenterol Hepatol Nutr 17: 61-73.
6. Castro AP, Pastorino AC, Gushken AK, Kokron CM, Filho UD, et al. (2015) Establishing a cut-off for the serum levels of specific IgE to milk and its components for cow's milk allergy: Results from a specific population. Allergol Immunopathol (Madr) 43: 67-72.
7. Boyce JA, Assa'ad A, Burks AW, Jones SM, Sampson HA, et al. (2011) Guidelines for the diagnosis and management of food allergy in the United States: summary of the NIAID-sponsored expert panel report. J Am Diet Assoc 111: 17.
8. Costa AJ, Sarinho ES, Motta ME, Gomes PN, de Oliveira de Melo SM, et al. (2011) Allergy to cow's milk proteins: what contribution does hypersensitivity in skin tests have to this diagnosis? Pediatr Allergy Immunol 22: e133-138.
9. Ishizaka T (1982) IgE and mechanisms of IgE-mediated hypersensitivity. Ann Allergy 48: 313-319.
10. Kersten W, Von Wahl PG, Lang CE, Wenning J (2000) Recommendations for in vitro diagnosis of allergy. Allergologie Jg, 23: 304-307.
11. Young DS, Friedman RB (2001) Effects of Disease on Clinical Laboratory Tests, (4th edn), Vol. 1 and 2: AACCC Press.
12. Rodkey FL (1965) Direct Spectrophotometric Determination Of Albumin In Human Serum. Clin Chem 11: 478-487.
13. Lifschitz C, Szajewska H (2015) Cow's milk allergy: evidence-based diagnosis and management for the practitioner. Eur J Pediatr, 174: 141-150.
14. Topal E, EÄYritaÄY O, Arga M, SarÄ S, Poyraz A, et al. (2013) Eosinophilic esophagitis and anaphylaxis due to cow's milk in an infant. Turk J Pediatr 55: 222-225.
15. van den Hoogen SC, van de Pol AC, Meijer Y, Toet J, van Klei C, et al. (2014) Suspected cow's milk allergy in everyday general practice: a retrospective cohort study on health care burden and guideline adherence. BMC Res Notes 7: 507.
16. Robert A, Wood, Scott H, Sicherer, Brian P, et al. (2013) The natural history of milk allergy in an observational cohort. J. Allergy Clin. 131: 805-812.
17. Sommanus S, Kerddonfak S, Kamchaisatian W, Vilaiyuk S, Sasisakulporn C, et al. (2014) Cow's milk protein allergy: immunological response in children with cow's milk protein tolerance. Asian Pac J Allergy Immunol 32: 171-177.
18. Mowszet K, Matusiewicz K, IwaÄczak B (2014) Value of the atopy patch test in the diagnosis of food allergy in children with gastrointestinal symptoms. Adv Clin Exp Med 23: 403-409.

19. Shek LP, Bardina L, Castro R, Sampson HA, Beyer K (2005) Humoral and cellular responses to cow milk proteins in patients with milk-induced IgE-mediated and non-IgE-mediated disorders. *Allergy* 60: 912-919.
20. Skripak JM, Matsui EC, Mudd K, Wood RA (2007) The natural history of IgE-mediated cow's milk allergy. *J Allergy Clin Immunol* 120: 1172-1177.
21. Merras-Salmio L, Kolho KL, Peikonen AS, Kuitunen M, Mäkelä MJ, et al. (2014) Markers of gut mucosal inflammation and cow's milk specific immunoglobulins in non-IgE cow's milk allergy. *Clin Transl Allergy* 4: 8.
22. BeÄyer OF, Sancak S, Erkan T, Kutlu T, CokuÄyraÄY H, et al. (2014) Can Fecal Calprotectin Level Be Used as a Markers of Inflammation in the Diagnosis and Follow-Up of Cow's Milk Protein Allergy? *Allergy Asthma Immunol Res* 6: 33-38.
23. Ahrens B, Lopes de Oliveira LC, Grabenhenrich L, Schulz G, Niggemann B, et al. (2012) Individual cow's milk allergens as prognostic markers for tolerance development? *Clin Exp Allergy* 42: 1630-1637.
24. Halcken S (2003) Early sensitisation and development of allergic airway disease - risk factors and predictors. *Paediatr Respir Rev* 4: 128-134.
25. Keet CA, Frischmeyer-Guerrero PA, Thyagarajan A, Schroeder JT, Hamilton RG, et al. (2012) The safety and efficacy of sublingual and oral immunotherapy for milk allergy. *J Allergy Clin Immunol* 129: 448-455, 455.
26. Bartnikas LM, Sheehan WJ, Hoffman EB, Permaul P, Dioun AF, et al. (2012) Predicting food challenge outcomes for baked milk: role of specific IgE and skin prick testing. *Ann Allergy Asthma Immunol* 109: 309-313.
27. Bellini F, Ricci G, Dondi A, Piccinno V, Angelini F, et al. (2011) End point prick test: could this new test be used to predict the outcome of oral food challenge in children with cow's milk allergy? *Ital J Pediatr* 37: 52.