

Study of the Intestinal Absorption Characteristics of Curcumin *In Vivo* and *In Vitro*

Xue M¹, Cheng Y², Xu L² and Zhang L^{2*}

¹College of Basic Sciences, Nanjing University of Chinese Medicine, China

²Department of Pharmacology, Nanjing University of Medicine, Nanjing 210046, China

*Corresponding author: Zhang L, Department of Pharmacology, Nanjing University of Chinese Medicine, Nanjing 210046, China, Tel: +86-25-85811724; E-mail: zhangl_1999@163.com

Received date: June 07, 2017; Accepted date: June 16, 2017; Published date: June 23, 2017

Copyright: © 2017 Xue M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Curcumin is the yellow pigment of turmeric. In addition to its positive safety profile, curcumin is reported to have beneficial pharmacologic effects as an antioxidant, antitumor, and anti-inflammatory agent, along with other promising pharmacologic effects on the cardiovascular and digestive systems. Curcumin is poorly absorbed, which limits its value in clinical application. In order to improve the poor bioavailability and enhance the pharmacologic action of curcumin, we studied its absorption mechanisms in an animal model *in vivo* and in a Caco-2 cell model *in vitro*. The absorption rates of curcumin at different concentrations in blank intestinal juice were not the same. The absorption rate of curcumin solution at a concentration of 5 µg/mL was the highest, followed by 10 µg/mL, and the minimum absorption occurred at 20 µg/mL. The absorption rate in the ileum decreased as the concentration of curcumin increased, which reminds us that absorption in the ileum does not result from simple passive diffusion but rather shows the characteristics of active transport. Curcumin may be a P-glycoprotein (P-gp) substrate and thus may be affected by P-gp efflux, and thus the addition of a P-gp inhibitor such as verapamil can promote the intestinal absorption of curcumin. A Caco-2 cell model was established to accurately study curcumin's absorption mechanisms. We found that, for curcumin in a 5 µg/mL solution, the Caco-2 cell monolayer transport was passive, and when the concentration was increased to 10 µg/mL efflux influenced the transport but not extensively. The transport mode of curcumin appears to be passive diffusion at concentrations <10 µg/mL, but at concentrations >10 µg/mL active transport is involved. In summary, curcumin is transported by a combination of passive diffusion and active transport, curcumin is a substrate for the intestinal transporter P-gp, and intestinal absorption of curcumin is regulated by intestinal P-gp transport.

Keywords: Curcumin; Intestinal absorption; P-glycoprotein; Caco-2 cell

Introduction

Curcumin, the yellow pigment of turmeric, is a natural active ingredient extracted from the root of turmeric or curcuma. It is soluble in methanol, ethanol, alkali, acetic acid, acetone, chloroform, and other organic solvents (Figure 1). Numerous studies have demonstrated its positive safety profile and beneficial pharmacologic effects as an antioxidant, its antitumor and anti-inflammatory activities, its free radical scavenging and antimicrobial effects, and many other pharmacologic effects on the cardiovascular and digestive systems [1]. In particular, recent findings suggest that curcumin is potentially a powerful tool to reverse cisplatin-induced toxicity, and it has attracted considerable interest over the last decades due to its beneficial effects for human health [2]. Studies of the absorption and metabolism of curcumin have been conducted since at least the 1980s [3]. Ravindranath studied the absorption and tissue distribution of curcumin in rats that were orally given 400 mg of curcumin. At 15 min to 24 h after oral administration of curcumin, only trace amounts of 5 mg·L⁻¹ were detected in the hepatic portal vein [4]. Eight volunteers were given 2 g curcumin, after which the content of curcumin in human serum was very low or undetectable; the peak concentration was only 0.006 ± 0.005 mg·L⁻¹; and at 3 h after administration, the serum concentration reached zero. Overall, curcumin is poorly

absorbed and has low bioavailability, both of which limit its clinical applications and therapeutic efficacy [5]. At present, the mechanism by which curcumin is absorbed *in vivo* is not clear. In order to find suitable methods to improve poor bioavailability and enhance its pharmacologic action, we studied the absorption mechanisms of curcumin.




Figure 1: Structure of curcumin.

In this research, we used an animal model *in vivo* and a Caco-2 cell model *in vitro* in order to study the absorption process and the characteristics of curcumin. At the same time, we wanted to provide a theoretical basis to solve the problem of the low absorption rate of curcumin via the oral route and to find a new way to improve the biological use of some drugs that are poorly absorbed in the gastrointestinal tract.

Materials and Methods

Drug and chemicals

Reference-grade curcumin (purity 99.8%) were purchased from Nanjing Zelang Medical Technology Co., Ltd. (ZL20090515A). Hank's buffered salt solution (HBSS) was prepared as follows: anhydrous calcium chloride (0.40 g), magnesium chloride hexahydrate (0.10 g), and magnesium sulfate heptahydrate (0.10 g) were dissolved in 1000 mL of distilled water, and then potassium chloride (0.40 g), potassium dihydrogen phosphate (0.06 g), sodium bicarbonate (0.35 g), water (1000 mL), disodium hydrogen phosphate (0.12 g), sodium chloride (8.00 g), and glucose (1.00 g) were added and mixed until completely dissolved. All other reagents were of High-performance Liquid Chromatography (HPLC) grade or were of the highest purity commercially available.

Experimental animals

Animal experiments in this study were used in compliance with standard ethical guidelines and were approved by Nanjing University of Chinese Medicine Animal Care and Use Committee. Forty male SD rats, weighing 220 ± 20 g, were purchased from Shanghai Silaike Experiment Animal Co., Ltd. (SCXK2007-0005; Shanghai, China) and were housed in a 23°C room with a 12 h/12 h light-dark cycle and were allowed to acclimate to the laboratory for 1 week before experiments.

Instrumentation and analytical conditions

The concentration of curcumin was analyzed with an LC-10A HPLC (Shimadzu Corporation, Kyoto, Japan) equipped with an SPD-10Avp UV detector, a quaternary pump, an online degasser, and an automatic plate sampler. Chromatographic separations were performed with a Dimma ODS-2 C18 column (150×4.6 mm $5 \mu\text{m}$ packing) with a column temperature of 35°C. The mobile phase was methanol: water (70:30). The flow-rate was 0.2 mL/min. The detection wavelength was 425 nm. Aliquots of 20 μL were injected into the HPLC system for analysis.

Reverted gut sac experiments *in vitro*

Precise amounts of curcumin were added to a 0.1% Tween 80 solution containing HBSS in order to make concentrations of 5, 10, and 20 g/mL HBSS solution. The rats were fasted 24 h (water was freely available) and were anesthetized by intraperitoneal injection of 20% urethane anaesthesia. First we fixed the animals' backs on the operating table, and then we made a 3-cm incision along the ventral midline and quickly removed a 20-cm-long segment of the ileum. Next, we removed the small intestine and immediately placed it in saline solution, washed the small intestine, and then transferred the intestinal segment to the HBSS fluid; surface fat and blood vessels were removed by stripping.

Then the ileum was divided into 2 sections, each portion of which was sealed at one end with a silk thread ligation. A fine glass rod with a diameter <5 mm of fine glass rod for outer intestine, the small intestinal mucosa of lateral outside, and the other end is tied to one end of the glass tube bulge. Blank HBSS buffer 0.5 mL injection; the gut sacs thus prepared were placed into the containers with HBSS buffer maintained at 37°C. HBSS buffer was placed in a 100-mL beaker, and 95% O₂ and 5% CO₂ were aspirated for 5 min, and then

the gut sac was placed into the oxygenated HBSS solution 100 mL containing 5, 10, or 20 g/mL of drugs of HBSS solution. 0.5 mL was taken out of the gut sac at 30 min, 60 min, 90 min, and 120 min; then we added an equivalent amount of fresh blank HBSS solution. The cumulative absorption amount per unit area (Q) was determined by the following formula:

$$Q_n = C \times V/S$$

Where n=the number; V=the solution volume (0.5 mL) in the intestine; C=sample concentration (g/mL), and S=intestinal segment area (cm²).

In situ intestinal perfusion experiment

Rats were fasted 16 h overnight. After anaesthesia, animals were placed for surgery, and the ileum was removed. Then glass tubes were inserted into both ends of the intestinal segments. The segments were fixed and ligated, cleaned with a 37°C saline flush, and the lengths of the intestinal segments were measured at the end of the test.

HBSS solutions containing 0.1% Tween 80 buffer solution were prepared to concentrations of 5 g/mL, 10 g/mL, and 20 g/mL curcumin solution as the test group. The ahead solution served as the blank control. The amount of 50 mL containing curcumin intestinal perfusate, adding a 100 mL flask circulation device. Researchers placed the flask in a 37°C water bath and opened the constant-flow pump.

The flow rate was adjusted to 0.2 mL/min, replaced the infusion vial and liquid vial and weighing every 15 min, then calculated the perfusate volume. Drug concentrations at each time point were determined by HPLC. The experiment lasted 120 min, when all the vials were collected for determination, and the rats were killed. The ileum cannula along the cut ends, gently on the droplet adding 37°C Krebs-Ringer solution, measuring intestinal length, measuring inner diameter with thread. Calculation of relevant data was calculated as follows:

$$K_a = \left(1 - \frac{C_{out} Q_{out}}{C_{in} Q_{in}}\right) \frac{Q}{V} \quad (1)$$

$$P_{app} = \frac{-Q \ln\left(\frac{C_{out} Q_{out}}{C_{in} Q_{in}}\right)}{2\pi r l} \quad (2)$$

Where K_a is absorption rate, P_{app} is apparent permeability coefficient, Q_{in} and Q_{out} are the volumes of intestinal fluid imported and exported (mL), respectively. C_{in} and C_{out} were the concentrations of curcumin in the intestinal fluid imported and exported ($\mu\text{g/mL}$), respectively. L and R were the length (cm) and the cross-section radius (cm) of the intestinal segment, Q was the perfusion rate (0.2 mL/min), t was the perfusion time (0.25 h), and V was the perfusion volume of the intestinal segment.

Effect of the P-gp inhibitor verapamil on the transport of curcumin

A solution of 5 $\mu\text{g/mL}$ of curcumin was used in the control group, and a 5 $\mu\text{g/mL}$ curcumin solution containing 500 $\mu\text{mol/L}$ verapamil hydrochloride was used in the experimental group. We conducted rat ileum perfusion experiments using the method just described, and both K_a and P_{app} values were calculated.

Study of the mechanism of curcumin activity in Caco-2 cell culture

Caco-2 cells from the Shanghai Cell Bank, Chinese Academy of Sciences were grown to <90% confluency in DMEM containing 10% FBS, 100 U/mL penicillin, 0.1 mg/mL streptomycin in a humidified atmosphere of 5% CO₂ in air maintained at 37°C. The cells were used from passage 50 to 90. After trypsinization, Caco-2 cells were seeded onto Millicell hanging cell culture inserts (Supplier, City, Country) in 24-well plastic plates at a density of 1 × 10⁵ cells/cm² for approximately 21 days.

The integrity of the cell monolayer was checked before and after each experiment by measuring transepithelial electrical resistance (TEER) using a volt-ohmmeter (EVOM; Millipore, Tokyo, Japan). Only monolayers with a TEER value >350 Ω·cm² were considered intact and suitable for transport studies. To obtain Caco-2 monolayers with various TEER values, the monolayers were treated with cytochalasin D (0.1 mM) from the apical side.

Transepithelial transport experiments

Curcumin was diluted to working concentrations of 5, 10, and 20 µg/mL, and we prepared a 500 M verapamil with HBSS solution containing 0.1% Tween 80. All experiments were performed in HBSS buffer. After washing the inserts 3 times with warm HBSS buffer for 30 min, researchers added transport medium containing curcumin (5, 10, or 20 µg/mL with 500 M verapamil) either to the apical or basolateral compartment (0.4 or 0.6 mL, respectively). After shaking at 55 rpm at 37°C, 100-µL samples were collected after 30, 60, 90, and 120 min from the receiver compartment then immediately replaced with an equal volume of prewarmed fresh HBSS (n=4). Simultaneous determinations of TEER values were made during the process of the experiments. Papp (apparent permeability coefficient) and ER (efflux ratios) were calculated:

$$P_{app} = Q / (t \cdot A \cdot C_0);$$

Q: Transport quantity in t (µg);

A: Surface area of the carbon film (cm²);

C₀: Initial concentration of the drug (µg/mL).

$$ER = P_{appB \rightarrow A} / P_{appA \rightarrow B}$$

P_{appB→A} Apparent permeability coefficient from B side to A side (cm/s);

P_{appA→B} Apparent permeability coefficient from A side to B side (cm/s).

Data analysis

The differences between the groups were analyzed using Student's t-test when only 2 groups were compared or one-way analysis of variance (ANOVA) when >2 groups were compared. P values <0.05 indicate statistical significance. All statistical analyses were performed using SPSS 17.0 software (SPSS, Chicago, IL, USA).

Results

HPLC method for curcumin assay

HPLC mapping was used to measure the intestinal absorption of curcumin in different concentrations. The curcumin standard curve is

shown by: $y = 31648x - 22253$, $R = 0.9996$. In the range of (0.5 to approximately 50 g/mL), the concentration of curcumin was in good linear relationship with the peak area. Compared with the initial value at 0 h, different concentrations of curcumin at different time points were not significantly different ($P > 0.05$). Compared with the initial concentration, the concentration at 2 h was 98%. The intestinal solution can maintain a steady state of curcumin, and the HPLC assay can quickly and accurately determine the concentrations of curcumin, which can be used for quantitative analysis.

Intestinal absorption rates of different concentrations of curcumin in reverted gut sac experiments

The absorption rates of curcumin at different concentrations were not the same. Results are shown in Figure 2. The absorption rate of the curcumin solution with a concentration of 5 µg/mL was the highest, followed by the second-highest absorption of the 10 µg/mL solutions, but the minimum absorption occurred with the 20 µg/mL solutions. The absorption rate of the ileum decreased as the concentration of curcumin increased. Absorption in the ileum is not the result of simple passive diffusion and instead shows the characteristics of active transport.




Figure 2: Absorption rate of curcumin in the ileum segment.

In situ intestinal perfusion experiment of different concentrations of curcumin in the presence of a P-gp substrate

When the concentration of curcumin increased from 5 µg/mL to 10 µg/mL, the K_a value decreased, and the K_a value continued to decrease at a concentration of 20 µg/mL. Similarly, the P_{app} value also decreased. The concentration continued to increase to 20 µg/mL when the P_{app} decreased slightly (Table 1).

Dose group	K_a	P_{app}
	($\times 10^{-2} \text{ min}^{-1}$)	($\times 10^{-3} \text{ cm} \times \text{min}^{-1}$)
20 µg/mL curcumin	11.92 ± 1.10	8.23 ± 0.29
10 µg/mL curcumin	15.99 ± 0.63	12.29 ± 3.51
5 µg/mL curcumin	20.22 ± 0.07	21.09 ± 0.42

Table 1: Absorption rate and apparent permeability coefficient of various concentrations of curcumin.

The experimental results show that the absorption rate (K_a) and apparent permeability coefficient (P_{app}) decreased when the concentration of curcumin was increased. We observed a high concentration of saturation phenomenon. When 5 $\mu\text{g}/\text{mL}$ of curcumin solution was added to the 5 M P-gp inhibitor verapamil, the K_a value was increased from $20.22 \pm 0.07 \times 10^{-2} \text{ min}^{-1}$ to $41.61 \pm 5.71 \times 10^{-2} \text{ min}^{-1}$ ($P < 0.01$), and the P_{app} value increased from $21.09 \pm 0.42 \times 10^{-3} \text{ cm} \times \text{min}^{-1}$ to $29.14 \pm 4.00 \times 10^{-3} \text{ cm} \times \text{min}^{-1}$ ($P < 0.05$). Results are shown in Figures 3 and 4. The effect of P-gp on the intestinal absorption of curcumin in the presence of verapamil is relatively large, and curcumin may be a substrate of P-gp and may be affected by the P-gp efflux function.

Transport across Caco-2 cells

The experimental results are shown in Tables 2 and 3. As the concentration of curcumin increased, TEER values gradually increased in a concentration-dependent manner: Following addition of 500 μM verapamil, the curcumin $P_{appA \rightarrow B}$ in the 10- $\mu\text{g}/\text{mL}$ and 20- $\mu\text{g}/\text{mL}$ solutions increased. There was no significant decrease in TEER. After verapamil was added to the 500- μM solution, the $P_{appA \rightarrow B}$ values between the 2 concentrations were almost equal and were quite close to the $P_{appA \rightarrow B}$ of 5 $\mu\text{g}/\text{mL}$. After adding P-gp inhibitors, the apparent permeability coefficient of curcumin $P_{appA \rightarrow B}$ value remained constant, indicating that passive diffusion was the main transport mechanism. For curcumin in a 5 $\mu\text{g}/\text{mL}$ solution, the Caco-2 cell monolayer transport mechanism is still passive transport; when the concentration rose to 10 $\mu\text{g}/\text{mL}$, the efflux exerted an influence on the transport, but the efflux effect was not very strong. The transport mode of curcumin involves passive diffusion at concentrations $< 10 \mu\text{g}/\text{mL}$, but when the concentration is $> 10 \mu\text{g}/\text{mL}$, active transport is involved. Overall, the transport mechanism of curcumin involves a combination of active and passive transport, and curcumin is a P glycoprotein substrate.




Figure 3: Absorption rate of various concentrations of curcumin.




Figure 4: Apparent permeability coefficient of various concentrations of curcumin.

Sample	Papp ($\times 10^{-6} \text{ cm/s}$)		Efflux ratio
	A \rightarrow B	B \rightarrow A	
5 $\mu\text{g}/\text{mL}$ curcumin	1.50 ± 0.43	1.17 ± 0.12	0.78
10 $\mu\text{g}/\text{mL}$ curcumin	1.23 ± 0.21	1.44 ± 0.09	1.17
20 $\mu\text{g}/\text{mL}$ curcumin	0.97 ± 0.23	1.56 ± 0.26	1.62
10 $\mu\text{g}/\text{mL}$ curcumin+500 μM verapamil	1.59 ± 0.17	-	-
20 $\mu\text{g}/\text{mL}$ curcumin+500 μM verapamil	1.59 ± 0.10	-	-

Table 2: Apparent permeability coefficient (P_{app}) and efflux ratios of curcumin transport across Caco-2 cell monolayers ($n=4$).

Time(min)	0	60	120	After 24 h of cultivation
A \rightarrow B				
5 $\mu\text{g}/\text{mL}$ curcumin	407 ± 56	460 ± 16	678 ± 33	848 ± 36
10 $\mu\text{g}/\text{mL}$ curcumin	435 ± 14	447 ± 6	665 ± 2	776 ± 183
20 $\mu\text{g}/\text{mL}$ curcumin	394 ± 11	450 ± 8	773 ± 83	544 ± 226
10 $\mu\text{g}/\text{mL}$ curcumin +500 μM verapamil	422 ± 9	434 ± 12	653 ± 26	728 ± 219
20 $\mu\text{g}/\text{mL}$ curcumin +500 μM verapamil	444 ± 16	454 ± 6	722 ± 27	980 ± 432
B \rightarrow A				
5 $\mu\text{g}/\text{mL}$ curcumin	437 ± 29	444 ± 6	655 ± 11	870 ± 212
10 $\mu\text{g}/\text{mL}$ curcumin	427 ± 21	437 ± 4	661 ± 14	804 ± 118
20 $\mu\text{g}/\text{mL}$ curcumin	407 ± 13	446 ± 14	755 ± 11	675 ± 18

Table 3: TEER of the Caco-2 cells monolayers in the Caco-2 transport study ($n=4$).

Discussion

Everted gut sacs are commonly used to study the intestinal absorption of drugs. The rate and amount of curcumin absorption can be studied by determining the composition of the intestinal absorption solution. Because of the lack of intestinal blood circulation and nerve control in this model, long-term *in vitro* intestinal mucosal functions can easily change, so an improved method was used in this experiment. The culture medium with HBSS was treated with 95% O₂/5% CO₂ so the exposed mucosal cells received an adequate supply of O₂ to ensure the continuing intestinal functions during the experiment. The everted sac model retains the characteristics of intact tissue and mucosa and can be used to simulate the absorption of drugs in different intestinal segments *in vitro*. Our study of the absorption mechanism in the ileum is perhaps the most helpful and persuasive portion of this experiment. On the basis of the experiment, we further investigated the mechanism of intestinal absorption of curcumin by using *in vivo* intestinal perfusion model. In situ intestinal perfusion is a technique to determine the absorption rate of a drug in situ in rats, which has recently become one of the main research models of drug absorption [6-8].

P-gp is a membrane transport protein that pumps the drug out of the cell, thereby reducing the bioavailability of drugs. P-gp is an ATP-dependent efflux pump that is highly expressed in multi-drug resistant tumor cells and the blood-brain barrier [9]. P-gp inhibitors can slow the drug efflux function of P-gp, thereby reversing drug resistance and increasing the amount of drug in the tissues. In order to improve the bioavailability of drugs, the combination of curcumin and P-gp inhibitors such as verapamil and cyclosporine can be considered. The key factor to ensure the effect of an oral medicine is that it must be absorbed in the digestive tract, and the intestinal transport protein P-gp plays a very important role in the transmembrane transport (including absorption and secretion) of lipid-soluble drugs. Therefore, it is necessary to strengthen the research on P-gp and its substrates [10].

The Caco-2 cell model has been widely used to study the absorption and transport mechanisms of drugs in the small intestine. Caco-2 cells were first isolated from human colon cancer cells in the 1970s, and the model was originally proposed by Markowska et al. [11]. Caco-2 cells possess the same cell polarity and compact single-cell layer as do human intestinal epithelial cells. The morphology and function of cells were similar to those of human intestinal epithelial cells. Because most of the absorption processes and mechanisms of action of Chinese herbal medicines are not yet elucidated, the Caco-2 cell model has wide applications in the study of the oral absorption of Chinese medicines [12-14].

The P_{app} value reflects the ability of drug molecules to penetrate the intestinal tract, and ER reflects the role of the efflux protein to expel drugs—the greater the ER, the more powerful the efflux effect. Based on our findings using a Caco-2 cell model to study curcumin, we determined that the transmembrane transport of curcumin can be enhanced by the presence of a P-gp inhibitor, and the Papp value was increased after the addition of verapamil. From the results, we see that in a monolayer of Caco-2 cells, the transport mode of curcumin is passive diffusion at concentrations <10 µg/mL and when the concentration is >10 µg/mL, active transport is involved. The transport mechanism for curcumin is a combination of active and passive processes. This is confirmed in the experiments by the addition of an inhibitor, and the value is close to P_{appA→B} of 5 µg/mL.

Curcumin is present in many foods and traditional Chinese medicines and is an important material basis for food nutrition and traditional Chinese medicine. In this paper, the experimental results reveal that, when the content of curcumin in food and medicinal materials is low, curcumin can be absorbed into the body at low dose levels by passive diffusion and is a substrate for the drug intestinal transporter P-gp. Careful use of other P-gp components can improve a drug's competitive transport into the gastrointestinal tract. The results of this study can also be used to explain some of the reasons for the poor performance of certain formulations of curcumin as a high-purity compound. Chinese foods and Chinese herbal medicines containing curcumin can still play a role in nutrition and health.

Conclusion

In summary, curcumin transport results from a combination of passive diffusion and active transport. Curcumin is a substrate of the intestinal transporter P-gp, and the mechanism by which intestinal absorption of curcumin takes place is related to the mechanism of intestinal P-gp transport.

Declaration of interest

The authors declare that they have no competing interests in this work.

Acknowledgments

This thesis is sponsored by Jiangsu Province and the Opening Foundation of Key Laboratory of acupuncture and moxibustion (No. KJA200802)

References

1. Jiang S, Han J, Li T, Xin Z, Ma Z, et al. (2017) Curcumin as a potential protective compound against cardiac diseases. *Pharmacol Res* 119: 373-383.
2. Rezaee R, Momtazi AA, Monemi A, Sahebkar A (2017) Curcumin: a potentially powerful tool to reverse cisplatin-induced toxicity. *Pharmacol Res* 117: 218-227.
3. Wang YJ, Pan MH, Cheng AL, Lin LI, Ho YS, et al. (1997) Stability of curcumin in buffer solutions and characterization of its degradation products. *J Pharm Biomed Anal* 15: 1867-1876.
4. Ravindranath V, Chandrasekhara N (1980) Absorption and tissue distribution of curcumin in rats. *Toxicology* 16: 259-265.
5. Shoba G, Joy D, Joseph T, Majeed M, Rajendran R, et al. (1998) Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers. *Planta Med* 64: 353-356.
6. Carmona A (1998) A simple *in vitro* perfusion system to measure intestinal nutrient uptake. *J Nutri Biochem* 9: 52-57.
7. Levet-Trafit B, Gruyer MS, Marjanovic M, Chou RC (1996) Estimation of oral drug absorption in man based on intestine permeability in rats. *Life Sci* 58: 359-363.
8. Abuhelwa AY, Williams DB, Upton RN, Foster DJ (2017) Food, Gastrointestinal pH, and Models of Oral Drug Absorption. *Eur J Pharm Biopharm* 112: 234-248.
9. Li M, de Graaf IA, van de Steeg E, de Jager MH, Groothuis GM (2017) The consequence of regional gradients of P-gp and CYP3A4 for drug-drug interactions by P-gp inhibitors and the P-gp/CYP3A4 interplay in the human intestine *ex vivo*. *Toxicol In Vitro* 40: 26-33.
10. Huo X, Liu Q, Wang C, Meng Q, Sun H, et al. (2013) Enhancement effect of P-gp inhibitors on the intestinal absorption and antiproliferative activity of bestatin. *Eur J Pharm Sci* 50: 420-428.

-
11. Markowska M, Oberle R, Juzwin S, Hsu CP, Gryszkiewicz M, et al. (2001) Optimizing Caco-2 cell monolayers to increase throughput in drug intestinal absorption analysis. J Pharmacol Toxicol 46: 51-55.
 12. Walgren RA, Karnaky KJ, Lindenmayer GE, Walle T (2000) Efflux of dietary flavonoid quercetin 4'- β -glucoside across human intestinal Caco-2 cell monolayers by apical multidrug resistance-associated protein-2. J Pharmacol Exp Ther 294: 830-836.
 13. Wang J, Ma W, Tu P (2015) The mechanism of self-assembled mixed micelles in improving curcumin oral absorption: *in vitro* and *in vivo*. Colloid Surface B 133: 108-119.
 14. Shen C, Chen R, Qian Z, Meng X, Hu T, et al. (2015) Intestinal absorption mechanisms of MTBH, a novel hesperetin derivative, in Caco-2 cells, and potential involvement of monocarboxylate transporter 1 and multidrug resistance protein 2. Eur J Pharm Sci 78: 214-224.