

Transgenesis through Blastomeres Transfection

Islam M. Saadeldin*

Department of Physiology, Faculty of Veterinary Medicine, Zagazig University, Zagazig, Korea

Clinical Image

Parthenogenetically activated porcine oocytes were subjected to pronase 0.3% (in PBS w/v) for 90 seconds to digest the zona pellucida. Zona-free blastomeres were transfected with piggybac transposon DNA using Fugene HD transfection reagent (Promega Corp., Madison, WI, USA).

In contrary to the conventional transgenesis techniques, such as somatic cell nuclear transfer, sperm-mediated transfection and pronuclear injection; this method paves the way to modify the embryo genome in a simple and efficient manner in addition to increase blastocyst total cell count caused by blastomeres aggregation.




Figure 1: Tnsfection of zona pellucida free blastomeres aggregates with piggyback transposon (pB-CA-cherry red). A. Day 2 control blastomeres with fugene HD transfection reagent and without the DNA. B. Day 2 transfected blastomeres with piggybac DNA, C. transgenic blastocyst resulted from aggregated blastomeres transfection showing cherry red fluorescence. A'-C': under visible light images. Scale bar=200 μ m.

*Corresponding author: Islam M. Saadeldin, Department of Physiology, Faculty of Veterinary Medicine, Zagazig University, 44519, Zagazig, Korea, Tel: +20-11000-24182; E-mail: islamms@snu.ac.kr

Received June 28, 2015; Accepted June 29, 2015; Published July 01, 2015

Citation: Saadeldin IM (2015) Transgenesis through Blastomeres Transfection Baculovirus Infection. Clon Transgen 4: i103. doi:10.4172/2168-9849.1000i103

Copyright: © 2015 Saadeldin IM. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.