

Zeolite-based Photocatalysts: A Promising Strategy for Efficient Photocatalysis

Yanan Guo, Baiyi Zu and Xincun Dou*

Key Laboratory of Functional Materials and Devices for Special Environments, Laboratory of Environmental Science and Technology, Xinjiang Technical Institute of Physics & Chemistry, Chinese Academy of Sciences, Urumqi, 830011, China

A photocatalyst is generally defined as a material that has the capability to induce chemical reactions of substances of interest adsorbed on the surface by light-generated active radicals [1]. The efficiency of the photocatalyst is a function of the balance between charge separation, ease of interfacial electron transfer and energy-wasting charge recombination [2]. With the inspiration of improving photocatalysis efficiency and selectivity, people are trying hard to develop new photocatalysts.

Zeolites are crystalline nanoporous inorganic materials with well-defined interconnected channels or cavities in the nanometre or subnanometre length scale, termed as micropores (0.5-2 nm) [3,4]. With the uniform pore size, polar environment, high surface area, the internal active sites and excellent adsorption capability [5], zeolites could enhance the efficiency and selectivity of photocatalysts either by photoactivating the zeolite framework or by encapsulating with semiconductor oxides.

Photoactivating the zeolite framework through the incorporation of heteroatoms (Ti and other transition metals) can make the structure a photocatalyst [6-9]. The most representative example of titanium containing zeolite is titanosilicalite TS-1, which has the MFI structure [10]. There is a characteristic absorption band at 225 nm, which originates from the incorporation of the tetrahedral Ti atoms and the electron transfer in ligand-to-metal (-O-Ti). The photocatalytic properties of TS-1 was further proved by the decomposition of 4-nitrophenol in aqueous solution when illuminated with a 500 W high pressure Hg lamp [11]. The tetrahedral Ti atoms tripodally connected to the TS-1 framework are able to form the titanohydroperoxy species when interacting with H₂O₂, photochemically generating the •OH radicals. It was found that Si/Ti ratios (from 3.3 to 26.3) in TS-1 can greatly affect the photocatalytic efficiency. The TS-1 with the highest Si/Ti ratio exhibited a remarkable enhancement of the photocatalytic activity in the presence of H₂O₂ [11]. TS-1 has also been used to degrade monoethanolamine in aqueous solution by forming ethanolamine-Ti complex [12]. It was found that the photocatalytic activity of TS-1 is comparable with TiO₂ per weight of photocatalyst and the intrinsic activity per Ti atom in TS-1 is higher than that in TiO₂.

Besides TS-1, Ti-Beta zeolite could be used for the selective oxidation of alkenes in the presence of organic hydroperoxides [13]. It was found that the hydrophilic or hydrophobic properties of the zeolite cavities could control the reactivity and selectivity in the photocatalytic reduction of CO₂ with H₂O to produce CH₄ and CH₃OH on these Ti-Beta zeolite catalysts [14]. This result opened a new way to improve the photocatalytic activity by modifying the surface properties of zeolite. Furthermore, a novel titanosilicate ETS-10 contains photoexcitable Ti-O-Ti 1-dimensional quantum wires and 3-dimensional 12-ring channel was reported recently [15]. It was found that this ETS-10 could act as a shape-selective photocatalyst for the degradation of a mixture of phenols of different sizes [16]. The compounds which are small enough to access the interior of the micropores become protected and are degraded more slowly, while the others that are too large to enter the pores are degraded preferentially.

It is relatively difficult to obtain zeolites with photocatalytically active framework, and the amount of these zeolites is pretty few. Another class of zeolite-based photocatalysts, which incorporates semiconductor oxides into the cavities either by ion exchange or by hydrothermal method, has been widely studied [17-20]. In this type of semiconductor @ zeolite photocatalysts, the unique characteristics of zeolite could be fully utilized, such as, the crystal structure determines the size of the encapsulated particles and the optical absorption bandgap, [21] the polar environment favors the photoinduced electron transfer and minimizes the electron-hole recombination [22]. The most representative semiconductor @ zeolite photocatalyst is TiO₂ @ Zeolite, which has been widely studied recently. It was found that TiO₂ @ Zeolite is efficient for the decomposition of NO, [23] for the efficient removal of NH₃ and H₂S from air [24] and for the reduction of CO₂ [25]. TiO₂ @ Zeolite is also efficient for selectivity photocatalyze CH₃OH with minor product concentration of CO and O₂ [26]. Furthermore, it was suggested that different semiconductor oxides located in different cages of zeolite may further synergistically enhance the photocatalytic performance of semiconductor @ zeolite photocatalyst [19]. For example, a maximum water splitting efficiency of (ZnS-CdS) @ Y photocatalyst was achieved at a Cd/Zn ratio of 0.25 and when CdS and ZnS clusters were located in α -cages and β -cages, respectively [27,28].

The photocatalytic efficiency of zeolites could be further enhanced by incorporating the photoactive zeolite framework and semiconductor photocatalysts together. For example, microporous titanosilicates (ETS-4, ETS-10) have been utilized to incorporate CdS, in which the Ti-O-Ti backbone acts as a nanowire capable of accepting electrons from the photoexcited CdS [29,30]. The encapsulation of CdS in ETS zeolite is effective for improving the activity as well as the stability of CdS.

In conclusion, by photoactivating the zeolite framework and by encapsulating with semiconductor oxides, the efficiency and selectivity of zeolite-based photocatalysts could be enhanced effectively. However, little is known about the photocatalysis mechanism on a molecular level, to which people should pay special attention in the future.

References

1. Mills A, Hunte SL (1997) An overview of semiconductor photocatalysis. *J Photochem Photobiol A-Chem* 108: 1-35.

*Corresponding author: Xincun Dou, Key Laboratory of Functional Materials and Devices for Special Environments, Laboratory of Environmental Science and Technology, Xinjiang Technical Institute of Physics & Chemistry, Chinese Academy of Sciences, Urumqi, 830011, China, Tel: +86-991-367-7875; Fax: +86-991-383-8957; E-mail: xcdou@ms.xjb.ac.cn

Received July 16, 2013; Accepted July 17, 2013; Published July 20, 2013

Citation: Guo Y, Zu B, Dou X (2013) Zeolite-based Photocatalysts: A Promising Strategy for Efficient Photocatalysis. *J Thermodyn Catal* 4: e120. doi: [10.4172/2157-7544.1000e120](https://doi.org/10.4172/2157-7544.1000e120)

Copyright: © 2013 Guo Y, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

2. Linsebigler AL, Lu G, Yates Jr JT (1995) Photocatalysis on TiO₂ surfaces: principles, mechanisms, and selected results. *Chemical reviews* 95: 735-758.
3. Breck DW (1973) *Zeolite Molecular Sieves: Structure, Chemistry, and Use*. New York, Wiley.
4. Xu R, Pang W, Yu J, Huo Q, Chen J (2007) *Chemistry of zeolites and related porous materials: synthesis and structure*. Singapore, Wiley.
5. Corma A, Garcia H (2004) Zeolite-based photocatalysts. *Chem Commun* 1443-1459.
6. Notari B (1991) Titanium silicalite: a new selective oxidation catalyst. *Studies in Surface Science and Catalysis* 67: 243-256.
7. Hashimoto S (2003) Zeolite photochemistry: impact of zeolites on photochemistry and feedback from photochemistry to zeolite science. *Journal of Photochemistry and Photobiology C: Photochemistry Reviews* 4: 19-49.
8. Krissanasaerane M, Wongkasemjit S, Cheetham AK, Eder D (2010) Complex carbon nanotube-inorganic hybrid materials as next-generation photocatalysts. *Chemical Physics Letters* 496: 133-138.
9. Zhang J, Zhao DS, Yang LY, Li YB (2010) Photocatalytic oxidation dibenzothiophene using TS-1. *Chemical Engineering Journal* 156: 528-531.
10. Clerici MG (1991) Oxidation of saturated hydrocarbons with hydrogen peroxide, catalysed by titanium silicalite. *Applied Catalysis* 68: 249-261.
11. Lee GD, Jung SK, Jeong YJ, Park JH, Lim KT, et al. (2003) Photocatalytic decomposition of 4-nitrophenol over titanium silicalite (TS-1) catalysts. *Applied Catalysis A: General* 239: 197-208.
12. Ban T, Kondoh S, Ohya Y, Takahashi Y (1999) Degradation reaction of monoethanolamine using TS-1 zeolite as a photocatalyst. *Physical Chemistry Chemical Physics* 1: 5745-5752.
13. Corma A, Cambor M, Esteve P, Martinez A, Perezpariente J (1994) Activity of Ti-Beta catalyst for the selective oxidation of alkenes and alkanes. *Journal of Catalysis* 145: 151-158.
14. Ikeue K, Yamashita H, Anpo M, Takewaki T (2001) Photocatalytic reduction of CO₂ with H₂O on Ti-β zeolite photocatalysts: effect of the hydrophobic and hydrophilic properties. *J Phys Chem* 105: 8350-8355.
15. Howe RF, Krisnandi YK (2001) Photoreactivity of ETS-10. *Chem commun* 1588-1589.
16. Calza P, Pazé C, Pelizzetti E, Zecchina A (2001) Shape-selective photocatalytic transformation of phenols in an aqueous medium. *Chemcommun* 20: 2130-2131.
17. Liu ZF, Liu ZC, Wang Y, Li YB, Qu L, et al. (2012) Photocatalysis of TiO₂ nanoparticles supported on natural zeolite. *Materials Technology: Advanced Performance Materials* 27: 267-271.
18. Huang M, Xu C, Wu Z, Huang Y, Lin J, et al. (2008) Photocatalytic discolorization of methyl orange solution by Pt modified TiO₂ loaded on natural zeolite. *Dyes and Pigments* 77: 327-334.
19. White JC, Dutta PK (2011) Assembly of nanoparticles in zeolite Y for the photocatalytic generation of hydrogen from water. *The Journal of Physical Chemistry C* 115: 2938-2947.
20. Liu XS, Lu KK, Thomas JK (1993) Preparation, characterization and photoreactivity of titanium (IV) oxide encapsulated in zeolites. *J Chem Soc Faraday Trans* 89: 1861-1865.
21. Cosa G, Galletero MS, Fernández L, Márquez F, García H, et al. (2002) Tuning the photocatalytic activity of titanium dioxide by encapsulation inside zeolites exemplified by the cases of thianthrene photooxygenation and horseradish peroxidase photodeactivation. *New J Chem* 26: 1448-1455.
22. Bossmann SH, Turro C, Schnabel C, Pkhrel MR, Payawan LM, et al. (2001) Ru(bpy)₃²⁺/TiO₂-codoped zeolites: synthesis, characterization, and the role of TiO₂ in electron transfer photocatalysis. *J Phys Chem B* 105: 5374-5382.
23. Yamashita H, Ichihashi Y, Anpo M, Hashimoto M, Louis C, et al. (1996) Photocatalytic decomposition of NO at 275 K on titanium oxides included within Y-zeolite cavities: The structure and role of the active sites. *J Phys Chem* 100: 16041-16044.
24. Rasmussen SB, Portela R, Suárez S, Coronado JM, Rojas-Cervantes ML, et al. (2010) Hybrid TiO₂-SiMgOX Composite for Combined Chemisorption and Photocatalytic Elimination of Gaseous H₂S. *Ind Eng Chem Res* 49: 6685-6690.
25. Wang C, Thompson RL, Baltrus J, Matranga C (2009) Visible Light Photoreduction of CO₂ Using CdSe/Pt/TiO₂ Heterostructured Catalysts. *J Phys Chem Lett* 1: 48-53.
26. Ikeue K, Yamashita H, Anpo M (1999) Photocatalytic reduction of CO₂ with H₂O on titanium oxides prepared within the FSM-16 mesoporous zeolite. *Chemistry Letters* 28: 1135-1136.
27. Suárez S (2013) *Design of Advanced Photocatalytic Materials for Energy and Environmental Applications*. Green Energy and Technology Springer, 171-194.
28. Tel'biz G, Shwets A, Gunko V, Stoch J, Tamulajtis G, et al. (1994) Preparation and Properties Quantized Semiconductor Particles in Zeolites. *Studies in Surface Science and Catalysis* 84: 1099-1106.
29. Guan G, Kida T, Kusakabe K, Kimura K, Fang X, et al. (2004) Photocatalytic H₂ evolution under visible light irradiation on CdS/ETS-4 composite. *Chemical Physics Letters* 385: 319-322.
30. Guan G, Kida T, Kusakabe K, Kimura K, Abe E, et al. (2005) Photocatalytic activity of CdS nanoparticles incorporated in titanium silicate molecular sieves of ETS-4 and ETS-10. *Applied Catalysis A: General* 295: 71-78.