

Thermodynamic Mechanism of Protein Bindings

Iizuka Matsui*

Department of Diabetes and Endocrinology, University of Gifu, Gifu, Japan

DESCRIPTION

Protein binding is a fundamental concept in biochemistry and pharmacology that plays a pivotal role in various biological processes and has significant implications for drug development and the understanding of disease mechanisms. Protein binding, also known as protein-ligand binding or molecular binding, refers to the specific interaction between a protein molecule and another molecule, which could be a ligand, substrate, ion, or another protein [1]. This interaction occurs through non-covalent forces, primarily hydrogen bonds, electrostatic attractions, van der Waals forces, and hydrophobic interactions. The resulting complex, known as a protein-ligand complex, plays a pivotal role in many biological processes. Proteins, the workhorses of biology, have diverse structures and functions. They include enzymes, receptors, transporters, antibodies, and structural proteins, among others. Protein binding can modulate the function of these proteins, leading to various cellular responses. Understanding protein-ligand interactions at a molecular level is crucial for unravelling the mechanisms behind biological processes and for designing effective therapeutic interventions [2,3]. Protein binding can be categorized into several types based on the nature of the interacting molecules and their functional roles. Enzymes are proteins that catalyze biochemical reactions. Enzyme-substrate binding is a specific interaction where the enzyme binds to its substrate at the enzyme's active site. This binding is essential for catalysis and the conversion of substrates into products [4]. Receptors are proteins that recognize and bind specific ligands, such as hormones, neurotransmitters, or drugs. The binding of a ligand to a receptor triggers a cellular response, initiating signal transduction pathways or altering gene expression. Proteins can also bind to other proteins, forming complexes with distinct functions [5,6]. These interactions underlie processes such as protein signaling, cell adhesion, and regulation of gene expression. Examples include the binding of transcription factors to Deoxyribonucleic Acid (DNA) or the interaction between components of a signaling cascade. Transport proteins, like hemoglobin, bind to specific molecules for transport within the body [7]. Hemoglobin, for instance, binds to oxygen in the lungs and releases it in oxygen-deprived tissues. The protein binding plays a pivotal role in

several key biological processes and has profound implications for various scientific and medical fields [8]. Understanding protein-ligand interactions is crucial in drug discovery and development. Drugs often function by binding to specific proteins to modulate their activity [9,10]. The binding affinity and specificity of drugs to their target proteins determine their therapeutic efficacy. Protein binding influences the distribution and elimination of drugs in the body. When drugs bind tightly to plasma proteins, they may have a longer duration of action but may also be less available for their intended pharmacological effect. Protein-protein interactions are fundamental to molecular biology. They regulate processes such as gene expression, Deoxyribonucleic Acid (DNA) replication, and signal transduction [11]. Understanding these interactions helps elucidate cellular mechanisms. Dysregulation of protein binding can lead to various diseases. For example, mutations in the binding sites of ion channels can result in channelopathies, and aberrant receptor-ligand interactions can contribute to cancer or autoimmune disorders. Protein binding studies are essential for elucidating the three-dimensional structures of protein-ligand complexes [12]. Techniques like X-ray crystallography and Nuclear Magnetic Resonance (NMR) spectroscopy provide insights into the atomic-level details of these interactions. Computational methods, such as molecular docking and molecular dynamics simulations, predict protein-ligand interactions by modelling their atomic interactions and dynamics. The development of drug resistance, where proteins mutate to reduce drug binding, is a significant concern. Combating drug resistance requires innovative approaches, including the development of next-generation therapeutics [13]. Protein binding is a multifaceted concept that underlies essential biological processes and has far-reaching implications for fields such as drug discovery, molecular biology, and disease research. Understanding the diverse types of protein binding, its significance, and the techniques [14].

REFERENCES

1. Naba S, Numoto N, Ogawa S, Morii H, Ikura T, Abe R, et al. Crystal structures and thermodynamic analysis reveal distinct mechanisms of CD28 phosphopeptide binding to the Src homology 2 (SH2) domains of three adaptor proteins. *J Biol Chem.* 2017;292(3): 1052-60.

Correspondence to: Iizuka Matsui, Department of Diabetes and Endocrinology, University of Gifu, Gifu, Japan, E-mail: matusi999@gmail.jp

Received: 03-May-2025, Manuscript No. JTC-23-26246; **Editor assigned:** 05-May-2025, Pre QC No. JTC-23-26246(PQ); **Reviewed:** 19-May-2025, QC No. JTC-23-26246; **Revised:** 26-May-2025, Manuscript No JTC-23-26246 (R)**Published:** 02-Jun-2025, DOI: 10.32548/2157-7544.25.16.442

Citation: Matsui L (2025) Thermodynamic Mechanism of Protein Bindings. *J Thermodyn.* 16:442.

Copyright: © 2025 Matsui L. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

2. Ziegler A, Seelig J. Interaction of the protein transduction domain of HIV-1 TAT with heparan sulfate: binding mechanism and thermodynamic parameters. *Biophys J.* 2004;86(1):254-63.
3. Milenković D, Dimić D, Avdović E, Simijonović D, Vojinović R, Marković Z et al. A thermodynamic and kinetic HO radical scavenging study and protein binding of baicalein. *The J Phys Chem B.* 2023 Jun:107110.
4. Manley LJ, Lin MM. Kinetic and thermodynamic allostery in the Ras protein family. *bioRxiv.* 2022:2022-11.
5. Racz CP, Racz LZ, Floare CG, Tomoaia G, Horovitz O, Riga S, et al. Curcumin and whey protein concentrate binding: Thermodynamic and structural approach. *Food Hydrocoll.* 2023;139:108547.
6. Patni D, Jha SK. Thermodynamic modulation of folding and aggregation energy landscape by DNA binding of functional domains of TDP-43. *Biochim Biophys Acta Proteins Proteom.* 2023;1871(4):140916.
7. Paul BK. Thermodynamic insight into the interaction mechanism of an ES IPT drug with a serum transport protein: Slower solvent-relaxation dynamics explored through wavelength-sensitive fluorescence behavior. *Chem Phys Impact.* 2023;6:100225.
8. Saikia S, Das PK. Monolayer protein adsorption on cationic gold nanoparticles: A thermodynamic study. *Chemical Physics Letters.* 2023;825:140595.
9. Shpilt Z, Tshuva EY. Binding of the anticancer Ti (IV) complex phenolTi to serum proteins: Thermodynamic and kinetic aspects. *J Inorg Biochem.* 2022;232:111817.
10. Meng Y, Wei Z, Xue C. Deciphering the interaction mechanism and binding mode between chickpea protein isolate and flavonoids based on experimental studies and molecular simulation. *Food Chem.* 2023;429:136848.
11. Han C, Zheng Y, Zhou C, Liang Y, Huang S, Sun Y, et al. Pressurization-induced molten globule-like protein as a regulatory target for off-flavor controlling: Thermodynamic evidence and molecular modeling. *Biochemistry.* 2023:110022.
12. Zhao L, Ouyang D, Cheng X, Zhou X, Lin L, Wang J, et al. Multi-frequency ultrasound-assisted cellulase extraction of protein from mulberry leaf: Kinetic, thermodynamic, and structural properties. *Ultrason Sonochem.* 2023:106554.
13. Xu L, Wan Y, Liu X, Qin Z, Zhao Y, Fu X, et al. Insights on the binding mechanism between specified aldehydes and flaxseed protein using multispectral image and molecular docking. *Food Chem.* 2023;422:136256.
14. Xie G, Timasheff SN. The thermodynamic mechanism of protein stabilization by trehalose. *Biophys Chem.* 1997;64(1-3):25-43.