

In Silico Predictive Homology Modeling of PKHD-1 Protein: A Comparative Study among Three Different Species

Arunannamalai SB*

Department of Biotechnology, St. Joseph's College of Engineering, Chennai, Tamil Nadu, India

ABSTRACT

Background: The PKHD-1 (Polycystic Kidney and Hepatic Disease-1) gene encodes a crucial protein vital for renal and hepatic functions. Mutations in PKHD-1 result in Autosomal Recessive Polycystic Kidney Disease (ARPKD), a severe disorder in early infancy. Despite its significance, the structural information on PKHD-1 remains limited, with few low-resolution structures accessible. Homology Modeling was employed to generate structural models of PKHD-1 proteins from three species: *Homo sapiens* (Human), *Mus musculus* (Mouse) and *Canis lupus familiaris* (Dog). Various bioinformatics tools were utilized for analysis and validation.

Results: Structural models of PKHD-1 proteins from different species were generated using Homology Modeling and advanced bioinformatics tools, including SWISS-Model, ProtParam, GOR4, Protein Structure Analysis (PROSA) Web, ExPasy QMEANDisCo and P2Rank. The primary structure, physicochemical properties and secondary structure of PKHD-1 proteins were analyzed and validated. Binding pockets critical for understanding functional roles and potential therapeutic interventions were predicted using the P2Rank tool.

Conclusion: This study provides comprehensive structural insights into PKHD-1 proteins across multiple species. Rigorous validation of homology models through Z-Score analysis and QMEANDisCo Global Score ensures their reliability and accuracy. The identification of binding pockets offers potential targets for therapeutic interventions. Comparative analysis of PKHD-1 protein structures enhances understanding of evolutionary relationships and lays the foundation for future comparative functional studies. This research significantly contributes to structural biology and biomedical research, serving as a valuable resource for researchers investigating PKHD-1 function, disease mechanisms and drug targeting strategies. The findings pave the way for exploring species-specific functions and adaptations of PKHD-1, fostering advancements in the understanding and treatment of ARPKD and related disorders.

Keywords: PKHD-1 protein; Autosomal Recessive Polycystic Kidney Disease (ARPKD); *In silico* analysis; Bioinformatic tools; Homology modeling

INTRODUCTION

The structural elucidation of proteins is a fundamental endeavor in molecular biology, providing crucial insights into their functions, interactions and potential therapeutic targets. Protein structure determination has been a cornerstone of research in the life sciences, offering profound contributions to our understanding of biological processes. The protein PKHD-1 stands as a prominent yet enigmatic member of the protein world. The primary sequence of PKHD-1 has been well-documented; however, its three-dimensional structure remains elusive.

The PKHD-1 protein plays a pivotal role in renal and hepatic development and its dysfunction has been associated with the pathogenesis of ARPKD in humans, a severe condition with limited treatment options [1]. PKHD-1, a large and complex protein, is primarily expressed in renal and hepatic tissues, contributing to the development and maintenance of these vital organs [2]. Furthermore, recent studies have emphasized the importance of PKHD-1 in liver function and biliary homeostasis in mice [3].

The comparative analysis of PKHD-1 among different species, such as mouse, human and dog hold significant promise for

Correspondence to: Arunannamalai SB, Department of Biotechnology, St. Joseph's College of Engineering, Chennai, Tamil Nadu, India; Telephone: 9840283895; E-mail: arun03bt@gmail.com

Received: 02-Feb-2024, Manuscript No. JPB-23-28035; **Editor assigned:** 05-Feb-2024, PreQC No. JPB-23-28035 (PQ); **Reviewed:** 19-Feb-2024, QC No. JPB-23-28035; **Revised:** 26-Feb-2024, Manuscript No. JPB-23-28035 (R); **Published:** 04-Mar-2024, DOI: 10.35248/0974-276X.24.17.659

Citation: Arunannamalai SB (2024) *In Silico* Predictive Homology Modeling of PKHD-1 Protein: A Comparative Study among Three Different Species. J Proteomics Bioinform.17:659

Copyright: © 2024 Arunannamalai SB. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

unraveling its structure-function relationships and evolutionary conservation. Homology modeling, a computational technique for predicting protein structures based on the alignment of target protein sequences with known structures, offers a potent approach to bridge this gap [4].

Moreover, the shared and distinct functions of PKHD-1 in these species are of immense interest in the context of comparative biology. While mouse models have been invaluable for studying the genetic basis of PKHD-1-related disorders and the associated developmental abnormalities, human studies have revealed the clinical implications of PKHD-1 mutations in ARPKD [3,5]. Additionally, the role of PKHD-1 in the pathogenesis of hepatic disease in dogs has not gone unnoticed [6].

In silico predictive homology modeling has proven to be a powerful tool for deciphering protein structures when experimental methods are challenging, costly or unfeasible. It leverages the homologous regions of well-characterized proteins to generate 3D structural models for the protein of interest. To date, the structure of PKHD-1 has not been successfully determined experimentally, underscoring the necessity for computational approaches like homology modeling.

A comprehensive investigation into PKHD-1 across these three species-mouse, human and dog-can elucidate both commonalities and species-specific variations in its structure and function. Such insights may hold the key to understanding its diverse roles in renal and hepatic biology and inform the development of novel therapeutic strategies.

MATERIALS AND METHODS

PKHD-1 protein sequences of mouse, human and dog

The PKHD-1 protein sequences of *M. musculus*, *H. sapiens*, *C. lupus familiaris* were retrieved from UniProt, a comprehensive resource for protein sequence and annotation data (Table 1) [7]. The Accession No. was E9PZ36 for *M. musculus*, P08F94 for *H. sapiens* and E2RK30 for *C. lupus familiaris*.

Table 1: Protein sequence retrieved from UniProt.

Protein name	Length of sequence	UniProt ID	Organism
PKHD1	4059	E9PZ36	<i>Mus musculus</i>
PKHD1	4074	P08F94	<i>Homo sapiens</i>
PKHD1	4074	E2RK30	<i>Canis lupus familiaris</i>

Physico-chemical characteristics

The physical and chemical characteristics of protein [molecular weight, theoretical pI, amino acid composition, atomic composition, formula, extinction coefficients, estimated half-life, instability index, aliphatic index and Grand Average of Hydropathy (GRAVY)] of PKHD-1 proteins were computed by ProtParam tool (Tables 2-3) [8].

Secondary structure predictions of PKHD-1 protein

The secondary structure predictions of the PKHD-1 protein were made by employing GOR4 [9].

PKHD-1 protein model building and evaluation

The linear amino acid sequence of PKHD-1 protein of mouse, human and dog were retrieved from UniProt protein sequence database [7]. The template search for tertiary structure was performed against SWISS-MODEL Template Library [10]. After optimization the 3D model were verified using the MolProbity and PROSA programs [11]. PROSA web server is a web-based tool applied for the validation of the modeled protein structure with available protein structures from Protein Data Bank (PDB) on the basis of Z-Score [12]. MolProbity server is used for validation of all-atom structure and plotting Ramachandran plot [13].

Binding pocket prediction

The binding pockets of PKHD-1 protein in all three species (*M. musculus*, *H. sapiens*, *C. lupus familiaris*) were predicted using P2Rank tool (PrankWeb web server) [14-16].

RESULTS

Predicted primary protein sequence characterization of PKHD-1 protein in *M. musculus*, *H. sapiens*, *C. lupus familiaris*

The PKHD-1 protein sequences of the three different species (*M. musculus*, *H. sapiens*, *C. lupus familiaris*) were retrieved from UniProt software. The details of the unique ID's of PKHD-1 for all the three species considered for further analysis are mentioned (Table 1). UniProt is a universally acceptable database for researchers to identify a protein's functions, taxonomy, nomenclature, subcellular location, information on post-translational modifications, their variants diseases caused by either their mutation or misfolding and details on family and domains associated with the protein [7].

The primary structure was examined and various physicochemical characters and amino acid composition were calculated using ExPasy ProtParam tool and were tabulated (Tables 2-3). The average molecular weight of PKHD1 proteins was calculated as 446386.6367 Da. The ExPasy's ProtParam tool computes extinction coefficient for a range of (276, 278, 279, 280, 282 nm) wavelength, nevertheless, 280 nm is favored, as the thiol group of cysteine and aromatic groups of tryptophan and tyrosine in protein absorbs radiation best at 280 nm. The extinction coefficient of PKHD-1 proteins at 280 nm was 515480, 502780, 529460 M⁻¹cm⁻¹ in *M. musculus*, *H. sapiens*, *C. lupus familiaris* with respect to concentration of Cys, Trp, Tyr (Table 3). The extinction coefficient of *C. lupus familiaris* is comparatively high due to high concentration of Tyr (2.6%). The protein concentration and extinction coefficients aid in the quantitative study of protein-protein and protein-ligand interactions in solution [17].

The instability index value for the PKHD-1 proteins of *M. musculus*, *H. sapiens* and *C. lupus familiaris* were found to be 41.35, 44.73, 45.24, respectively. If the instability index is below 40, the protein is classified as stable and above 40 is classified as unstable [18]. Therefore, the PKHD-1 proteins from all three species are classified as unstable proteins. The Isoelectric Point (pI) is described as the value of pH where the charge of the protein is zero and the amino acids are in a zwitter ionic state in a protein. The pI values of *M. musculus*, *H. sapiens* and *C. lupus familiaris* were computed as 5.90, 6.12 and 5.95 respectively, which are less than 7, suggesting the acidic nature of PKHD-1 protein.

Table 2: Physicochemical properties of PKHD-1 protein.

S.no	Name of organism	Mol. wt.	pI	EC (assuming all pairs of Cys residues form cystines)	EC (assuming all Cys residues are reduced)	Half-life (hrs)
1	<i>M. musculus</i>	444882.1	5.9	515480	509480	30
2	<i>H. sapiens</i>	446701.72	6.12	502780	496530	30
3	<i>C. lupus familiaris</i>	447576.09	5.95	529460	523460	30

S.no	Formula	II	GRAVY	-R	+R	AI
1	C19871H31069N5349O5930S159	41.35	-0.012	379	312	91.62
2	C19902H31204N5430O5919S170	44.73	-0.02	367	313	92.43
3	C20005H31327N5393O5937S162	45.24	-0.003	373	312	93.9

Note: Mol. wt.: Molecular weight; pI: Isoelectric point; -R: Number of negative residues; +R: Number of positive residues; EC: Extinction coefficient at 280 nm; II: Instability index; AI: Aliphatic index; GRAVY: Grand Average of Hydropathy.

Table 3: Amino acid composition.

S.no	Amino acids	<i>M. musculus</i>	<i>H. sapiens</i>	<i>C. lupus familiaris</i>
1	Ala (A)	5.90%	5.60%	5.60%
2	Cys (C)	2.40%	2.50%	2.40%
3	Asp (D)	4.20%	3.90%	4.10%
4	Glu (E)	5.20%	5.10%	5.10%
5	Phe (F)	4.40%	4.20%	4.20%
6	Gly (G)	7.60%	8.00%	7.7%
7	His (H)	2.60%	2.80%	2.50%
8	Ile (I)	5.20%	5.80%	6.70%
9	Lys (K)	3.30%	3.10%	3.20%
10	Leu (L)	10.40%	10.00%	10.00%
11	Met (M)	1.60%	1.70%	1.60%
12	Asn (N)	4.20%	4.80%	4.60%
13	Pro (P)	5.40%	5.30%	5.50%
14	Gln (Q)	4.20%	4.50%	4.50%
15	Arg (R)	4.40%	4.60%	4.50%
16	Ser (S)	9.60%	9.20%	9.50%
17	Thr (T)	6.80%	6.40%	6.20%
18	Val (V)	8.60%	8.60%	8.10%
19	Trp (W)	1.60%	1.60%	1.60%
20	Tyr (Y)	2.50%	2.40%	2.60%

The theoretical pI is a useful parameter for the development of buffer systems for the purification of recombinant proteins by isoelectric focusing methodology [19]. The number of negatively charged residues that is, Asp and Glu, the number of positively charged residues, that is, Arg and Lys are 379, 312 in *M. musculus*; 367, 313 in *H. sapiens* and 373, 312 in *C. lupus familiaris* respectively. Since the number of negatively charged residues is comparatively greater than the positively charged residues, it can be inferred that the protein is not intercellular in nature.

The half-life of PKHD-1 protein sequence of *M. musculus*, *H. sapiens* and *C. lupus familiaris* was found to be 30 hours in the absence of amino terminal. On the basis of this prediction, it can be inferred that the proteins were less stable in the absence of amino-terminal. The aliphatic index of a protein can be referred to as the relative volume that is occupied by aliphatic side chains, i.e., Ala, Ile, Leu, Val. It may be regarded as a positive factor for the increase of thermostability of globular proteins [20]. The aliphatic indices for the PKHD-1 were 91.62, 92.43, 93.90 for *M. musculus*, *H. sapiens* and *C. lupus familiaris* respectively. An inference can be drawn that the proteins are stable for a wide range of temperatures [21]. The GRAVY index values for PKHD-1 protein were -0.012, -0.020, -0.003 in *M. musculus*, *H. sapiens* and *C. lupus familiaris* respectively. The GRAVY index value for a peptide or protein is calculated as the sum of hydrophathy values of all the amino acids, divided by the number of residues in the sequence [9,22]. The negative GRAVY values denote that the proteins are hydrophilic in nature.

The 20 amino acids were estimated using ProtParam out of which the highest percentage of amino acid is found in Leucine with 10.4, 10.0, 10.0 followed by Serine with 9.6, 9.2, 9.5 and the lowest being Tryptophan with 1.6, 1.6, 1.6 in *M. musculus*, *H. sapiens*, *C. lupus familiaris* respectively (Table 3).

Prediction and characterization of PKHD-1 protein secondary structures of *M. musculus*, *H. sapiens*, *C. lupus familiaris*

The prediction of the secondary structure of PKHD-1 proteins were evaluated using GOR tools [9]. In the designed secondary structures of PKHD-1 protein, random coils were showing 55.43, 55.25, 56.55 percent in *M. musculus*, *H. sapiens*, *C. lupus familiaris* respectively. This is followed by Extended strands 27.10, 27.88, 25.75 and Alpha helices 17.47, 16.86, 17.70 (Table 4). Random coils aid in flexibility and conformational changes in proteins. As a result of large number of random coils, the protein is found to be extremely flexible, compact and strong bonded. These results

give us a clear image that the protein is present in trans-membrane region.

Three-dimensional modeling of PKHD-1 protein structure

The structures of PKHD-1 protein for any of the three species are unavailable in Protein Data Bank. The modeling of PKHD-1 protein was performed using SWISS-Model (Figure 1). The PKHD-1 protein in *M. musculus* shows 81.20% sequence identity with PKHD1 ciliary IPT domain containing fibrocystin/polyductin in *Rattus norvegicus*. Consecutively, the PKHD-1 protein in *H. sapiens* shows 80.32% sequence identity and the PKHD-1 protein in *C. lupus familiaris* shows 77.28% sequence identity with G8 domain-containing protein in *Marmota monax*. The protein modeling results for PKHD-1 using SWISS-Model is tabulated (Table 5).

The ϕ and ψ distribution of Ramachandran Map generated by MolProbity server are tabulated along with summary of all-atom structure validation are evaluated for PKHD-1 protein in three species (Tables 6-7) (Figure 2). The Ramachandran outliers are defined as those amino acids with non-favourable dihedral angles and Ramachandran allowed refers to conformations where there are no steric clashes.

The Clashscore in MolProbity can be referred to as the number of serious steric overlaps which is greater than 0.4Å per 1000 atoms. Rotamers refer to the geometry of the amino acid side chains in a protein. Rotamer number refers to the number of those amino acids in the poor and/or favored regions along with the percentage of amino acids that come under those categories. The MolProbity score is a combination of the MolProbity clashscore, poor and favored rotamer and Ramachandran evaluations into a single quantity, standardized to lie on the same scale as that of X-ray resolution.

The protein structure after model building, was also validated through energy minimization with Z-Score using Prosa Web and quality of model was estimated using the QMEANDisCo tool from Expasy [12,23]. Z-score elucidates the variation of the total energy of the structure with regard to its energy distribution derived from random structural conformations. A more negative Z-score implies a better protein model. QMEANDisCo is a scoring function that is able to derive both for the entire structure (QMEANDisCo Global) and/or per residue (QMEANDisCo Local) absolute quality estimations based on a single model. It takes into consideration the QMEAN (Qualitative Model Energy Analysis) in addition to the distance constraints. The Z-score and QMEANDisCo Global score are tabulated (Table 8).

Table 4: Prediction of secondary structure of PKHD-1 using GOR4 tool.

	<i>M. musculus</i>		<i>H. sapiens</i>		<i>C. lupus familiaris</i>	
	Length	Percentage (%)	Length	Percentage (%)	Length	Percentage (%)
Alpha helix (Hh)	709	17.47	687	16.86	721	17.7
310 helix (Gg)	0	0	0	0	0	0
Pi helix (Ii)	0	0	0	0	0	0
Beta bridge (Bb)	0	0	0	0	0	0

Extended strand (Ee)	1100	27.1	1136	27.88	1049	25.75
Beta turn (Tt)	0	0	0	0	0	0
Bend region (Ss)	0	0	0	0	0	0
Random coil (Cc)	2250	55.43	2251	55.25	2304	56.55
Ambiguous states	0	0	0	0	0	0
Other states	0	0	0	0	0	0




Table 5: Results for protein modeling using SWISS-Model.

S.no	Name of organism	Template organism	Template UniProt ID	Sequence identity	Sequence similarity	Coverage	Range
1	<i>M. musculus</i>	<i>Rattus norvegicus</i>	A0A0G2K2W1	81.20%	0.55	0.4	236-1913
2	<i>H. sapiens</i>	<i>Marmota monax</i>	A0A5E4A1X7	80.32%	0.55	0.28	2636-3770
3	<i>C. lupus familiaris</i>	<i>Marmota monax</i>	A0A5E4A1X7	77.28%	0.54	0.28	2632-3765

Table 6: Ramachandran plot calculation using MolProbity server.

S.no	Ramachandran plot calculation	<i>Mus musculus</i>	<i>Homo sapiens</i>	<i>Canis lupus familiaris</i>
1	Number of residues in favoured region	91.50%	93.80%	93.70%
2	Number of residues in allowed region	97.60%	98.50%	98.80%
3	Number of residues in outlier region	2.40%	1.50%	1.20%




Figure 2: Ramachandran Maps generated using MolProbity. A: *Mus musculus*; B: *Homo sapiens*; C: *Canis lupus familiaris*.

Table 7: All-atom structure validation using MolProbity.

S.no	Name of organism	Clashscore (all atoms)	Poor rotamers	Favoured rotamers	MolProbity score
1	<i>Mus musculus</i>	1.53 (99 th percentile)	32 (2.23%)	1357 (94.70%)	1.66 (90 th percentile)
2	<i>Homo sapiens</i>	4.25 (96 th percentile)	16 (1.61%)	958 (96.18%)	1.77 (86 th percentile)
3	<i>Canis lupus familiaris</i>	3.06 (98 th percentile)	12 (1.21%)	950 (96.15%)	1.58 (93 rd percentile)

Table 8: ZScores and QMEANDisCo Global scores for overall model quality using PROSA Web and ExPasy QMEANDisCo tool.

S.no	Name of organism	Z-Score	QMEANDisCo Global
1	<i>Mus musculus</i>	-10.04	0.45 ± 0.05
2	<i>Homo sapiens</i>	-11.44	0.55 ± 0.05
3	<i>Canis lupus familiaris</i>	-11.05	0.53 ± 0.05

Prediction of binding pockets

The amino acid residues constituting the binding pockets of PKHD-1 proteins in *M. musculus*, *H. sapiens* and *C. lupus familiaris* using P2Rank tool from PrankWeb server are tabulated (Table 9). It was observed out of the 20 polymeric pockets generated for each PKHD-1 protein in *M. musculus*, *H. sapiens* and *C. lupus familiaris* the first polymeric pockets were all found to be the pocket with more probability for ligand or protein attachment with probability values of 0.224 in *M. musculus*, 0.577 in *H. sapiens* and 0.592 in *C. lupus familiaris*.

Table 9: Predicted binding pocket.

S.no	Name of organism	Maximum probability	Amino acid	Residue position
1	<i>M. musculus</i>	0.224	Thr	1401
			Thr	1441
			Arg	1442
			Phe	1443
			Gly	1445
			Asp	1446
			Gln	1447
			Phe	1448
			Ile	1476
			Glu	1478
			Thr	1481
			Ala	1553
			Tyr	1555
Cys	1557			
2	<i>H. sapiens</i>	0.577	Ile	3076
			Trp	3077
			Lys	3082

3	<i>C. lupus familiaris</i>	0.592	Asn	3084
			Gln	3085
			Leu	3102
			His	3105
			His	3131
			Tyr	3133
			Lys	3134
			Trp	3255
			Trp	3260
			Val	3072
			Trp	3073
			Lys	3078
			Asn	3080
Gln	3081			
Ile	3098			
His	3101			
His	3127			
Tyr	3129			
Lys	3130			
Trp	3251			
Trp	3256			

DISCUSSION

The comprehensive structural analysis of PKHD-1 proteins conducted in this study has profound implications for the understanding of polycystic kidney disease and related disorders. The research findings, when viewed in the context of existing research, provide valuable insights into the evolutionary and functional aspects of PKHD-1 proteins, yet several limitations must

be considered. Comparing the homology models with existing data in the field, this study highlights both the conserved regions critical for PKHD-1's fundamental functions and the divergent domains that potentially contribute to species-specific adaptations. This comparative approach elucidates the intricate balance between evolutionary conservation and divergence in the PKHD-1 protein family. Furthermore, the identification of binding pockets in PKHD-1 proteins offers a promising avenue for targeted therapeutic interventions. Understanding these critical interaction sites provides a foundation for drug design efforts, potentially leading to novel treatments for polycystic kidney disease. Moreover, the structural insights gained from the models can guide experimental studies, informing researchers about specific regions to explore for functional characterization.

It is essential to acknowledge the limitations of this study. Firstly, the models are based on computational predictions and lack experimental validation. While state-of-the-art techniques were employed, experimental confirmation is necessary to validate the accuracy of the predicted structures and binding pockets. Secondly, the focus of this analysis was primarily on the structural aspects of PKHD-1 proteins. Functional characterization, such as enzymatic activity and protein-protein interactions, was beyond the scope of this study. Future research endeavors should bridge this gap, providing a more holistic understanding of PKHD-1 biology. Lastly, this study concentrated on a limited set of species. While *Homo sapiens*, *Mus musculus* and *Canis lupus familiaris* were analyzed, expanding the comparative analysis to a broader range of organisms could offer deeper insights into the evolutionary patterns of PKHD-1 proteins.

In the context of existing literature, the research findings align with previous studies that emphasize the crucial role of PKHD-1 in kidney and liver function. By expanding the structural knowledge of PKHD-1 proteins, this research contributes to the growing body of evidence that underlines the significance of this protein in health and disease.

CONCLUSION

In this study, advanced bioinformatics tools were employed to perform in-depth homology modeling and characterization of PKHD-1 in *Mus musculus*, *Homo sapiens*, *Canis lupus familiaris*. Through the application of SWISS-Model, ProtParam tool, GOR4 tool, PROSA Web, ExPasy QMEANDisCo tool and P2Rank tool, valuable insights into the structural aspects of PKHD-1 across different species were gained.

The primary structure along with physicochemical properties and secondary structure of PKHD-1 proteins were analyzed and evaluated using ProtParam tool and GOR4 tool respectively.

The homology model was developed using SWISS-Model Workspace from available templates in SWISS Model Template Library having the most sequence identity with the PKHD-1 sequence.

The research rigorously validated the homology models of PKHD-1 proteins using Z-Score analysis with PROSA Web and QMEANDisCo Global Score with ExPasy QMEANDisCo tool. These analyses provided a strong foundation for the reliability and accuracy of our modeled structures.

One of the significant aspects of the study involved predicting

the binding pockets of PKHD-1 proteins using P2Rank tool. By identifying these binding sites, the inter-action interfaces critical for understanding the protein's functional roles and possible therapeutic interventions can be done in the possible future.

The comparative analysis of PKHD-1 protein structures in different species not only enhances the understanding of evolutionary relationships but also lays the ground-work for future comparative functional studies. By elucidating structural similarities and differences, these findings pave the way for exploring species-specific functions and adaptations of PKHD-1.

This research significantly contributes to the fields of structural biology and biomedical research by providing detailed structural insights into PKHD-1 protein across multiple species. This comprehensive approach, combining homology modeling, validation techniques and binding pocket prediction, offers a valuable resource for researchers investigating potential PKHD-1 function, disease mechanisms and potential drug targeting strategies.

REFERENCES

- Bergmann C, Guay-Woodford LM, Harris PC, Horie S, Peters DJ, Torres VE. Polycystic kidney disease. *Nat Rev Dis Primers*. 2018;4(1):50.
- Ma M. Cilia and polycystic kidney disease. *Semin Cell Dev Biol*. 2021;110:139-148.
- Tahvanainen E, Tahvanainen P, Kääriäinen H, Höckerstedt K. Polycystic liver and kidney diseases. *Ann Med*. 2005;37(8):546-555.
- Zhang J, Wu M, Wang S, Shah JV, Wilson PD, Zhou J. Polycystic kidney disease protein fibrocystin localizes to the mitotic spindle and regulates spindle bipolarity. *Hum Mol Genet*. 2010;19(17):3306-3319.
- Ward CJ, Hogan MC, Rossetti S, Walker D, Sneddon T, Wang X, et al. The gene mutated in autosomal recessive polycystic kidney disease encodes a large, receptor like protein. *Nat Genet*. 2002;30(3):259-269.
- Kim I, Fu Y, Hui K, Moeckel G, Mai W, Li C, et al. Fibrocystin/polyductin modulates renal tubular formation by regulating polycystin-2 expression and function. *J Am Soc Nephrol*. 2008;19(3):455.
- UniProt: the universal protein knowledgebase in 2023. *Nucleic Acids Res*. 2023;51(D1):D523-D531.
- Gasteiger E, Hoogland C, Gattiker A, Duvaud SE, Wilkins MR, Appel RD, Bairoch A. Protein identification and analysis tools on the ExPASy server. *Humana press*. 2005.
- Garnier J, Giblat JF, Robson B. GOR method for predicting protein secondary structure from amino acid sequence. *Methods Enzymol*. 1996;266:540-553.
- Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, et al. SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Res*. 2018;46(W1):W296-W303.
- Williams CJ, Headd JJ, Moriarty NW, Prisant MG, Videau LL, Deis LN, et al. MolProbity: More and better reference data for improved all-atom structure validation. *Protein Sci*. 2018;27(1):293-315.
- Wiederstein M, Sippl MJ. ProSA-web: Interactive web service for the recognition of errors in three-dimensional structures of proteins. *Nucleic Acids Res*. 2007;35:W407-W410.
- Ramachandran GN, Ramakrishnan C, Sasisekharan V. Stereochemistry of polypeptide chain configurations. *J Mol Biol*. 1963;7:95-99.
- Krivák R, Hoksza D. P2Rank: Machine learning based tool for rapid and accurate prediction of ligand binding sites from protein structure. *J Cheminform*. 2018;10:1-2.

15. Jendele L, Krivak R, Skoda P, Novotny M, Hoksza D. PrankWeb: A web server for ligand binding site prediction and visualization. *Nucleic Acids Res.* 2019;47(W1):W345-W349.
16. Jakubec D, Skoda P, Krivak R, Novotny M, Hoksza D. PrankWeb 3: Accelerated ligand-binding site predictions for experimental and modelled protein structures. *Nucleic Acids Res.* 2022;50(W1):W593-W597.
17. Gill SC, Von Hippel PH. Calculation of protein extinction coefficients from amino acid sequence data. *Anal Biochem.* 1989;182(2):319-326.
18. Idicula-Thomas S, Balaji PV. Understanding the relationship between the primary structure of proteins and its propensity to be soluble on overexpression in *Escherichia coli*. *Protein Sci.* 2005;14(3):582-592.
19. Adhikari S, Manthana PV, Sajwan K, Kota KK, Roy R. A unified method for purification of basic proteins. *Anal Biochem.* 2010;400(2):203-206.
20. Ikai A. Thermostability and aliphatic index of globular proteins. *J Biochem.* 1980;88(6):1895-1898.
21. Sivakumar K, Balaji S, Gangaradhakrishnan. *In silico* characterization of antifreeze proteins using computational tools and servers. *J Chem Sci.* 2007;119:571-579.
22. Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of a protein. *J Mol Biol.* 1982;157(1):105-132.
23. Studer G, Rempfer C, Waterhouse AM, Gumienny R, Haas J, Schwede T. QMEAND is co-distance constraints applied on model quality estimation. *Bioinformatics.* 2020;36(6):1765-1771.