

Functional Annotation of Non-Coding RNAs through Transcriptomic Approaches

Nneka Okafor*

Department of Molecular Biosciences, University of Lagos, Lagos, Nigeria

DESCRIPTION

Functional annotation of non-coding Ribonucleic Acid (RNA) through transcriptomic approaches has become a critical area of modern molecular biology, reshaping the understanding of genome functionality beyond protein-coding genes. Although only a small fraction of the genome encodes proteins, a vast majority is actively transcribed into non-coding RNA molecules that play essential roles in gene regulation, chromatin organization, cellular signaling, and disease development. Advances in transcriptomic technologies, particularly RNA sequencing, have enabled researchers to systematically identify, quantify, and non-coding RNAs at an unprecedented scale, opening new avenues for functional annotation and biological discovery. RNA sequencing serves as the foundation for functional annotation of non-coding RNAs. By sequencing total RNA or enriched RNA fractions, researchers can capture both abundant and low-expression non-coding transcripts. Unlike traditional microarray-based methods, RNA sequencing does not rely on predefined probes, allowing discovery of novel non-coding RNA species. Long non-coding RNAs represent one of the most extensively studied classes of regulatory RNAs. These transcripts, typically longer than 200 nucleotides, are involved in chromatin remodelling, transcriptional regulation, and post-transcriptional control. Transcriptomic profiling has revealed that long non-coding RNAs exhibit highly cell-type-specific expression patterns, suggesting specialized roles in development and disease. Functional annotation studies have shown that many long non-coding RNAs act as scaffolds for chromatin-modifying complexes or as molecular decoys that regulate transcription factor activity.

Circular RNAs have also gained significant attention in recent years due to their unique covalently closed loop structure. Transcriptomic analysis has shown that circular RNAs are often expressed in a tissue-specific manner and can function as microRNA sponges, regulators of transcription, or even templates for translation in some cases. Functional annotation of circular RNAs through transcriptomics continues to expand understanding of their biological roles. Enhancer RNAs represent another emerging class of non-coding RNAs

transcribed from enhancer regions of the genome. These RNAs are often transiently expressed and are associated with active regulatory elements. Transcriptomic profiling has demonstrated that enhancer RNA expression correlates with target gene activation, suggesting a role in transcriptional regulation. Although their exact mechanisms are still being studied, enhancer RNAs are believed to contribute to chromatin looping and enhancer-promoter interactions. Single-cell transcriptomic technologies have significantly advanced the functional annotation of non-coding RNAs by enabling analysis at cellular resolution. This has revealed that non-coding RNA expression is highly heterogeneous across individual cells, even within the same tissue. Single-cell approaches have also helped identify rare cell populations expressing unique non-coding RNA signatures that may be important in development or disease.

Spatial transcriptomics further enhances functional annotation by preserving the spatial context of non-coding RNA expression within tissues. This allows researchers to map where specific non-coding RNAs are expressed and how their expression correlates with tissue architecture. Spatial analysis has revealed that certain non-coding RNAs are enriched in specific anatomical regions, such as tumor microenvironments or developing embryonic tissues, suggesting localized regulatory functions. Computational approaches are essential for interpreting transcriptomic data related to non-coding RNAs. Bioinformatics pipelines are used to assemble transcripts, quantify expression levels, and predict functional interactions. Machine learning methods are increasingly applied to classify non-coding RNAs based on sequence features, expression patterns, and structural properties. Many non-coding transcripts lack clear functional, and experimental validation is often required to confirm computational predictions. Additionally, low expression levels and high tissue specificity make detection and analysis difficult. Distinguishing functional non-coding RNAs from transcriptional noise continues to be a major area of research. Integration of multi-omics data has significantly improved the annotation process. By combining transcriptomic data with Epigenomic, proteomic, and chromatin interaction data, researchers can better infer the regulatory roles of non-coding RNAs. Correlation between non-coding RNA expression

Correspondence to: Nneka Okafor, Department of Molecular Biosciences, University of Lagos, Lagos, Nigeria, E-mail: nneka.okafor@unilag.edu.ng

Received: 01-Dec-2025, Manuscript No. TOA-25-41968; **Editor assigned:** 03-Dec-2025, PreQC No. TOA-25-41968 (PQ); **Reviewed:** 16-Dec-2025, QC No. TOA-25-41968; **Revised:** 23-Dec-2025, Manuscript No. TOA-25-41968 (R); **Published:** 30-Dec-2025, DOI: 10.35248/2329-8936.25.11.225.

Citation: Okafor N (2025). Functional Annotation of Non-Coding RNAs through Transcriptomic Approaches. Transcriptomics. 11:225.

Copyright: © 2025 Okafor N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and histone modification patterns can indicate involvement in transcriptional regulation. Similarly, interactions with RNA-binding proteins can provide functional clues. Recent technological advancements, including long-read sequencing,

have further enhanced non-coding RNA discovery by enabling full-length transcript reconstruction. This has improved the accuracy of transcript annotation and revealed complex isoform diversity among non-coding RNAs.