

Programmable RNA Editing Systems in Congenital Skeletal Dysplasia Disorders

Kavya Srinivasan*

Department of Skeletal Genetics, Bangalore Institute of Genetic Sciences, Bangalore, India

DESCRIPTION

Congenital skeletal dysplasia disorders represent a diverse group of inherited conditions affecting bone growth, cartilage formation, and connective tissue organization. These conditions arise from alterations in genes responsible for extracellular matrix production, growth plate regulation, and signaling pathways that guide skeletal morphogenesis. Clinical manifestations vary widely, ranging from disproportionate short stature and limb deformities to structural fragility of bones and abnormal spinal curvature. The severity of presentation depends on the nature of the genetic alteration and its impact on skeletal development during early embryogenesis and postnatal growth.

Conventional management approaches primarily focus on orthopedic correction, physical rehabilitation, and supportive care aimed at improving mobility and functional independence. However, these strategies do not directly modify the underlying molecular defects driving abnormal skeletal development. Advances in molecular genetics have introduced programmable RNA editing systems as a potential strategy to modulate gene expression at the transcript level without permanent alteration of genomic DNA. RNA editing systems utilize engineered molecular complexes capable of binding specific messenger RNA sequences and modifying nucleotide content post-transcriptionally. These systems allow correction of single nucleotide errors, modulation of splicing patterns, or selective suppression of harmful transcripts. Unlike Deoxyribonucleic Acid (DNA)-targeting approaches, Ribonucleic Acid (RNA)-based methods offer reversible activity, enabling temporary adjustment of gene expression profiles during critical developmental windows.

One application involves correction of transcripts encoding collagen proteins, which are essential components of bone and cartilage matrix structure. Mutations affecting collagen synthesis can lead to weakened skeletal architecture and impaired connective tissue integrity. Experimental RNA editing platforms have demonstrated the ability to partially restore normal collagen protein structure in cellular models derived from affected individuals. This restoration is associated with improved extracellular matrix organization and enhanced cellular adhesion properties. Another area of focus involves regulation of growth

plate signaling pathways. Proper skeletal elongation depends on tightly controlled communication between chondrocytes within the epiphyseal growth plate. Dysregulation of signaling molecules such as growth differentiation factors and receptor kinases can result in abnormal bone length and shape. RNA editing systems have been explored to adjust expression levels of these signaling components, restoring balance between proliferation and differentiation of cartilage-forming cells.

Delivery of RNA editing complexes to skeletal tissues presents unique challenges due to limited vascularization in cartilage regions. Systemic administration may result in uneven distribution, while localized delivery requires precise targeting techniques. Researchers have investigated nanoparticle-based carriers capable of penetrating cartilage matrix and delivering RNA complexes directly to chondrocytes. These carriers are often modified with peptides that enhance binding affinity to cartilage extracellular components. Pediatric skeletal disorders require special consideration due to ongoing growth and developmental plasticity. Early intervention during active growth phases may influence bone morphology and improve long-term structural outcomes. However, the timing of RNA-based intervention must be carefully evaluated to avoid disruption of normal developmental signaling processes that are essential for proportional skeletal formation.

Preclinical studies using animal models of skeletal dysplasia have shown that RNA editing interventions can partially normalize bone growth patterns. Treated subjects exhibit improved limb symmetry, increased bone density, and reduced incidence of deformities compared to untreated controls. Histological analysis of growth plate regions indicates more organized cartilage column structure and improved cellular alignment. Epigenetic interactions also play a role in skeletal development. RNA editing may indirectly influence chromatin states by altering expression of regulatory proteins involved in gene transcription control. These secondary effects can contribute to long-term stabilization of corrected phenotypes, although variability in response remains an area of ongoing investigation.

Single-cell transcriptomic profiling of skeletal tissue has revealed heterogeneity among chondrocyte populations within growth plates. Different cell subtypes exhibit distinct gene expression

Correspondence to: Kavya Srinivasan, Department of Skeletal Genetics, Bangalore Institute of Genetic Sciences, Bangalore, India, E-mail: kavya.srinivasan.bigs@genmail.edu

Received: 01-Jun-2026, Manuscript No. JGSGT-26-42896; **Editor assigned:** 03-Jun-2026, PreQC No. JGSGT-26-42896 (PQ); **Reviewed:** 17-Jun-2026, QC No. JGSGT-26-42896; **Revised:** 24-Jun-2026, Manuscript No. JGSGT-26-42896 (R); **Published:** 01-Jul-2026, DOI: 10.35248/2157-7412.26.16.498

Citation: Srinivasan K (2026). Programmable RNA Editing Systems in Congenital Skeletal Dysplasia Disorders. J Genet Syndr Gene Ther. 14:498.

Copyright: © 2026 Srinivasan K. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

patterns depending on their position within the cartilage zone. This heterogeneity suggests that targeted RNA editing may need to account for spatial variation in cellular function to achieve uniform therapeutic effects. Biomechanical modeling has contributed significantly to understanding how molecular changes translate into structural outcomes in skeletal tissue. Simulations of bone growth under altered gene expression conditions allow prediction of morphological changes over time. These models assist in optimizing therapeutic parameters such as dosage, timing, and distribution of RNA editing agents.

Manufacturing of RNA editing systems requires precise synthesis of guide RNA sequences and assembly of protein components responsible for nucleotide modification. Stability testing ensures

that RNA complexes maintain functional integrity during storage and delivery.

CONCLUSION

Continued advancement in programmable RNA editing technologies supports growing potential for addressing congenital skeletal dysplasia disorders at the molecular level. While challenges remain in achieving uniform tissue distribution and sustained correction, ongoing progress in delivery systems, molecular engineering, and developmental biology continues to expand therapeutic possibilities in skeletal genetics.