

Lipid Oxidation in Meat

Cheng JH*

Department of Nutrition and Health Sciences, Kainan University, Taoyuan City, Taiwan

*Corresponding author: Cheng JH, Department of Nutrition and Health Sciences, Kainan University, Taiwan, Tel: 886222193813; E-mail: jhcheng115@yahoo.com.tw

Received date: Mar 09, 2016; Accepted date: Apr 19, 2016; Published date: Apr 22, 2016

Copyright: © 2016 Cheng JH. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Lipids are primarily responsible for both desirable and undesirable flavors and aromas in meat. However, lipid oxidation in most cases deteriorates the quality of meat and causes unacceptable flavor for consumers. Lipids can be oxidized by enzymatic and non-enzymatic reactions and there are many mechanisms to explain these complex reactions in meat. Autoxidation is a continuous free-radical chain reaction and is the most important mechanism of lipid oxidation in meat. The three basic phases of this mechanism consists of initiation, propagation, and termination and these steps are usually used to explain the autoxidation system. Antioxidants are classified as (1) free radical scavengers (2) oxygen scavengers (3) chelators.

Keywords: Chelators; Oxygen; Scavengers; Antioxidants

Introduction

Lipids are primarily responsible for both desirable and undesirable flavors and aromas in meat. Basically, Nawar [1] indicated that the desirable flavor of cheese is caused by lipid oxidation. However, lipid oxidation in most cases deteriorates the quality of meat and causes unacceptable flavor for consumers. Buckley et al. [2] reported that an increase of lipid oxidation would cause a decrease of flavor, color, texture, nutritional value and acceptability in meat. There are also some other detrimental effects regarding lipid oxidation, including decrease of shelf life, increase of off-flavor, change of the functional and sensory characteristics, and sometimes formation of carcinogenic substances [3,4]. Malonaldehyde, which is a degradation product of lipid oxidation, has been criticized as a carcinogenic factor in food [5].

Lipids can be oxidized by enzymatic and non-enzymatic reactions and there are many mechanisms to explain these complex reactions in meat. However, autoxidation is a continuous free-radical chain reaction and is the most important mechanism of lipid oxidation in meat [6]. The definition of a "free radical" is an atom or molecule with an unpaired electron; therefore, they are very unstable and reactive. After free radicals attract electrons from stable compounds, the previously stable materials become free radicals. These newly formed free radicals then react with other compounds that become free radicals to react with other compounds. Therefore, these free-radical reactions work like a chain reaction [7]. It is also indicated that oxygen free radicals usually cause biological damage of lipids, nucleic acids, enzymes, and proteins. Especially, oxygen free radicals react with polyunsaturated fatty acids to produce lipid peroxides in cell membranes.

The three basic phases of this mechanism consists of initiation, propagation, and termination and these steps are usually used to explain the autoxidation system. Min [8] also outlined these three major chemical reactions of autoxidation. It is called a triplet oxygen oxidation: initiation is the formation of free alkyl radicals; propagation is the chain reaction of free alkyl radicals and peroxy radicals; and

termination is the formation of nonradical products. These reactions can be illustrated as follows:


Initiation:



Propagation:



Termination:



RH=Unsaturated fatty acid; R=Alkyl radical; ROO.=Peroxy radical; RO.=Alkoxy radical; ROOH=Hydroperoxide (Figure 1).

Hsieh and Kinsella [9] proposed that the reactions that cause decomposition of hydroperoxides to form secondary oxidative substances are very complex. Hydroperoxides, the primary products formed from autoxidation, are very unstable. It is, therefore, easy to break these products down into secondary oxidative products, such as aldehydes, alkenes, ketones and alcohols that cause off-flavors and off-odors in meat. Lillard [10] indicated that hydroperoxides, which are odorless and tasteless, do not contribute directly to flavor. Moreover, Ho and Chen [11] reported that aldehydes were the largest contributors to volatile flavors by cleavage of hydroperoxides formed from unsaturated fatty acids.

There are many factors associated with lipid oxidation in meat; for example, heat and light; catalysts; oxygen content and types of oxygen; phospholipids; unsaturated fatty acids; condition of pre-slaughter; processes that destroy muscle membranes; and pH [3,12-14]. Preventing these factors from causing oxidation is very important when making nitrite-free meat products.

Phospholipids, which are located in the cell membranes, are sensitive to oxidation in meat due to their more unsaturated fatty acids compared to other lipids. Lean meat contains a high percentage of phospholipids that makes it sensitive to oxidation [15]. Therefore, phospholipids act as the major contributors to oxidative rancidity in lean meat. However, lipid oxidation is also influenced by the degree of unsaturation of the fatty acids in the phospholipid and triglyceride fractions.

Meat from various livestock species has different sensitivity to lipid oxidation because the different species contain various proportions of unsaturated fatty acids; for example, chicken contains a higher percentage of unsaturated fatty acids and therefore has a faster rate of lipid oxidation than pork. Because pork contains more unsaturated fatty acids than beef, pork is more sensitive to lipid oxidation compared to beef. Therefore, the development of lipid oxidation varies due to the specie [16]. The processes of grinding, chopping, flaking, emulsification and cooking also accelerate lipid oxidation. The major reason is that processing liberates the membrane-bound phospholipids so they can be more easily oxidized. These processing steps disrupt muscle structures, causing unsaturated fatty acids to react with oxygen that is incorporated with the air, and to increase contact with enzymes, and heme pigments which contain metal ions such as irons that promote autoxidation in the meat systems [17]. There are, thus, many factors to influence lipid oxidation and off-flavor in muscle foods. Control of these factors is the best way to retard lipid oxidation and off-flavor in meat products.




Figure 1: Autoxidation in meat.




The chemical compounds used to delay the initiation step or slow the speed of lipid oxidation are called antioxidants [8]. However, antioxidants do not reverse autoxidation, if it occurs. On the other hand, Decker [18] indicated that antioxidants could have prooxidant activity to promote lipid oxidation under different conditions. According to their functions of inhibiting lipid oxidation, the classification of antioxidants could be primary antioxidants such as free radical scavengers or synergists that contain oxygen scavengers and chelators. The mechanism of primary antioxidants is to stabilize the free radicals by donating hydrogens to the free radicals or

accepting electrons from the free radicals with the formation of a complex. These reactions of the mechanism are as follows:

Hydrogen donation:



Formation of a complex (electron acceptance)



R.=Alkyl radical; ROO.=Peroxy radical; RO.=Alkoxy radical; AH=Antioxidant [8].

Synergistic antioxidants are the compounds used to improve the function of primary antioxidant and improve lipid stability in food. The functions of synergists are to regenerate the primary antioxidants, to react with oxygen (oxygen scavengers), and to chelate prooxidant metals (chelators) such as iron and copper [19]. Therefore, antioxidants are classified as (1) free radical scavengers (2) oxygen scavengers (3) chelators.

Consumers are demanding "label friendly" food products. Use of synthetic antioxidants could be related to health concerns; however, natural antioxidants have limited used in food systems due to special color, flavor and high cost [20]. Natural antioxidants are usually made from seed, grains, nuts, and vegetable oil; therefore, it is very important for processor to prevent the original flavor, aroma and color from the raw material of natural antioxidants that makes them unacceptable in meat products.

The ability of antioxidants for maintaining oxidative stability is influenced by the food system in which they are used. The first of practical difficulties encountered is to make a uniform dispersion at low concentration required by regulation of the antioxidants. Hydrophilic and hydrophobic characteristics of the antioxidants also affect distribution of antioxidants in meat systems. Therefore, the delivery of antioxidants in meat product becomes very important for antioxidative efficacy.

References

- Nawar WW (1996) Lipids. In Food Chemistry, (3rd edn) Fennema OR (eds), Marcel Dekker, Inc., New York.
- Buckley DJ, Gray JI, Ashgar A, Price JF, Crackel RL, et al. (1989) Effects of dietary antioxidants and oxidized oil on membral lipid stability and pork product quality. J Food Sci Off Publ Inst Food Technol 54: 1193-1197.
- Ahn DW, Wolf FH, Sim JS, Kim DH (1992) Packaging cooked turkey meat patties while hot reduces lipid oxidation. J Food Sci Off Publ Inst Food Technol 57: 1075-1077.
- Shahidi F (1994) Assessment of lipid oxidation and off-flavor development in meat and meat products. In "Flavor of meat and meat products", Shahidi F (eds). Blackie Academic & Professional, Landon, UK, p: 253.
- Kurechi T, Kikugawa K, Ozawa M (1980) Effect of malondialdehyde on nitrosamine formation. Food Cosmet Toxicol 18: 119-122.
- Pearson AM, Gray JI, Wolzak AM, Horenstein NA (1983) Safety implications of oxidized lipids in muscle foods. Food Technol 37: 121-129.

7. Hamilton RJ (1994) The chemistry of rancidity in foods. In "Rancidity in Foods", Allen JC, Hamilton RJ (Eds), Blackie Academic and Professional, an imprint of Chapman & Hall NY, p: 1-21.
8. Min DB (1998) Food Lipid. A course note. Food Science & Technology, The Ohio State University, Columbus, OH, p: 57-71.
9. Hsieh RJ, Kinsella JE (1989) Oxidation of polyunsaturated fatty acids: mechanisms, products, and inhibition with emphasis on fish. Adv Food Nutr Res 33: 233-341.
10. Lillard DA (1987) Oxidative deterioration in meat, poultry, and fish. In: "Warmed-Over Flavor in Meat," At Angelo AJ, Bailey MF (eds.) Academic Press, Orlando, FL, pp: 41-67.
11. Ho CT, Chen O (1993) Lipids in food flavors. In: "Lipids in Food flavors," Ho CT, Hartman TG (eds). American Chemical Society, Washington, DC pp: 1-15.
12. Buckley DJ, Morrissey PA, Gray JI (1995) Influence of dietary vitamin E on the oxidative stability and quality of pig meat. J Anim Sci 73: 3122-3130.
13. Stoick SM, Gray JI, Booren AM, Buckley DJ (1991) Oxidative stability of restructured beef steaks processed with oleoresin rosemary, tertiary butylhydroquinone, and sodium tripolyphosphate. J Food Sci 56: 597-600.
14. Lee SK, Mei L, Decker EA (1996) Lipid oxidation in cooked turkey as affected by added antioxidant enzymes. J Food Sci 61: 726-728.
15. Igene JO, Pearson AM, Dugan Jr LR, Price JF (1980) Role of triglycerides and phospholipids on development of rancidity in model meat systems during frozen storage. Food Chem 5: 263-265.
16. Igene JO, Pearson AM, Gray JI (1981) Effects of length of frozen storage, cooking and holding temperatures upon component phospholipids and fatty acid composition of meat triglycerides and phospholipids. Food Chem 7: 289-303.
17. Gray JI, Pearson AM (1987) Rancidity and warmed-over flavor. In: "Advances in Meat Research. Restructured Meat and Poultry Products," Pearson AM, Dutton TR (edn). Van Nostrand Reinhold Company, New York, NY, pp: 22.
18. Decker EA (1998) Strategies for manipulating the pro-oxidative/antioxidative balance of foods to maximize oxidative stability. Trends in Food Sci and Technol 9: 241-248.
19. Rajalakshmi D, Narasimhan S (1996) Food antioxidants; sources and methods of evaluation. In: "Food antioxidants: technological, toxicological, and health perspectives. Ed. by Madhavi, Deshpande DL, Salunkh SS, Inc, NY.
20. Hettiarachchy NS, Glenn KC, Gnanasambandam R, Johnson MG (1996) Natural antioxidant extract from fenugreek (*Trigonella foenumgraecum*) for ground beef patties. J Food Sci 61: 516-519.