

A Short Review on Infectious Viruses in Cultural Shrimps (*Penaeidae* Family)

Mohammedsaeed Ganjoor*

Genetic and breeding research center for cold water fishes-shahid motahari, 75914-358, Yasuj, Iran

Abstract

A major constraint limiting the shrimp production is diseases. Shrimp aquaculture is an important industry in many countries especially Southeast Asia and Iran. In cultured pond, the shrimp may be infected with several pathogens such as several viruses. There are at least six lethal viruses affecting penaeid shrimps production in the world especially Southeast Asia and Thailand. However, known viral pathogen in shrimp is about 20. They have been identified from 1970. Incidence of infection in artificial condition is more than nature. The 6 viruses are very important and they cause serious problem for shrimp cultivation and economic losses. They are consisting of HPV, IHNV, MBV, TSV, WSSV and YHV. Two of them are highly pathogenic and lethal in shrimp such as WSSV and TSV. Shrimp aquaculture is a successful activity. Despite this success, annual production decreased in the latter because of widespread epidemics (epizootics) caused by new viral pathogens. Molecular diagnostic methods such as PCR are tools to detection viral diseases in shrimp in many parts of the world. Pathological methods and electron microscopy are good tools to detection viral disease especially at the first outbreak. Sanitary methods are the best way to control and prevention of viral diseases.

Keywords: Shrimp; Viruses; Pathogen

Introduction

Shrimp farming in the Asia-Pacific region is one of the most lucrative aquaculture sectors. Asia leads the world in cultivated shrimp production with export earnings in the order of billions of US dollars per year. The pioneers for shrimp cultivation have been Japan, India, Thailand, China, Philippines, Vietnam, Iran, Ecuador, Taiwan and some other countries in the Southeast of Asia and South America and so Australia. Thailand alone has been the world's leading producer since 1992 with its export earnings alone reaching more than 1 billion US dollars per year [1-3]. Iran export shrimp. It is about 6000 metric ton annually and its value is about 30 M\$ per year. Iran not only cultivated shrimp but also is cultivating other aquatic animals such as Surgeon, Rainbow trout, Carp and Tilapia.

Viruses are the most common biological agents in the marine environment and it is known that they infect Fish, Shrimp and other aquatic animals. Marine crustaceans can be simultaneously infected by more than one type of virus [4-6].

The major viruses of concern in shrimps and fresh water shrimp are mention in the following [1,2,7,8]:

- 1) White-spot syndrome virus (WSSV or PmNOBII a mistake name which called for WSSV).
- 2) Monodon baculovirus (MBV).
- 3) Yellow-head virus (YHV).
- 4) Hepatopancreatic parvovirus (HPV).
- 5) Related Australian lymphoid organ virus (LOV).
- 6) Gill associated virus (GAV).
- 7) Infectious hypodermal and hematopoietic necrosis virus (IHNV).
- 8) Taura syndrome virus (TSV).
- 9) Mourilyan virus (MOV).

- 10) Laem Singh virus (LSNV).
- 11) Baculovirus midgut gland necrosis virus (BMNV).
- 12) Monodon slow growth syndrome (MSGs).
- 13) Infectious myonecrosis virus (IMNV).
- 14) Macrobrachium rosenbergii nodavirus (MrNV).
- 15) Extra small virus (XSV).

More than 15 viruses have been reported to infect marine shrimp [1,9]. They cause disease in shrimp specially penaeid shrimp family as species as *Penaeus monodon*, *Litopenaeus vannamei*, *Fenneropenaeus indicus*, *Litopenaeus stylirostris*, *Marsupenaeus japonicus* and etc [10-13]. Nine viruses are responsible for main considerable economic losses. These include white spot syndrome virus (WSSV), infectious hypodermal and hematopoietic necrosis virus (IHNV), monodon baculovirus (MBV), hepatopancreatic parvovirus (HPV), yellowhead virus (YHV), gill-associated virus (GAV), Taura syndrome virus (TSV), infectious myonecrosis virus (IMNV), and Mourilyan virus (MoV) [1]. Although these viruses were no cause for alarm to human health, authorities find that they were economically crippling for Asian shrimp farmers [2].

Initially, *Penaeus monodon* was the main cultivated species in Asia but this has changed markedly since 2002 when *Litopenaeus vannamei* (formerly called *Penaeus vannamei*) started to be cultivated in many Asian countries. Since 2004, it has been the main cultivated species in the world [2].

*Corresponding author: Mohammed Saeed Ganjoor, Genetic and breeding research center for cold water fishes-shahid motahari, P.O.Box:75914-358, Yasuj, Iran, Tel: 0098 919 008 1580; E-mail: ganjoor.s@gmail.com

Received April 22, 2015; Accepted June 22, 2015; Published June 25, 2015

Citation: Ganjoor M (2015) A Short Review on Infectious Viruses in Cultural Shrimps (*Penaeidae* Family). Fish Aquac J 6: 136. doi:10.4172/2150-3508.1000136

Copyright: © 2015 Ganjoor M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Viral infection found not only in cultivated shrimp but also in wild shrimp. Different viruses have found in wild shrimps for example at a research which done in Brunei waters, Over 270 *P. monodon* were collected from the South China Sea, screened and spawned. Of the nine viruses assessed, infectious hypodermal and hematopoietic necrosis virus (IHHNV) was most commonly detected (19.6%), followed by monodon baculovirus (MBV) (7.4%), hepatopancreatic parvovirus (HPV) (3.8%), and Mourilyan virus (MoV) (0.9%). The only multiple viral infections found were a combination of IHHNV and MBV (2.2%). Two most infectious viruses for *P. monodon*, white spot syndrome virus (WSSV) and yellowhead virus (YHV) were not distinguished in any shrimp [1]. Additional research in Thailand display number of viral infection in 42 shrimp samples from central and southern areas of Thailand using multiplex RT-PCR technique. Percentage of infection in examined shrimp with different viruses was: HPV 4.8%, TSV 7.1%, YHV 2.4%, MBV 2.4%, IHHNV 2.4%, WSSV 40.5%, and Mix-infection 2.4% [14].

White Spot Syndrome Virus (WSSV) is the causative agent of widespread disease related with high mortality rate in cultured shrimp [12]. It causes up to 100% mortality within 10 days in commercial shrimp farmhouses, resulting in huge losses to the shrimp farming industry [15]. About 4–6 billion US\$ of economic losses have been estimated in Asia and more than 1 billion US\$ in America, between 1992 and 2001 and presently the disease has spread worldwide. Conventional control strategies such as improvement of environmental conditions, stocking of specific pathogen free (SPF) shrimp post-larvae and augmentation of disease resistance by oral immune stimulants, are currently employed to contain WSSV infections. However, extreme virulence of this virus and its wide host range including many other crustaceans make the transmission control and prevention to be problematic [16-21].

Pathogenic DNA viruses in shrimp

DNA viruses which cause infection in shrimp contain: IHHNV, HPV, WSSV, MBV and BP.

Pathogenic RNA viruses in shrimp

RNA viruses which are infectious for shrimp contain: YHV, GAV, LOV, TSV, IMNV, MOV, MrNV, XSV and LSNV. Several positive sense RNA (+ssRNA) viruses have been reported from shrimp. Most notably, these include yellow head complex viruses (YHV), and Taura syndrome virus (TSV) [6,17,22,23].

Viruses

Properties of known viruses in shrimp summarized in below:

Monodon baculovirus (MBV)

Introduction: Even though MBV is not a severe pathogen for the black tiger prawn (*Penaeus monodon*), it should be eliminated from the farming system because it is unlikely that shrimp could carry such heavy viral infections without paying some price [5].

Virion: MBV is a DNA-virus like white spot syndrome virus (WSSV) and hepatopancreatic parvovirus (HPV). It is a double stranded DNA (dsDNA) virus [2,14,24].

Signs: MBV did not cause shrimp mortality so long as rearing conditions were good. The infection of this virus retards growth of the shrimp which named stunted-growth and resulted in economic losses [2,14,25-27].

Diagnostic: The viral inclusions can be seen directly through the

cuticle of early PL specimens using the light microscope. In tissue sections stained with hematoxylin and eosin (H&E). Transmission electron microscopy is a potent tool to recognition viral particles in specimens. Polymerase chain reaction (PCR) is a rapid method for viral detection. However, experience has shown that the Taiwanese primers do not work with MBV from Thailand and Australia [2,28,29]. Properties of the virus summarized in Table 1.

Yellow-head virus (YHV)

Introduction: YHV was first mistakenly considered to be a baculovirus but it was soon discovered during purification and characterization that its morphology differed from that of baculoviruses [30,31]. Now, it is classified as *Roniviridae* [32,33].

Virion: Rod-shaped, enveloped virion. Its genome contain positive sense single-stranded RNA (+ssRNA) [20].

Signs: Gross signs of disease that included a yellowish cephalothorax and very pale overall coloration of moribund, infected shrimp [15].

Diagnosis: Histologically, YHV infections can be easily recognized by densely basophilic inclusions, particularly in H&E stained gill sections and rapidly stained whole gills, or by staining of hemolymph smears. Diagnosis is currently best by RT-PCR method rather than in situ hybridization. There is RT-PCR recognition kit based on the work done in Thailand and Australia. The kit is not useful to detect the types of YHV found in India. In addition to nucleic acid-based tests for YHV group viruses, monoclonal antibody assays have also been developed for diagnosis by immunohistochemistry. Dot blot assay and lateral flow chromatographic assay are accessible. Additional diagnostic rule is Gold-labeled MAb (Monoclonal Antibody) test strips [2,19,34-36]. The yellow head virus properties summarized in Table 2.

White-spot syndrome virus (WSSV)

Introduction: Historically, WSSV was the second viral disease to seriously disturb Thai shrimp farmers. It has been demonstrated that spawning induces WSSV replication in *Penaeus monodon* [15,37,38].

Virion: This is a tailed, rod shaped, double stranded DNA virus with a very large circular genome in the order of 300 kbp. White spot syndrome virus belongs to the new virus family which called

Name of virion		Monodon baculovirus (MBV).
Viral family		Baculoviridae
Name of disease		Spherical baculovirus (MBV- Disease).
Host		<i>Penaeus monodon</i> .
Properties of Virion	Sensitive growth	Post larvae
	stage of shrimp.	
	(Shape and Genome)	Icosahedral, Rod-shape, dsDNA,
Epidemiology	(Outbreaks)	Mid 1980s Taiwan, 1990 Thailand,
Virulence and Signs		Acute, MBV did not cause mortality, but retards growth of the shrimp.
Current Diagnosis Methods		Pathological method (H&E Staining), Fluorescence microscopy, Monoclonal antibody, Immunohistochemistry, PCR, Nested PCR, Multiplex PCR, Dot-blot hybridization (Immuno dot-blot assay and Western blot test), in-situ hybridization, Scanning electron microscope, Transmission electron microscopy.

Table 1: Properties of MBV.

Name of virion		Yellow-head virus (YHV).
Viral family		Roniviridae
Name of disease		Yellow head disease (YHD).
Host		<i>Litopenaeus vannamei Litopenaeus stylirostris, Penaeus monodon, P. styliiferus, Macrobrachium sintangense and M. lanchesteri.</i>
Properties of Virion	(Size, Shape, Enveloped and	40-50×150-200 nm, Rod shape
	Genome)	(Bacilliform), Enveloped, +ssRNA.
Epidemiolog	Reservoir and Vectors	Invertebrates
	(Outbreaks)	1990 and 1992 Thailand
Virulence and Signs		Acute, YHV can cause high mortality in cultured shrimp. It is sometimes accompanied by the gross signs of yellowing of the cephalothorax (from which the disease got its name) and general bleaching of body color.
	Current Diagnosis Methods	Histopathology of lymphoid organs and gills, PCR, <i>in-situ</i> hybridization, Monoclonal antibody assay, Immunohistochemistry (Dot-blot assay and Lateral flow chromatographic assay), Electron microscopy.

Table 2: Properties of YHV.

Nimaviridae and genus Whispovirus [2,33,39,40]. The G+C ratio of white spot syndrome virus (WSSV) is about 41% [41].

Signs: WSS-virus is highly pathogenic affecting various crustaceans. The infection in shrimp causes mortality up to 90–100% within 3–7 days post-infection. White spot disease (WSD) has been reported to cause severe mortality in farmed shrimp especially black tiger shrimp in many countries. Mass mortalities began to be reported with characteristic gross signs of WSSV infection [2,14,18,42].

Diagnosis: Histopathology with the light and electron microscopes can indicate infection. On the basis of gross signs of disease, histopathology with the light and electron microscopes and molecular method based DNA methods used to distinguish virus. In situ DNA hybridization tests with cultivated shrimp of various species from several Asian countries that showed gross signs of white spot syndrome. PCR methods have been described for WSSV either singly [2]. Methods for real-time PCR and isothermal DNA amplification have also been described. After development of DNA hybridization probes for WSSV in Thailand primers for PCR detection of WSSV were quickly developed. In addition to PCR tests, immunological tests have also been described, lateral flow chromatographic detection strips are now accessible from both Japan and Thailand [2,14,18,42-44].

Vaccine (Prevention and Control): Trials show a new era for shrimp vaccination although shrimp immune system isn't like vertebrates. The results of some trials indicate the possibility of vaccination of kuruma shrimp with recombinant proteins against WSSV. However, new recombinant vaccine from VP28-Protein against WSSV is under studying. There is growing evidence that invertebrates, including crustaceans, possess some form of 'immune memory' or 'priming' mechanism that can be stimulated by past exposure to an infectious agent [24,45-50]. The properties of WSSV briefed in Table 3.

Hepatopancreatic parvovirus (HPV)

Introduction: HPV was first described from farmed marine shrimp in Singapore [51]. Also, It is proposed to call this virus *P. merguensis* densovirus (PmergDnv) (<http://talk.ictvonline.org/media/p/380.aspx>) [52,53]. Hepatopancreatic parvovirus (HPV) is pathogen for cultivated and wild penaeid shrimp species including *Penaeus monodon* and *Fenneropenaeus chinensis* [11,54,55]. Hepatopancreatic parvovirus infects the hepatopancreas in *Penaeus monodon* and is associated with slow growth that decreases profitability for shrimp farmers [25,52].

Virion: HPV is a single-stranded DNA (ssDNA) virus that is non-enveloped icosahedral virus averaging 22–23 nm in diameter and containing linear ssDNA. As such it belongs in the family Parvoviridae in the densovirus group. Two Asian types have been characterized at the molecular level, one in *F. chinensis* from Korea and the other in *P. monodon* from Thailand. They differ in total genome length (approximately 4 and 5 kb, respectively) [2,14,44,54-57]. HPV genome consisted of 6321 nucleotides [54]. HPV belongs to the Parvoviridae family, Parvoviruses are unique among all known viruses in having single-stranded DNA genomes which are linear. Virions are non-

Name of virion		White-spot syndrome virus (WSSV) or White spot virus (WSV).
Viral family		Nimaviridae
Name of disease		White spot disease (WSD), HHNBV.
Host		<i>Marsupenaeus japonicus, Penaeus monodon, P. penicillatus, P. semisulcatus, P. aztecus, Fenneropenaeus indicus, Litopenaeus vanname, Fenneropenaeus merguensis, Fenneropenaeus chinensis, Farfantepenaeus duorarum, Litopenaeus schmittii, L. setiferus, L. stylirostris, Metapenaeus ensis, Macrobrachium rosenbergii.</i>
Properties of Virion	(Size, Shape, Enveloped and Genome)	80-120×250-380 nm, Rod shape to elliptical, with a tail-like extension, ovoid (bacilliform), Enveloped, dsDNA, Circular 290 to 305 Kbp.
Epidemiology	Reservoir and Vectors	Invertebrates, Decapods (Crabs), Copepods, Crayfish, Aquatic insect larvae.
	(Outbreaks)	1992-1993 Asia (southeast Asia and India), 1993 Japan, 1999 Central America, (Caused pandemic epizootic).
Virulence and Signs		Very Acute, 100% mortality within 3–10 days, shrimp with WSD are reported to show a rapid reduction in food consumption, become lethargic, have a loose cuticle with some showing characteristic white spots of 0.5–2.0 mm in diameter.
Current Diagnosis Methods		Histological methods (H&E Staining), Antibody based methods, Immunohistochemical tests, <i>in-situ</i> hybridization (<i>in-situ</i> DNA hybridization), Molecular methods (PCR, One-step PCR, nested PCR, real-time PCR) and Electron microscopy, Lateral flow chromatographic test.
Appendix		The largest of the known penaeid shrimp virus. Previously called: Penaeid rod shape DNA virus (RDV), Rod shaped nuclear virus of <i>Ma. Japonicus</i> (RV-PJ), Hypodermal and hematopoietic Systemic ectodermal and mesodermal baculovirus or PmNOBII, CBV.

Table 3: Properties of WSSV.

enveloped, containing a single copy of the small (4-6 kb) viral chromosome encapsidated in a rugged icosahedral protein capsid 18-26 nm in diameter [58-61]. However, the classification of HPV viruses is still uncertain, because of their unusual capsid proteins and genome organizations [54,58]. There are dissimilar isolates for HPV. They are HPV-chinensis (HPVchin), HPVmonodon (PmDENV) and HPVsemisulcatus (HPVsemi) [53,62].

Epidemiology: HPV may have been of Indo-Pacific origin but later spread to wild shrimp in the Americas via importation of live Asian shrimp for aquaculture. In any case, it is now considered worldwide in distribution. Horizontal and Vertical transmission of HPV have been described. It is proposed that HPV may have an unknown reservoir and carriers [2,11].

Signs: The infection of this virus delays growth of the shrimp and resulted in economic losses [14,27]. Heavy infections caused in poor growth, which decreases shrimp production and without visible inflammatory response [2,25,54].

Diagnosis: Gross signs are not sufficient for HPV identification and other tests such as histological analysis or PCR testing are essential. DNA sequence analysis has revealed that there are different geographical kinds of HPV. It is recommended that different PCR primers be used for their detection [2,54,56]. In addition to PCR, monoclonal antibodies for HPV detection have recently been produced and it is hoped that a lateral flow chromatographic test will soon be available [63]. Positive diagnosis is based on the presence of hepatopancreatic lesions showing basophilic inclusions within enlarged nuclei of tubule epithelial cells, and sometimes adjacent mid-gut cells [2,11]. Traditional PCR, PCR-ELISA, real-time PCR and histopathology for detection of shrimp hepatopancreatic parvovirus (PmDENV) is applicable. Loop-mediated isothermal amplification (LAMP) combined with amplicon detection by chromatographic lateral flowdipsticks (LFD) allowed simpler detection. In situ hybridization is another method to diagnosis [52,54]. A number of diagnostic methods were established for detection of this virus including histological method (H&E staining), Transmission electron microscopic (TEM), in situ hybridization, and polymerase chain reaction (PCR) [1,2,48,64]. Properties of the virus summarized in Table 4.

Infectious hypodermal and hematopoietic necrosis virus (IHHNV)

Introduction: The virus was first described in blue shrimp *Litopenaeus stylirostris* and white shrimp *L. vannamei* (Formerly called *Penaeus stylirostris* and *P. vannamei*) in the Americas in the early 1980s [2,65,66].

Virion: IHHN-virus is a non-enveloped, icosahedral virus with 22–23 nm in diameter and holding linear ssDNA of 4.1 kb. IHHNV is single-stranded DNA virus. Thus it is a typical densovirus classified in the Parvoviridae family [14,44,54,58,67,68].

Signs: Although IHHNV is low virulence virus in adult shrimp [37]. But, it causes acute epidemics and mass mortality only with juveniles and sub-adults of *L. stylirostris* [2,67]. The infection of this virus retards growth of shrimp so resulted in economic losses [14]. In *L. vannamei*, it causes reduced, irregular growth and cuticular deformities that are collectively referred to as “runt-deformity syndrome” (RDS) [69-72]. The infectious hypodermal and hematopoietic necrosis virus is very pathogenic for *Litopenaeus stylirostris* whereas infection in *Litopenaeus vannamei* is known to induce development and growth abnormalities and cause economic losses that range between 10% and 50% [12,37].

Diagnosis: IHHNV can be identified by routine histological technique with H&E staining and in situ DNA hybridization assays with a specific IHHNV probe. A polymerase chain reaction (PCR) assay is also described in the OIE Aquatic Animal Health Manual. By TEM icosahedral virions can be seen in the cytoplasmic region of infected cells [2,73, 74]. Several clinical evaluations had shown that hemolymph is the best tissue sample for reliable diagnosis of IHHNV infection [37].

Epidemiology: In *L. stylirostris* the virus can transmit by vertical and horizontal route. IHHNV vertical transmission from infected females was clearly established [37,67]. Properties of this virus briefed in Table 5.

Name of virion		Hepatopancreatic parvovirus (HPV) or <i>Penaeus monodon</i> densovirus (PmDENV and PstDENV).
Viral family		Parvoviridae
Name of disease		HPI (Hepatopancreatic parvovirus infection)
Host		<i>Penaeus monodon</i> , <i>Macrobrachium rosenbergii</i> , <i>Marsupenaeus japonicus</i> , <i>Fenneropenaeus merguensis</i> , <i>Fenneropenaeus indicus</i> , <i>Fenneropenaeus chinensis</i> , <i>Penaeus orientalis</i> .
Properties of Virion	Sensitive growth stage of shrimp.	Larvae, Post larvae and Juvenile.
	(Size, Shape, Enveloped and Genome)	22-24 nm, Isometric (Icosahedral), Nonenveloped, -ssDNA, Linear, 4-5 Kbp, There are different geographical type of HPV.
Epidemiology	Reservoir and Vectors	Vertebrates, Invertebrates
	(Outbreaks, Transmission route)	1980s Singapore, Horizontal and Vertical.
Virulence and Signs		Slow growth that reduces profitability for shrimp farmers (reduced growth rates of prawns during the juvenile stages and overt mortalities). However, there are no specific gross signs for HPV so diagnosis may be difficult, High acute for larvae (The virus is lethal to shrimp larvae during the first month after stocking).
Current Diagnosis Methods		Histological examination (H&E staining), PCR and nested-PCR, PCR- ELISA detection method, The loop-mediated isothermal amplification (LAMP), Transmission electron microscopy (TEM), in-situ hybridization.
Appendix		The Australian HPV isolate from <i>P. merguensis</i> is the fourth strain of penaeid prawn HPV to be partially sequenced. We have therefore proposed to name this virus <i>P. merguensis</i> densovirus (PmergDENV), following the convention of the International Committee for the Taxonomy of Viruses. The other three are HPVchin from <i>P. chinensis</i> of Korea, PmDENV from <i>P. monodon</i> of Thailand and HPVsemi from <i>P. semisulcatus</i> of India. An additional strain of HPV has been reported in the freshwater prawn <i>M. rosenbergi</i> .

Table 4: Properties of HPV.

Taura syndrome virus (TSV)

Introduction: Taura syndrome was first known as a new disease in the Americas in 1992 but its viral etiology was not established until 1994 [75-77]. It is a recent viral pathogenic agent to arrive on the Asian scene [11,78,79].

Virion: It is a naked (without envelop) 32 nm icosahedral virus containing ssRNA molecule with 10.2 kb length and positive sense [80]. However, it was later allocated to the family Dicistroviridae near the genus Cripavirus (cricket paralysis virus) [32,81]. GC content of the viral RNA ranging from 35 to 45%. RNA constitutes about 30% of the virion weight. A small genome-linked virus protein (VPg), is covalently attached to the 5' end of the genome [41].

Signs: Experimental bioassays have exposed that low-grade mortalities can occur and that *P. monodon* can carry asymptomatic infections [1,82,83]. Properties of TSV summarized in Table 6.

Mourilyan virus (MoV)

Introduction: This virus was discovered accidentally during the study of GAV, an Australian virus from the yellow head virus complex [84].

Epidemiology: It appears to be endemic in populations of *Penaeus monodon* from Queensland in Australia, from Malaysia and from Thailand [2].

Name of virion		Infectious hypodermal and hematopoietic virus (IHNV).
Viral family		Parvoviridae
Name of disease		Infectious hypodermal and hematopoietic necrosis (IHNV) OR: Runt-deformity syndrome (RDS).
Host		<i>Litopenaeus stylirostris</i> , and <i>Litopenaeus vannamei</i> .
Properties of Virion	Sensitive growth stage of shrimp.	Juvenile and Subadult.
	(Size, Shape, Enveloped and Genome)	22-23 nm, Icosahedral, Nonenveloped, ssDNA, Linear, Length: 4.1 Kb, Variants: IHNV-I, IHNV-II, IHNVIII
Epidemiology	Reservoir and Vectors	Shrimp
	(Outbreaks, Transmission route)	1980s Taiwan & 1981 America, Pass onto other population by Horizontal & Vertical transmission.
Virulence and Signs		Acute epizootics and mass mortalities (>90%) in <i>L. stylirostris</i> , (In <i>L. vannamei</i> , It cause reduced-irregular growth and cuticular deformities on the other hand reported that it is asymptomatic without significant mortalities in <i>L. vannamei</i>)
Current Diagnosis Methods		It cause reduced-irregular growth and cuticular deformities on the other hand reported that it is asymptomatic without significant mortalities in <i>L. vannamei</i>) Histopathological staining and examination, in-situ DNA hybridization, Dot-Blot test, PCR, real-time PCR.
Appendix		The smallest of the known penaeid shrimp viruses.

Table 5: Properties of IHNV.

Name of virion		Taura syndrome virus (TSV).
Viral Family		Dicistroviridae (Previously classified as: <i>Picomiviridae</i>).
Name of disease		Taura syndrome disease (TSD).
Host		<i>Litopenaeus vannamei</i> , <i>Litopenaeus stylirostris</i> , <i>L. setiferus</i> , <i>L. schmitti</i> . Other penaeid (<i>Farfantepenaeus aztecus</i> , <i>Fa. duorarum</i> , <i>Fenneropenaeus chinensis</i> , <i>Penaeus monodon</i> and <i>Marsupenaeus japonicus</i>) have been experimentally infected. <i>Penaeus monodon</i> .
Properties of Virion	Sensitive growth stage of shrimp.	Post larvae, Juvenile and Sub adult.
	(Size, Shape, Enveloped and Genome)	32 nm, Isometric (Icosahedral), Nonenveloped, +ssRNA, 10.2 Kb.
Epidemiology	Reservoir and Vectors	Invertebrates, Shrimp eating birds and Flying aquatic insects.
	(Outbreaks, Transmission route)	1991-1992 Ecuador & 1999 Asia (1998 Taiwan), Horizontal and Vertical, Cause pandemic epizootic.
Virulence and Signs		Acute, Cumulative mortalities due to TSV epizootics have ranged from 40 to >90% in cultured population of <i>L. vannamei</i> .
Current Diagnosis Methods		Pathological methods (H&E staining), Monoclonal antibody based methods, in-situ hybridization, Dot-blot method (Western blot), Immunohistochemistry methods, DNA amplification method (PCR).

Table 6: Properties of TSV.

Disease: There is some indication that it may be associated with gradual, progressive mortality in pond-reared *Marsupenaeus japonicus* in Australia [85].

Virion: MoV size is about 85-100 nm in diameter. It is an enveloped virus with spherical to ovoid virions. The MoV genome is contained of single-stranded, negative sense RNA divided into 4 fragments. The morphology, genome type, genome fragmentation and genome organization most closely look like features of viruses in the family Bunyaviridae [2].

Diagnosis: A nested RT-PCR method has been developed [86].

Table 7 shows the properties of MoV.

Laem singh virus (LSNV)

Introduction: Laem Singh virus (LSNV) is a new shrimp virus from Thailand [6].

Virion: This virus hasn't envelopment. Its shape is icosahedral which size is about 27 nm diameter, similar to the size of viruses in the family Luteoviridae. On the other hand, LSNV is like family Barnaviridae based on some its properties [2].

Diagnosis: In situ hybridization test and RT-PCR apply to detect infected shrimp [2]. Features of LSNV summarized in Table 8.

Baculovirus midgut gland necrosis virus (BMNV)

Introduction: BMNV was acute pathogen of larval stages of *M. japonicus* in Japan in the early period of shrimp culture development but was excluded from the cultivation system after the mode of transmission from infected broodstock was established, and thorough washing of the eggs or nauplii was employed as a routine preventative measure [87]. Characteristics of BMNV summarized in Table 9.

Name of virion		Mourilyan virus (MOV).
Viral family		The morphology, genome type, genome fragmentation and genome organization most closely resemble features of viruses in the family Bunyaviridae.
Host		<i>Penaeus monodon</i> , <i>Marsupenaeus japonica</i> .
Properties of Virion	(Size, Shape, Enveloped and Genome)	85-100 nm, Spherical (ovoid), Enveloped, -ssRNA (4 Fragments).
Epidemiology	(Outbreaks)	Australia, Malaysia, Thailand.
Virulence and Signs		There is some indication that it may be associated with gradual, progressive mortality in pond-reared <i>P. japonicus</i> in Australia. The link to specific disease in <i>P. monodon</i> is less clear.
Current Diagnosis Methods		Nested RT-PCR
Appendix		This virus was discovered accidentally during the study of GAV.

Table 7: Properties of MOV.

Name of virion		Laem Singh virus (LSNV).
Viral family		It is mostly from the family Luteoviridae. but also to mushroom bacilliform virus (Luteoviridae or Barnaviridae).
Name of disease		Monodon slow growth syndrome (MSGS)
Host		<i>Penaeus monodon</i> .
Properties of Virion	Sensitive growth stage of shrimp. (Size, Shape, Enveloped and Genome)	27 nm, Icosahedral, Nonenveloped, More genome information will be needed for proper classification of this virus.
Epidemiology	(Outbreaks)	2002 Asia.
Virulence and Signs		Probably, It cause monodon slow growth syndrome (MSGS) but more research will be needed.
Current Diagnosis Methods		<i>In-situ</i> hybridization, RT-PCR, Electron microscopy.
Appendix		Tests using both in situ hybridization and RT-PCR revealed the presence of LSNV in both MSGS ponds and normal growth ponds, indicating that it was probably not the direct cause of MSGS. There still remains the possibility the MSGS is related to the prevalence or severity of MSGS infections in a shrimp culture pond.

Table 8: Properties of LSNV.

Monodon slow growth syndrome (MSGS caused by laem singh virus (LSNV))

Introduction: Monodon slow growth syndrome (MSGS) was first observed by shrimp farmers in cultured black tiger shrimp in 2002 [6,88]. It may be a viral disease of shrimp. Several scientists acclaim that Laem Singh virus (LSNV) is an agent that cause MSGS [6].

Signs: Unusual slow growth in cultivated *P. monodon*. Properties of LSN-virus summarized in Table 10.

Infectious myonecrosis virus (IMNV)

Introduction: IMNV is the most recent of the identified shrimp viruses to arrive in Asia, and is thought to have been introduced with contaminated *L. vannamei* stocks from Brazil [1,89,90]. While

L. vannamei is the primary host for IMNV, other species such as *P. monodon* are vulnerable by experimental infection [1,91].

Signs: No gross signs or mortalities have been reported [1]. Features of the virus summarized in Table 11.

Gill associated virus (GAV)

Introduction: Gill-associated virus (GAV) is a RNA virus. In Australia, Gill-associated virus has been linked to morbidity and mortalities in cultivated *Penaeus monodon* [92-95]. GAV also can to cause disease and mortalities similar to that caused by the more highly virulent Yellow-head virus (YHV) that continues to cause production losses in shrimp farmed in South-east Asia. GAV has a 26.2 kb ssRNA genome and is the type species of the Okavirus genus in the Roniviridae [92]. Table 12 show properties of gill-associated virus.

Macrobrachium rosenbergii nodavirus (MrNV)

Macrobrachium rosenbergii is fresh water shrimp. White tail disease (WTD) caused by both *Macrobrachium rosenbergii*-nodavirus (MrNV) and extra small viruses (XSV). MrNV is a satellite virus particles are found in *Macrobrachium rosenbergii* (giant river prawn) infected with *Macrobrachium rosenbergii* nodavirus (MrNV; a virus not yet classified but clearly related to viruses in the family *Nodaviridae*) [41]. WTD is a major problem. It is responsible for severe mortality in post-larvae of *M. rosenbergii* in the hatcheries and nurseries [8,96]. White tail disease has been observed in freshwater prawn hatcheries and nursery

Name of virion		Baculovirus midgut gland necrosis virus
Viral family		Baculoviridae
Name of disease		BMN (It have five names more).
Host		<i>Marsupenaeus japonicus</i> , <i>Penaeus monodon</i> .
Properties of Virion	(Size)	36×250 nm
Virulence and Signs		Acute

Table 9: Properties of BMNV.

Name of virion		Laem Singh virus (LSNV).
Viral family		It is mostly from the family Luteoviridae. but also to mushroom bacilliform virus (Luteoviridae or Barnaviridae).
Name of disease		Monodon slow growth syndrome (MSGS)
Host		<i>Penaeus monodon</i> .
Properties of Virion	(Size, Shape, Enveloped and Genome)	27 nm, Icosahedral, Nonenveloped, More genome information will be needed for proper classification of this virus.
Epidemiology	(Outbreak)	2002 Asia.
Virulence and Signs		Probably, It cause monodon slow growth syndrome (MSGS) but more research will be needed.
Current Diagnosis Methods		<i>In-situ</i> hybridization, RT-PCR, Electron microscopy.
Appendix		Tests using both in situ hybridization and RT-PCR revealed the presence of LSNV in both MSGS ponds and normal growth ponds, indicating that it was probably not the direct cause of MSGS. There still remains the possibility the MSGS is related to the prevalence or severity of MSGS infections in a shrimp culture pond.

Table 10: Properties of LSNV which cause MSGS.

Name of virion		Infectious myonecrosis virus (IMNV).
Viral family		Totiviridae (A toti-like virus)
Name of disease		Infectious myonecrosis (IMN).
Host		<i>Litopenaeus vannamei</i> .
Properties of Virion	Sensitive growth stage of shrimp.	Juvenile and sub adult.
	(Size, Shape and Genome)	40 nm, Icosahedral, A single dsRNA with 7560 bp.
Epidemiology		(Outbreaks) 2004 America & 2006 Asia,
Virulence and Signs		IMN presents as a disease in <i>L. vannamei</i> with an acute onset of gross signs and elevated mortalities, but it progresses with a more chronic course accompanied by persistent moderate mortalities.
Current Diagnosis Methods		Histological examination (H&E staining), PCR (One step PCR, Nested PCR, real-time PCR), in-situ hybridization.

Table 11: Properties of IMNV.

Name of virion		Gill associated virus (GAV), GAV is the Australian strain of YHV.
Viral family		Roniviridae
Name of disease		Yellow head disease (YHD).
Host		<i>Penaeus monodon</i> .
Properties of Virion	(Genome)	+ssRNA, 26.2 Kb.
Current Diagnosis Methods		See: YHV details.
Appendix		LOV and GAV share approximately 95% DNA sequence identity and 100% amino acid identity, establishing that they are the same virus type, while GAV and YHV share approximately 85% DNA sequence identity and 96% amino acid identity indicating that they are different types.

Table 12: Properties of GAV.

ponds in different parts of India, causing high mortalities and huge economic losses [96,97].

Virion: MrNV is a small, icosahedral, non-enveloped virus. Its size is 26–27 nm in diameter. The genome of MrNV composed of two pieces of ssRNA (RNA1 and RNA2) of 2.9 and 1.26 kb, respectively, and there is a single polypeptide of 43 kDa in the capsid [8,98].

Signs: MrNV cause white tail disease (WTD) of freshwater prawns [8].

Diagnosis: A number of diagnostic methods have been established to identify this virus including histopathology, immunological methods, reverse transcriptase-polymerase chain reaction technique (RT-PCR) and in-situ dot blot hybridization method using nucleic acid probes. A sandwich enzyme-linked immunosorbent assay have been developed to detect MrNV in freshwater prawns [99]. Recently, genome-based methods, dot-blot hybridization and RT-PCR have been developed to detect MrNV [8,100].

Epidemiology: White tail disease was first described in the French West Indies, later in China, India, then in Thailand and recently in Taiwan [97,98,101-103]. Characteristics of the virus summarized in Table 13.

Extra small virus (XSV)

Introduction: The XSV (extra small virus) is a satellite virus particles are about 15 nm in diameter and serologically unrelated to those of MrNV. XSV is a positive-sense single-stranded RNA, about 800 bases in size, encoding a 17 kDa capsid protein. The mixed infection of MrNV and XSV is implicated in white spot disease of prawns. Extra small virus (XSV) may cause disease of shrimp and may produce white tail disease (WTD). It is may responsible for mortality in post-larvae of *M. rosenbergii* in the hatcheries and nurseries [8,96]. It has reported the presence of XSV in addition to MrNV in WTD-infected postlarvae of freshwater prawns in India. XSV does not cause mortality in marine shrimp as observed in adult freshwater prawn [8,104].

Virion: XSV is a virus-like particle (A satellite virus particles), icosahedral in shape and 15 nm in diameter, with a linear ssRNA [98].

Diagnosis: Various methods have been developed to detect this virus including: histopathology, immunological methods, reverse transcriptase-polymerase chain reaction technique (RT-PCR), in-situ dot blot hybridization method, sandwich enzyme-linked immunosorbent assay, genome-based methods, dot-blot hybridization and RT-PCR [8,97,100,104]. Table 14 show properties of the virus.

Miscellaneous other viruses

A number of other viruses have been reported from cultivated shrimp in Asia and have been adequately covered in previous reviews. However, with the exception of baculovirus midgut gland necrosis virus (BMNV), none have been reported to be the cause of serious or widespread economic losses, and they are not covered in this review.

Result and Conclusion

It is about 20 viruses which have found in shrimps from 1980 to

Name of virion		<i>Macrobrachium rosenbergii</i> nodavirus (MrNV).
Viral family		<i>Nodaviridae</i>
Name of disease		White tail disease (WTD).
Host		<i>Macrobrachium rosenbergii</i> (Freshwater prawns).
Properties of Virion	Sensitive growth stage of shrimp.	Larvae and Postlarvae.
	(Size, Shape, Enveloped and Genome)	26–27 nm, Icosahedral, Nonenveloped, ssRNA, (2 pieces, 2.9 and 1.26kb).
Epidemiology	Reservoir and Vectors	It is possible of the marine shrimp (<i>Fenneropenaeus indicus</i> , <i>Marsupenaeus japonicus</i> and <i>Penaeus monodon</i>) acting as reservoir for MrNV.
	(Outbreaks)	French West Indies, Taiwan, China, India.
Virulence and Signs		Causing high mortalities and huge economic losses in hatcheries and nursery ponds.
Current Diagnosis Methods		Histopathology, immunological methods, reverse transcriptase-polymerase chain reaction technique (RT-PCR) and in-situ dot blot hybridization method.
Appendix		The results of a study indicate the possibility of marine shrimp acting as reservoir for MrNV and XSV and maintaining their virulence in tissue system of marine shrimp.

Table 13: Properties of MrNV.

Name of virion		Extra small virus (XSV).
Viral family		No classified. It is a satellite virus particles.
Name of disease		White tail disease (WTD).
Host		<i>Macrobrachium rosenbergii</i> (Freshwater prawns).
Properties of Virion	Sensitive growth stage of shrimp.	Larvae and Postlarvae.
	(Size and Genome)	15 nm, ssRNA, Linear.
Epidemiology	Reservoir and Vectors	It is possible of the marine shrimp (<i>Fenneropenaeus indicus</i> , <i>Marsupenaeus japonicus</i> and <i>Penaeus monodon</i>) acting as reservoir for XSV.
	(Outbreaks)	China & India.
Current Diagnosis Methods		Histopathology, immunological methods, reverse transcriptase-polymerase chain reaction technique (RT-PCR) and in-situ dot blot hybridization method.
Appendix		Have reported the presence of XSV in addition to MrNV in WTD-infected postlarvae of freshwater prawns in India.

Table 14: Properties of XSV.

2011 as well as pathogens. On the other words, a new type of viral pathogen found in shrimp every 1-2 years. Shrimp cultivation is going to progress more and more in several countries and scientists have been found new viruses more, every several years. The most important viral pathogens in shrimp are WSSV, YHV, MBV, TSV, IHNV and HPV which some of them cause severe mortality in ponds. There is not report with human illness by viral disease in shrimps, but it is need more research.

Viruses of shrimps often cause no gross signs of disease in shrimp, especially in the natural environment. In stressful environments such as culture systems, some of these viruses can become more virulent and cause significant economic loss by mortality or retarded growth. But some of them are really lethal such as WSSV [4,13].

Viruses outbreak which had been seen and reported are contain: 1992 Thailand (HPV), 1995 Thailand (YHV), 1996-7 Thailand (WSSV), 1993 Japan and China (WSSV), 1980s Taiwan (MBV), 1990 Thailand (MBV), 1984 Singapore (HPV), 2003 India (HPV), 1980s America (IHNV), 1992 America (TSV), 2002 Thailand (MSGs), 1999 French West Indies (WT-Disease). In some cases it is possible the ponds be infected with 2 or more viruses simultaneously. Shrimp feeding behavior specially cannibalism may also be a serious problem for shrimp farmers because it caused horizontal transmission of viruses [2].

Methods which applied to detection viral disease in shrimp is different they are listed in the following:

1. Histology (H&E staining- Light Microscopy).
2. TEM (Transmission electron microscopy).
3. Non-nested PCR.
4. Nested PCR.
5. Multiplex PCR.
6. Multiplex reverse transcription-polymerase chain reaction (mRT-PCR).
7. Real-time RT-PCR.

8. Multiplex RT-nested PCR.
9. Miniarray.
10. Single-step multiplex PCR.
11. Single PCR.
12. ELISA (Monoclonal antibody assay based test).
13. PCR-ELISA.
14. Fluorescence microscopy (Eosin formula used contains some phloxine dye to detect occlusion bodies).
15. *In situ* hybridization (A type of Nucleic acid-based test).
16. Monoclonal antibody assay based tests: (ELISA, Dot blot assay, Lateral flow chromatographic assay).

The best way to control and prevention of viral diseases is following of hygienic rules in breeding, nursery and ponds as well as sanitary methods. Application and improvement of SPF stocks is another way. Also, stocking and Cultivation SPR shrimp may be useful. Although immune system in shrimp is primitive but vaccination may be useful method for prevention of viral disease in future.

References

1. Claydon K, Tahir RAH, Said HM, Lakim MH, Tamat W (2010) Prevalence of shrimp viruses in wild *Penaeus monodon* from Brunei Darussalam. *Aquaculture* 308: 71-74.
2. Flegel TW (2006) Detection of major penaeid shrimp viruses in Asia, a historical perspective with emphasis on Thailand. *Aquaculture* 258: 1-33.
3. De Silva SS, Mohan CV, Phillips MJ (2007) A different form of dumping: The need for a precautionary approach for yet another new species for shrimp farming in Asia. *Aquaculture Asia*. Network of Aquaculture Centres in Asia-Pacific.
4. Flegel TW (2001) The shrimp response to viral pathogens. In: *The New Wave, Proceedings of the special session on sustainable shrimp aquaculture*, World Aquaculture Society, Orlando, Boca Raton.
5. Flegel TW, Nielsen L, Thamavit V, Kongtim S, Pasharawipat T (2004) Presence of multiple viruses in non-diseased, cultivated shrimp at harvest. *Aquaculture* 240: 55-68.
6. Sritunyalucksana K, Apisawetakan S, Boon-nat A, Withyachumnarkul B, Flegel TW (2006) A new RNA virus found in black tiger shrimp *Penaeus monodon* from Thailand. *Virus Res* 118: 31-38.
7. Lightner DV (2011) Virus diseases of farmed shrimp in the Western Hemisphere (the Americas): A review. *Journal of Invertebrate Pathology* 106: 110-130.
8. Sudhakaran R, Musthaq SS, Haribabu P, Mukherjee SC, Gopal C, Hameed ASS (2006) Experimental transmission of *Macrobrachium rosenbergii* nodavirus (MrNV) and extra small virus (XSV) in three species of marine shrimp (*Penaeus indicus*, *Penaeus japonicus* and *Penaeus monodon*). *Aquaculture* 257: 136-141.
9. Bonami JR (2008) Shrimp Viruses. In: Mahy, B.W.J., Regenmortel, M.H.V.v. (Eds.), *Encyclopedia of Virology*. Academic Press, Oxford, pp. 567-576.
10. Loh PC, Tapay LM, Lu Y, Nadala EC (1997) Viral pathogens of the penaeid shrimp. *Adv Virus Res* 48: 263-312.
11. Lightner DV (1996) *A Handbook of Pathology and Diagnostic Procedures for Diseases of Penaeid Shrimp*. World Aquaculture Society, Baton Rouge, LA.
12. Lightner DV, Redman RM (1998) Shrimp diseases and current diagnostic methods. *Aquaculture* 164, 201-220.
13. Sritunyalucksana K, Srisala J, McColl K, Nielsen L, Flegel TW (2006b) Comparison of PCR testing methods for white spot syndrome virus (WSSV) infections in penaeid shrimp. *Aquaculture* 255: 95-104.
14. Khawsak P, Deesukon W, Chaivisuthangkura P, Sukhumsirichart W (2008) Multiplex RT-PCR assay for simultaneous detection of six viruses of penaeid shrimp. *Molecular and Cellular Probes* 22: 177-183.

15. Flegel TW (1997) Special topic overview; major viral diseases of the black tiger prawn (*Penaeus monodon*) in Thailand. World J Microbiol Biotechnol 13: 433-42.
16. Rout N, Kumar S, Jaganmohan S, Murugan V (2007) DNA vaccines encoding viral envelope proteins confer protective immunity against WSSV in black tiger shrimp. Vaccine 25: 2778-2786.
17. Robles-Sikisaka R, Garcia DK, Klimpel KR, Dhar AK (2001) Nucleotide sequence of 3'-end of the genome of Taura syndrome virus of shrimp suggests that it is related to insect picornaviruses. Arch Virol 146: 941-952.
18. Kanchanaphum P, Wongteerasupaya C, Sitidilokratana N, Boonsaeng V, Panyim S, Tassana, et al. (1998) Experimental transmission of white spot syndrome virus (WSSV) from crabs to shrimp *Penaeus monodon*. Dis Aquat Organ 34: 1-7.
19. Wongteerasupaya C, Boonsaeng V, Panyim S, Tassanakajon, Withyachumanankul B, Flegel TW (1997) Detection of yellowhead virus (YHV) of *Penaeus monodon* by RT-PCR amplification. Dis Aquat Org 31: 181-186.
20. Wongteerasupaya C, Sriurairatana S, Vickers JE, Anutara A, Boonsaeng V, Panyim S, et al. (1995a) Yellow-head virus of *Penaeus monodon* is an RNA virus. Dis Aquat Org 22: 45-50.
21. Wongteerasupaya C, Vicker JE, Sriurairatana S, Nash GL, Akarajamorn A, and Boonsaeng V, et al. (1995b) A non-occluded, systemic baculovirus that occurs in cells of ectodermal and mesodermal origin and causes high mortality in the black tiger prawn, *Penaeus monodon*. Dis Aquat Organ 21: 69-77.
22. Walker PJ, Cowley JA, Spann KM, Hodgson RAJ, Hall MR (2001) Yellow head complex viruses: transmission cycles and topographical distribution in the Asia-Pacific region. In: Browdy, C.L., Jory, D.E. (Eds.), The New Wave, Proceedings of the special session on sustainable shrimp culture, Aquaculture (2001) World Aquaculture Society, Baton Rouge, LA, 292-302.
23. Mari J, Bonami JR, Lightner DV (1998) Taura syndrome of penaeid shrimp: cloning of viral genome fragments and development of specific gene probes. Dis Aquat Org 33: 11-17.
24. Chang PS, Lo CF, Kou GH, Lu CC, Chen SN (1993) Purification and amplification of DNA from *Penaeus monodon*-type baculovirus (MBV). J Invertebr Pathol, 62: 116-20.
25. Flegel TW, Thamavit V, Pasharawipas T, Alday-Sanz V (1999) Statistical correlation between severity of hepatopancreatic parvovirus (HPV) infection and stunting of farmed black tiger shrimp (*Penaeus monodon*). Aquaculture 174: 197- 206.
26. Fegan DF, Flegel TW, Sriurairatana S, Waiakrutra M (1991) The occurrence development and histopathology of monodon baculovirus in *Penaeus monodon* in Southern Thailand. Aquaculture 96: 205-217.
27. Tang KF, Lightner DV, (2006) Infectious hypodermal and hematopoietic necrosis virus (IHHNV)-related sequences in the genome of the black tiger prawn *Penaeus monodon* from Africa and Australia. Virus Res, 118: 185-91.
28. Satidkanitkul A, Sithigorngul P, Sang-oum W, Rukpratanporn S, Sriurairatana S, Withayachumnarnkul B, Flegel TW (2005) Synthetic peptide used to develop antibodies for detection of polyhedrin from monodon baculovirus (MBV). Dis Aquat Org 65: 79-84.
29. Belcher CR (1998) Molecular detection and characterisation of monodon baculovirus. Thesis, University of Queensland, Brisbane.
30. Boonyaratpalin S, Supamattaya K, Kasornchandra J, Direkbusaracom S, Ekpanithanpong U, et al. (1993) Nonoccluded baculo-like virus, the causative agent of yellow head disease in the black tiger shrimp (*Penaeus monodon*). Fish Pathol 28: 103-109.
31. Chantanachookin C, Boonyaratpalin S, Kasornchandra J, Direkbusarakom S, Ekpanithanpong U, et al. (1993) Histology and ultrastructure reveal a new granulosis-like virus in *Penaeus monodon* affected by "yellow-head" disease. Dis Aquat Org 17: 145-157.
32. Fauquet CM, Mayo MA, Maniloff J, Desselberger U, Ball LA (2004) 8th Report of the International Committee on Taxonomy of Viruses. Elsevier, Amsterdam.
33. Mayo MA (2002) A summary of taxonomic changes recently approved by ICTV. Arch Virol 147: 1655-1656.
34. Cowley JA, Cadogan LC, Wongteerasupaya C, Hodgson RAJ, Boonsaeng V, et al. (2004) Multiplex RT-nested PCR differentiation of gill-associated virus (Australia) from yellow head virus (Thailand) of *Penaeus monodon*. J Virol Methods 117: 49-59.
35. Sithigorngul P, Rukpratanporn S, Longyant S, Chaivisuthangkura P, Sithigorngul W, et al. (2002) Monoclonal antibodies specific to yellow-head virus (YHV) of *Penaeus monodon*. Dis. Aquat Org 49: 71-76.
36. Sithigorngul P, Chauyuchwong P, Sithigorngul W, Longyant S, Chaivisuthangkura P, et al. (2000) Development of a monoclonal antibody specific to yellow head virus (YHV) from *Penaeus monodon*. Dis Aquat Org 42: 27-34.
37. Mottea E, Yugchaa E, Luzardo J, Castro F, Leclercq G, et al. (2003) Prevention of IHHNV vertical transmission in the white shrimp *Litopenaeus vannamei*. Aquaculture 219: 57-70.
38. Peng SE, Lo CF, Lin SC, Chen LL, Chang YS, et al. (2001) Performance of WSSV-infected and WSSV-negative *Penaeus monodon* postlarvae in culture ponds. Dis Aquat Org 46: 165-172.
39. ICTV (2013) <http://ictvonline.org/index.asp>
40. Du H, Xu Z, Wu X, Li W, Dai W (2006) Increased resistance to white spot syndrome virus in *Procambarus clarkii* by injection of envelope protein VP28 expressed using recombinant baculovirus. Aquaculture 260: 39-43.
41. King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (2012) Virus Taxonomy (Classification and Nomenclature of Viruses, Ninth Report of the International Committee on Taxonomy of Viruses). Elsevier academic press, Amsterdam p. 1273.
42. Wongteerasupaya C, Wongwisansri S, Boonsaeng V, Panyim S, Pratanpipat P, et al. (1996) DNA fragment of *Penaeus monodon* baculovirus PmNOBII gives positive in situ hybridization with viral infections in six penaeid shrimp species. Aquaculture 143: 23-32.
43. Kono T, Savan R, Sakai M, Itami T (2004) Detection of white spot syndrome virus in shrimp by loop-mediated isothermal amplification. J Virol Methods 115: 59-65.
44. Dhar AK, Roux MM, Klimpel KR (2001) Detection and quantification of infectious hypodermal and hematopoietic necrosis virus and white spot virus in shrimp using real-time quantitative pcr and sybr green chemistry. J Clin Microbiol 39: 2835-2845.
45. Pham LN, Dionne MS, Shirasu-Hiza M, Schneider DS (2007) A specific primed immune response in *Drosophila* is dependent on phagocytes. PLoS Pathogens 3: e26.
46. Schneider DS (2007) How and why does a fly turn it's immune system off? PLoS Biology 5: e247.
47. Little TJ, Kraaijeveld AR (2004) Ecological and evolutionary implications of immunological priming in invertebrates. Trends in Ecology & Evolution 19: 58-60.
48. Namikoshi A, Wu JL, Yamashita T, Nishizawa T, Nishioka T, et al. (2004) Vaccination trials with *Penaeus japonicus* to induce resistance to white spot syndrome virus. Aquaculture 229: 25-35.
49. Kurtz J, Franz K (2003) Innate defence: evidence for memory in invertebrate immunity. Nature 425 : 37-38.
50. Carius HJ, Little TJ, Ebert D (2001) Genetic variation in a host-parasite association: potential for coevolution and frequency-dependent selection. Evolution 55: 1136-1145.
51. Chong YC, Loh H (1984) Hepatopancreas chlamydial and parvoviral infections of farmed marine prawns in Singapore. Singap Vet J 9: 51-56.
52. Nimitphak T, Kiatpathomchai W, Flegel TW (2008) Shrimp hepatopancreatic parvovirus detection by combining loop-mediated isothermal amplification with a lateral flow dipstick. J Virol Methods 154: 56-60.
53. La Fauce KA, Elliman J, Owens L (2007) Molecular characterisation of hepatopancreatic parvovirus (PmergDNV) from Australian *Penaeus merguensis*. Virology 362: 397-403.
54. Sukhumsirichart W, Attasart P, Boonsaeng V, Panyim S (2006) Complete nucleotide sequence and genomic organization of hepatopancreatic parvovirus (HPV) of *Penaeus monodon*. Virology 346: 266 - 277.
55. Lightner DV (1993) Diseases of Cultured Penaeid Shrimp. CRC Press, Boca Raton, FL.
56. Sukhumsirichart W, Wongteerasupaya C, Boonsaeng V, Panyim S, Sriurairatana S, et al. (1999) Characterization and PCR detection of hepatopancreatic parvovirus (HPV) from *Penaeus monodon* in Thailand. Dis Aquat Org 38: 1-10.

57. Bonami JR, Mari J, Poulos BT, Lightner DV (1995) Characterization of hepatopancreatic parvo-like virus, a second unusual parvovirus pathogenic for penaeid shrimps. *J Gen Virol* 76: 813-817.
58. Bonami JR, Lightner DV (1991) Unclassified viruses of Crustacea. In: Adams, J.R., Bonami, J.R. (Eds.), *Atlas of Invertebrate Viruses*. CRC Press, Boca Raton, FL, 597-622.
59. Francki RIB, Fauquet CM, Knudson DL, Brown F (1991) Classification and nomenclature of viruses. Fifth report of the international committee on taxonomy of viruses. *Arch Virol Suppl* 2: 1-450.
60. Lightner DV, Redman RM (1985) A parvo-like virus disease of penaeid shrimp. *J Invert Pathol* 45: 47-53.
61. Cotmore SF, Tattersall P (1996) DNA Replication in Eukaryotic cells "Chapter 28: Parvovirus DNA Replication" <http://www.hixonparvo.info/Peter%20Parvo%20DNA%20replication%20Review%201996.pdf>).
62. Safeena MP, Tyagi A, Rai P, Karunasagar I, Karunasagar I (2010) Complete nucleic acid sequence of *Penaeus monodon* densovirus (PmDENV) from India. *Virus Research* 150: 1-11
63. Rukpratanpom S, Menasveta P, Sukhumsirichart W, Chaivisuthangkura P, Longyant S, et al. (2005) Generation of monoclonal antibodies specific to hepatopancreatic parvovirus (HPV) from *Penaeus monodon*. *Dis Aquat Org* 65: 85-89.
64. Sukhumsirichart W, Kiatpathomchai W, Wongteerasupaya C, Withyachumnarnkul B, Flegel TW, et al. (2002) Detection of hepatopancreatic parvovirus (HPV) infection in *Penaeus monodon* using PCR-ELISA. *Mol Cell Probes* 16: 409-413.
65. Lightner DV, Redman RM, Bell TA (1983 a) Detection of IHNV in *Penaeus stylirostris* and *P. vannamei* imported into Hawaii. *J World Maric Soc* 14: 212-225.
66. Lightner DV, Redman RM, Bell TA, (1983 b) Infectious hypodermal and hematopoietic necrosis, a newly recognized virus disease of penaeid shrimp. *J Invertebr Pathol* 42: 62-70.
67. Lightner DV, (1996a) Epizootiology, distribution and the impact on international trade of two penaeid shrimp viruses in the Americas. *Rev Sci Tech* 15: 579-601.
68. Bonami JR, Trumper B, Mari J, Brehelin M, Lightner DV (1990) Purification and characterization of the infectious hypodermal and hematopoietic necrosis virus of penaeid shrimps. *J Gen Virol* 71: 2657-2664.
69. Bell TA, Lightner DV (1987) IHNV disease of *Penaeus stylirostris*: effects of shrimp size on disease expression. *J Fish Dis* 10: 165-170.
70. Bell TA, Lightner DV (1984) IHNV: infectivity and pathogenicity studies in *Penaeus stylirostris* and *Penaeus vannamei*. *Aquaculture* 38: 185-194.
71. Brock JA, Main K (1994) *A Guide to the Common Problems and Diseases of Cultured Penaeus vannamei*. World Aquaculture Society, Baton Rouge, LA.
72. Kalagayan G, Godin D, Kanna R, Hagino G, Sweeney J (1991) IHNV as an etiological factor in runt deformity syndrome of juvenile *Penaeus vannamei* cultured in Hawaii. *J World Aquac Soc* 22: 235-243.
73. OIE (2003) *Manual of Diagnostic Tests for Aquatic Animals*. www.oie.int, World Animal Health Organization, Office International des Epizootics, Paris.
74. OIE (2000) *Diagnostic Manual for Aquatic Animal Diseases*. www.oie.int, World Animal Health Organization, Office International des Epizootics, Paris.
75. Brock JA, Gose RB, Lightner DV, Hasson KW (1997) Recent developments and an overview of Taura Syndrome of farmed shrimp in the Americas. In: Flegel TW, MacRae IH (eds.), *Diseases in Asian Aquaculture*, vol. III. Fish Health Section, Asian Fisheries Society, Manila, Philippines, 267-283.
76. Brock JA, Gose R, Lightner DV, Hasson KW (1995), LA, 84-94.
77. Hasson KW, Lightner DV, Poulos BT, Redman RM, White B L (1995) Taura Syndrome in *Penaeus vannamei*: demonstration of a viral etiology. *Dis Aquat Org* 23: 115-126.
78. Nielsen L, Sang-oum W, Cheevadhanarak S, Flegel TW (2005) Taura syndrome virus (TSV) in Thailand and its relationship to TSV in China and the Americas. *Dis Aquat Org* 63: 101-106.
79. Tu C, Huang HT, Chuang SH, Hsu JP, Kuo ST, et al. (1999) Taura syndrome in Pacific white shrimp *Penaeus vannamei* cultured in Taiwan. *Dis Aquat Org* 38: 159-161.
80. Bonami JR, Hasson KW, Mari J, Poulos BT, Lightner DV (1997) Taura syndrome of marine penaeid shrimp: characterization of the viral agent. *J Gen Virol* 78: 313-319.
81. Mayo MA (2005) Changes to virus taxonomy. *Arch Virol* 150: 189-198.
82. Srisuvan T, Tang KFJ, Lightner DV (2005) Experimental infection of *Penaeus monodon* with Taura syndrome virus (TSV). *Dis Aquat Organ* 67: 1-8.
83. Chang YS, Peng SE, Yu HT, Liu FC, Wang CH, et al. (2004) Genetic and phenotypic variations of isolates of shrimp Taura syndrome virus found in *Penaeus monodon* and *Metapenaeus ensis* in Taiwan. *J Gen Virol* 85: 2963-2968.
84. Cowley JA, McCulloch RJ, Spann KM, Cadogan LC, Walker PJ (2005a) Preliminary molecular and biological characterization of Mourilyan virus (MoV): a new bunyavirus-related virus of penaeid prawns. In: Walker PJ, Lester R, Bondad-Reantaso MB (Eds) *Diseases in Asian Aquaculture*, vol. V. Fish Health Section, Asian Fisheries Society, Manila, 113-124.
85. Sellars MJ, Keys SJ, Cowley JA, McCulloch RJ, Preston NP (2006) Association of Mourilyan virus with mortalities in farm pond-reared *Penaeus (Marsupenaeus) japonicus* transferred to maturation tank systems. *Aquaculture* 252: 242-247.
86. Cowley JA, McCulloch RJ, Rajendran KV, Cadogan LC, Spann KM, et al. (2005b) RT-nested PCR detection of Mourilyan virus in Australian *Penaeus monodon* and its tissue distribution in healthy and moribund prawns. *Dis Aquat Org* 66: 91-104.
87. Momoyama K, Muroga K (2005) Diseases of cultured kuruma shrimp in Japan: a review. *Fish Pathol* 40: 1-14.
88. Chayaburakul K, Nash G, Pratanpipat P, Sriuirairatana S, Withyachumnarnkul B (2004) Multiple pathogens found in growth-retarded black tiger shrimp *Penaeus monodon* cultivated in Thailand. *Dis Aquat Org* 60: 89-96.
89. NACA/FAO (2006) *Quarterly Aquatic Animal Disease Report (Asia and Pacific Region)*, Bangkok.
90. Senapin S, Phewsaiya K, Briggs M, Flegel TW (2007) Outbreaks of infectious myonecrosis virus (IMNV) in Indonesia confirmed by genome sequencing and use of an alternative RT-PCR detection method. *Aquaculture* 266: 32-38.
91. Tang KFJ, Pantoja CR, Poulos BT, Redman RM, Lightner DV (2005) In situ hybridization demonstrates that *Litopenaeus vannamei*, *L. stylirostris* and *Penaeus monodon* are susceptible to experimental infection with infectious myonecrosis virus (IMNV). *Dis Aquat Organ* 63: 261-265.
92. Underwood DJ, Cowley JA, Sellars MJ, Barnes AC, van Hulst MCW, et al. (2010) Gill-associated virus and recombinant protein vaccination in *Penaeus monodon*. *Aquaculture* 308: 82-88.
93. Callinan RB, Jiang L (2003) Fatal virus-associated peripheral neuropathy and retinopathy in farmed *Penaeus monodon* in eastern Australia. II. Outbreak descriptions. *Diseases of Aquatic Organisms* 53: 195-202.
94. Cowley JA, Hall MR, Cadogan LC, Spann KM, Walker PJ (2002) Vertical transmission of gill-associated virus (GAV) in the black tiger prawn *Penaeus monodon*. *Diseases of Aquatic Organisms* 50: 95-104.
95. Spann KM, Donaldson AR., Cowley JA, Walker PJ (2000) Differences in the susceptibility of some penaeid prawn species to gill-associated virus (GAV) infection. *Diseases of Aquatic Organisms* 42: 221-225.
96. Ravi M, Nazeer Basha A, Sarathi M, Rosa Idalia HH, Widada J S, et al. (2009) Studies on the occurrence of white tail disease (WTD) caused by MrNV and XSV in hatchery-reared post-larvae of *Penaeus indicus* and *P. monodon*. *Aquaculture* 292: 117-120.
97. Hameed ASS, Yoganandhan K, Sri Widada J, Bonami JR (2004a) Studies on the occurrence and RT-PCR detection of *Macrobrachium rosenbergii* nodavirus and extra small virus-like particles associated with white tail disease of *Macrobrachium rosenbergii* in India. *Aquaculture* 238: 127-133.
98. Qian D, Shi Z, Zhang S, Cao Z, Liu W, et al. (2003) Extra small virus-like particles (XSV) and nodavirus associated with whitish muscle disease in the giant freshwater prawn, *Macrobrachium rosenbergii*. *J Fish Dis* 26: 521-527.
99. Romestand B, Bonami JR (2003) A sandwich enzyme linked immunosorbent assay (S-ELISA) for detection of MrNV in the giant freshwater prawn, *Macrobrachium rosenbergii* (de Man). *J Fish Dis* 26: 71-75.
100. Sri Widada J, Bonami JR (2004) Characteristics of the monocistronic genome of extra small virus, a virus-like particle associated with *Macrobrachium*

-
- rosenbergii nodavirus: possible candidate for a new species of satellite virus. J. Gen. Virol 85: 643-646.
101. Arcier JM, Herman F, Lightner DV, Redman R, Mari J, et.al (1999) A viral disease associated with mortalities in hatchery-reared postlarvae of the giant freshwater prawn *Macrobrachium rosenbergii*. Dis Aquat Org 38: 177-181.
102. Yoganandhan K, Leartvibhas M, Sriwongpuk S, Limsuwan C (2006) White tail disease of the giant freshwater prawn *Macrobrachium rosenbergii* in Thailand. Dis Aquat Org 69: 255-258.
103. Wang CS, Chang JS, Wen CM, Shih HH, Chen SN (2008) *Macrobrachium rosenbergii* noda virus infection in *M. rosenbergii* (de Man) with white tail disease cultured in Taiwan. J Fish Dis 31: 415-422.
104. Hameed ASS, Yoganandhan K, SriWidada J, Bonami JR (2004b) Experimental transmission and tissue tropism of *Macrobrachium rosenbergii* nodavirus (MrNV) and extra small virus like-particles in *Macrobrachium rosenbergii*. Dis Aquat Org 62: 191-196.