

# Timely Initiation of Complementary Feeding Practice Among Mothers and Care Givers of Children Age 6 to 24 Months in Goba Town, Southeast Ethiopia

Yewalashet Alemayehu<sup>1</sup>, Tilahun Ermeko<sup>1\*</sup>, Abduljewad Hussien<sup>1</sup>, Abate Lette<sup>1</sup>, Teshome Gudisa<sup>1</sup>, Neway Assaye<sup>1</sup>, Tesfaye Eshetu<sup>1</sup>, Habtamu Gezahgn<sup>2</sup>, Musa Kumbi<sup>1</sup>, Ahmed Yasin<sup>1</sup>, Adem Abdulkadir<sup>1</sup>

<sup>1</sup>Department of Public Health, College of Medicine and Health Science, Madda Walabu University, Bale Goba, Ethiopia; <sup>2</sup>Department of Physiology, College of Medicine and Health Science, Madda Walabu University, Bale Goba, Ethiopia

## ABSTRACT

**Background:** Complementary feeding is the introduction of solid or semi-solid foods at six months of age with small amounts and increasing the quantity as the child gets older while maintaining frequent breastfeeding. The target range for complementary feeding is generally taking to be 6-24 months while breastfeeding continues for up to two years of age or beyond. Inappropriate initiation of complementary feeding practices is a major obstacle in attaining and maintaining child health.

**Objective:** To assess the prevalence of timely initiation of complementary feeding practice among mothers of children aged from 6 to 24 months living in Goba town, Bale zone, southeast Ethiopia from April 20 to June 20, 2019.

**Methodology:** Community-based cross-sectional study was conducted among 346 mothers or caregivers who have children 6-24 months of age living in 03 kebele, Goba town, selected by simple random sampling technique. Data were collected by a trained interviewer using semi-structured questionnaires filled for respondents after data collection, all collected data were analyzed by SPSS software version 20.

**Result:** A total of 346 mothers who had children 6 to 24 months of age were responded to the questionnaire. Ethnic composition shows that Oromo 175 (50.6%), Amharic accounts for, 151 (43.6%), Others 12 (3.5%) and Tigre 8 (2.3%). Regarding religion Orthodox 237 (68.5%), Muslim 89 (25.7), and protestant religion 20 (5.8%). Among the respondent's occupational status, 230 (66.5%) were housewives, 59 (17.1%) merchants, 47 (13.6%) governmental workers, and students are the least dominant. Marital status mother and caregiver 340 (98.3%) married and 4 (1.2%) and 2 (0.6%) of mother and caregiver were single and divorced respectively. About 292 (84.4%) mothers had started complementary feeding at the right time, that is 6 months of age, while 35 (10.1%) of a mother had started in the age of before six months, and the rest 19 (5.5%) mothers started by the age of after six months.

**Conclusion:** In this study, initiation of complementary feeding at the recommended time of six months was seen in the majority of children. Inadequate let down of breast milk, due to medical illness, and breast problem were some of the reasons mentioned by mothers who initiated early whereas lack of knowledge was they started complementary feeding, immature stomach, and the advice of family members were the reasons for late initiation of CF.

**Keywords:** Timely initiation; Complementary feeding practice; Mothers and caregivers; Children age 6 to 24 months

## BACKGROUND

Complementary feeding is the introduction of solid or semi-solid foods at six months of age with small amounts and increasing

the quantity as the child gets older while maintaining frequent breastfeeding. The target range for complementary feeding is generally taking to be 6-23 months while breastfeeding continues for up to two years of age or beyond [1]. Timely initiation of appropriate

**Correspondence to:** Tilahun Ermeko, College of Medicine and Health Science, Department of Public Health, Goba Referral Hospital, Madda Walabu University, Bale Goba, Ethiopia; Phone: +251909648032; E-mail: tilahunjimma2008@gmail.com

**Received:** June 24, 2021; **Accepted:** July 10, 2021; **Published:** July 30, 2021

**Citation:** Alemayehu Y, Ermeko T, Hussien A, Lette A, Gudisa T, Abdi H, et al. (2021) Timely Initiation of Complementary Feeding Practice Among Mothers and Care Givers of Children Age 6 to 24 Months in Goba Town, Southeast Ethiopia. *Gynecol Obstet (Sunnyvale)* 11:564.

**Copyright:** ©2021 Alemayehu Y, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

nutrient intake, in quantity, bioavailability, and timing in infancy is essential for optimal growth and development. Complementary food is food other than breast milk, infant formula, or follow-on formula was given to infants and these can be liquids, semi-liquids, and solids [1,2].

Appropriate complementary feeding depends on accurate information and skilled support from the family, community, and health care system. Inadequate knowledge about appropriate foods and feeding practices is often a greater determinant of malnutrition than the lack of food. Moreover, diversified approaches are requiring ensuring access to foods that will adequately meet the energy and nutrient needs of growing children. Appropriate complementary feeding is measured entails; introduction of complementary foods at 6 months with continued breastfeeding up to at least 2 years and beyond, correct feeding frequency for age, and consumption of a diverse diet [2].

In addition, low-cost complementary foods, prepared with locally available ingredients using suitable small-scale production technologies in community settings, can help to meet the nutritional needs of older infants and young children [3-25]

The global rate of exclusive breastfeeding remains low and is not improving-indeed in some of the largest countries IYCF results are deteriorating. high stunting rates persist in many countries, with feeding and maternal nutrition receiving little attention many countries do not implement effective, comprehensive, and large-scale IYCF programmers and only allocate a very small proportion of available nutrition resources [23].

World health organization recommends exclusive breastfeeding and delaying the introduction of solid foods to an infant's diet until 6 months post-partum [22]. This practice will displace breastfeeding and place the baby to develop diseases and finally malnutrition. However, in many countries, this recommendation this followed by few mothers, and earlier complementary feeding onto solid is a commonly reported global practice a prospective, observational study in Ireland revealed that among 401 pregnant women only one mother (0.2%) complied with the WHO recommendation to exclusive breastfeeding up to 6 months. ninety-one (22.6%) infants were prematurely complemented onto solid at 12 weeks [24].

Inadequate complementary feeding practices result in undernutrition which suffers millions of children globally. Inappropriate initiation of complementary feeding practices is a major obstacle in attaining and maintaining child health. It is a major cause of malnutrition which directly or indirectly results in 60% of the 10.9 million deaths annually among children under five Well over two-thirds of these deaths, which are often associated with inappropriate feeding practices, occur during the first year of life. No more than 35% of infants worldwide are exclusively breastfed during the first four months of life; complementary feeding frequently begins too early or too late, and foods are often nutritionally inadequate and unsafe [4].

In developing countries total of 152 million, 101 million, and 51 million children under the age of five are estimated to be stunted, underweight, and wasted, respectively which is resulted from late initiation complementary feeding practices. Over one-third of under-five mortality is caused by malnutrition related to inadequate complementary feeding [2].

In general, inappropriate and poor initiation of complementary feeding practices means that many children continue to be

vulnerable to irreversible outcomes of stunting, poor cognitive development, and significantly increased risk of infectious diseases leading to gastroenteritis, diarrhea, and acute respiratory infection [5].

Infant formula supplementation at any age is uncommon in Ethiopia. Among breastfeeding Children under age two, very few (2 percent) consume infant formula. However, a much higher proportion (18%) receives other milk or other liquids. The introduction of other liquids, such as water, juice, and formula, takes place earlier than the recommended introduction at age six months [21].

Inappropriate initiation of complementary feeding practice is a major cause of malnutrition in Ethiopia. Nationally, 10% of Ethiopian children are wasted, and 3% are severely wasted and 44% of children under age five are stunted, and 21% of children are severely stunted.

## METHODS

### Study area, design, and period

A community-based cross-sectional study was conducted from April 20, 2019-June20, 2019.

The study was conducted in Goba town, Bale zone, Oromia regional state, southeast Ethiopia. Goba town is located 445 km from Addis Ababa this town has a latitude and longitude of 7°0'N 39°59'E and an elevation of 2,743 meters above sea level. The total population is estimated to be over 32,000 in the area of the city (Ethiopia, 13 November 2011) among this population 15182 are male, 16843 female and 4710 children were under five. Regarding religion distribution orthodox the dominant religion which accounts 68.84% and Muslim and protestant accounts 23.12% and 5.4% respectively. The two largest ethnic groups reported were the Oromo 63.13%, and Amhara 33.3%; all other ethnic groups made up 3.5% of the population. Goba town has five kebele among this 03 was selected for this study and all children age 6-24 months whose randomly selected were included.

### Sample size determination

The sample size for the study was calculated using single population proportion formula by Considering  $p=71.2\%$  prevalence of timely initiation of complementary feeding in a study done in Wolaita Sodo town, southern Ethiopia and 95% confidence level, Margin of error, assumed to be 5%. By considering a 10% non-response rate the final sample size was 346.

### Sampling technique and procedure

A simple random and systematic sampling technique was employed for this study. The primary sampling unit was selected from all Goba towns, then the town divided 5 kebele among five kebele 03 kebele was selected by simple random sampling technique and 346 mothers were selected by systematic sampling technique. Therefore all mothers of children aged 6-24 months in selected kebele were included in this study.

### Data collection tools

Data were collected by semi-structured questionnaire is from literature, early initiation of complementary feeding, and associated factors among infants and young children, which is used

to assess the timely initiation of complementary feeding and associated factors. The questionnaire was arranged and modified to address based on the objectives. Therefore, the questionnaire was divided into two parts: the first part addressing the socio-demographic characteristics of the mother or guardian and the infant followed by the second part which covers maternal reproductive history, health service utilization, and complementary feeding. The questionnaire originally is prepared in English then translated to the local language orally to "Afan Oromo and Amharic" during an interview with the help of language translators in order to create a better understanding for the respondents.

### Data quality control

The principal investigators that mean who collect data and undergo quality assessment had ongoing supervision each day during data collection to ensure the quality of data by checking filled format for their completeness and consistency. The data collectors (investigators) were trained for one day by the supervisor before they are assigned to collect the data. The data was checked daily for its completeness and those with more than one incomplete data were discarded.

Before actual data collection, the questionnaire was pre-tested for validity and reliability on 5% mothers; thereby possible adjustment or modification was made accordingly. The pretest was done in Hasasa town among mothers having children aged 6-24 months.

### Data processing and analysis

After ensuring the completeness of each questionnaires data was

entered into the computer and analyzed, processed using and SPSS software version 20 for window package, and the descriptive statistic was used, then finally summarized and compiled in frequencies, percent, texts, tables, charts, and graphs.

### Operational definitions

**Early initiation of complementary feeding:** It is the initiation of additional food other than breast milk for a young child before six months of age.

**Family planning utilization:** Use of birth control for child spacing for better growth of the child.

**Kebele:** Is the smallest administrative unit of Ethiopia.

**Late initiation of complementary feeding:** This is the introduction of solid and semisolid foods after completed 6 months (180 days) of age according to WHO guidelines [1].

**Mother's knowledge on complementary feeding:** Assessing the mothers whether they know or not know the exact time when complementary feeding is started.

**Post-natal care:** Care given to mother from delivery to 42 days.

**Timely initiation of complementary Feeding:** It is the initiation of additional food other than breast milk for a young child at sixth months of age along with continued breastfeeding.

### ETHICAL CONSIDERATION

Before the actual data collection, and ethical clearance letter

**Table 1:** Socio-demographic characteristic and feeding practice of study participants.

Variable	Category	Frequency	Percent
Maternal age	15-24	103	29.7
	25-34	176	49.9
	Above 35	67	19.4
Educational status	Not read and write	51	14.7
	Read and write	30	8.7
	Primary	151	43.6
	Secondary	72	20.9
Ethnicity	Above secondary	42	12.1
	Oromo	175	50.6
	Amhara	151	43.6
	Tigre	8	2.3
Religion	Others	12	3.5
	Orthodox	237	68.5
	Muslim	89	25.7
	Protestant	20	5.8
Marital status	Married	340	98.3
	Single	4	1.2
	Divorced	2	0.6
Economic status	Below 500 ETB	59	17.1
	500-1500 ETB	133	38.4
	1501-2000 ETB	132	38.2
	More than 2000	22	6.4
Age during marriage	15-19	209	60.4
	20-24	95	27.5
	25-29	28	8.1
	Above 29	14	4.1
Occupational status	House wife	230	66.5
	Governmental workers	47	13.6
	Merchants	59	17.1
	Students	5	1.4

Table 2: Child characteristics and feeding patterns in Goba town, south Eastern Ethiopia.

Variable	Category	Frequency	Percentage
Age of child	6-10 months	167	48.3
	11-15 months	63	18.2
	16-20 months	14	4.0
	21-24 months	102	29.5
Sex of child	Male	172	49.7
	Female	174	50.3
Age category for time to initiate complementary feeding	at 6 months	292	84.4
	<6 months	35	10.1
	>6 Months	19	5.5
Time to initiate complementary	Timely	292	84.4
Reasons to initiate complementary feeding early	my breast milk is not sufficient	18	51
	due to breast problem	5	14
	due to medical illness	6	17.5
	other mothers practice it	3	8.75
	this is our culture	3	8.75
	Reasons to initiate complementary feeding lately	Breast milk is enough	8
Didn't know when to start		6	35.7
The advice of family members and another mother		2	11.4
The child may not able to digest it		3	12.9

was obtained from the student research program of Madda Walabu University College of medicine and health sciences. The respondents were informed about the purpose of the study, and their oral consent was obtained. The respondents' right to refuse or withdraw from participating in the interview was fully maintained and the information collected was not being described in relation to individuals' names. Moreover, a letter of permission was being issued from Madda Walabu University Goba referral hospital school of Health Sciences research & community service directorate, before the actual data collection.

## RESULTS

### Socio-demographic characteristics

A total of 346 mothers who had children 6 to 24 months of age were responded to the questionnaire generating a response rate of 100%. Ethnic composition shows that Oromo and Amhara account for 175 (50.6%), and 151 (43.6%) respectively. with the least dominant ethnic group was Tigre with the frequency of 8 (2.3%) respectively. Regarding religion, most of them were orthodox religion followers with a frequency of 237 (68.5%), and the least dominant religion is protestant religion with a frequency of 20 (5.8%). Regarding the educational status of the respondent, most of them about 151 (43.6%) are a primary school, 72 (20.8%) secondary, 42 (12.1%) above secondary, and 51 (14.7%) were not read and write. Among the entire respondent around 230 (66.5%) they were housewife which is the most dominant, and around 59 (17.1%) of the respondent are merchants, 47 (13.6%) of respondents are government workers and students are the least dominant. Around 340 (98.3%) that is the majority of the respondent are married and 4 (1.2%) and 2 (0.6%) of mothers and caregivers are single and divorced respectively (Table 1).

### Child characteristics and feeding patterns

Children 6-24 months were included in the study. 167 (48.3%) were 6-10months and 63 (18.2%) children were 11-15 months of

age. From this age group, 172 (49.7%) and 174 (50.3%) were male and female respectively. About 292 (84.4%) mothers had started complementary feeding (Figures 1-3 )at the right time, that is 6 months of age, while 35 (10.1%) of a mother had started in the age of before six months, and the rest 19 (5.5%) mothers started by the age of after six months. 18 (51%) mothers who started early believed that milk was not enough, 6 (17.5%) due to medical illness, 5 (14%) due to breast problem, other mothers practice it and cultural background accounts about 6 (17.5%). Among mothers started lately (35.7%) said didn't know when to start, (12.9%) stomach is not matured, (40%) breast milk is enough, (11.4%) advice of family and other mother practice it (Table 2).

### Maternal characteristics and medical history of children

Among the study participants interviewed, all 346 (100%) were biological mothers of the children. 58.2% of mothers had three to four antenatal care visits during the last pregnancy and 20.8% had 4 and more visits. From all interviewed mothers 171 (49.4%) mothers attend delivery at the hospital, health center 104 (30.1%), and 71 (20.5%) at home. The mean total family size was  $1.6387 \pm 0.74569$  (SD) and about 171 childbirth intervals were two years and 132 above two (Table 3).

## DISCUSSION

In this study, the prevalence of timely initiation of complementary feeding was 84.4% which is higher than the study done in India found that more than three fourth of mothers had started complementary feeding at 6 months [12]. In the present study, the median age of starting complementary feeding was six months. Whereas in a country like China 76% of mothers were introduced to complimentary food to their infant between 4 and 6 months of age and Timely complementary feeding was 41.6% [10]. Studies from the UK, Belgium, and Sweden highlight that a large proportion of infants are prematurely complemented at, 4 months of age [25]. However, in our study, only 10.1% prematurely started complementary feeding. The prevalence of timely initiation

Table 3: Maternal characteristics and medical history of children 6-24 months of age in Goba town, Eastern Ethiopia.

Variable	Category	Frequency	Percent
Maternal ANC follow up	One times	67	19.5
	Two times	201	58.2
	Three times	71	20.8
	Four and above	7	1.6
Place of delivery	Hospital	171	49.4
	Health center	104	30.1
	Home delivery	71	20.5
PNC follow up of mother	Yes	210	60.7
	No	136	39.3
Numbers of children in the house	0-3 children	177	51.2
	4-6 children	121	35
	7-10 children	44	12.7
	Above 10	4	1.2
Birth interval	<2 years	43	12.6
	2 year	171	49.1
	>2 year	32	38.4
Child immunization history	Yes	305	88.1
	No	39	11.9
Child immunization certificate	Yes	207	59.8
	No	139	40.2
Family planning utilization	Yes	246	71.1
	No	100	28.9




Figure 1: Time of introducing complementary feeding among child and caregiver of a child aged 6-24 months.




Figure 3: Type of food to start complementary feeding.




Figure 2: Source of knowledge on timely introducing of complementary feeding.

of complementary feeding in this study is much higher than the national prevalence of timely initiation of complementary feeding

(62.5) and also higher than the study done in Kamba Woreda Southwest Ethiopia revealed that among 392 respondents, 59.6% mothers start complementary feeding early before six months and 40.4% started complementary feeding at six months and relatively higher than the study done in northern Ethiopia, in mekele the prevalence of timely initiated complementary feeding at the age of six months was 62.8%. This relatively higher prevalence of timely initiation of complementary feeding at six months could be due to the communities' awareness towards the time of initiation of complementary feeding by different health information providers like health professional workers, HEW providing information which has contributed to high coverage of health institutional delivery and postnatal care. Media has also a great role in awareness creation to initiate complementary feeding on time in Goba town.

### CONCLUSION

In this study, initiation of complementary feeding at the recommended time of six months was seen in the majority of children. High institutional delivery accompanied with higher ANC follow-up contributes to this finding. The inadequate let down of breast milk, increased appetite to eat, and other information sources were some of the. Reasons mentioned by mothers who initiated early whereas immature stomach and the perception that breastfeeding is enough were the reasons for late initiation of CF.

## CONSENT FOR PUBLICATION

Not applicable.

## AVAILABILITY OF SUPPORTING DATA

Data will be available upon request.

## COMPETING INTERESTS

The authors have no competing interests and all have agreed on the manuscript for publication.

## FUNDING

This study has no specific funding. The authors have no competing interests and all have agreed on the manuscript for publication.

## ACKNOWLEDGMENTS

The authors are grateful to Madda Walabu University for supporting this study. We are also very grateful to data collectors for their cooperation to undertake this study.

## REFERENCES

- Joshi N, Agho KE, Dibley MJ, Senarath U, Tiwari K. Determinants of inappropriate complementary feeding practices in young children in Nepal: secondary data analysis of demographic and health survey 2006. *Matern Child Nutr.* 2012;8:45-59.
- Fabrizio CS, van Liere M, Peltó G. Identifying determinants of effective complementary feeding behaviour change interventions in developing countries. *Matern Child Nutr.* 2014;10:575-592.
- Chun-Ming CH, Yu-Ying WA, Chang SY. Effect of in-home fortification of complementary feeding on intellectual development of Chinese children. *Biomed Environ Sci.* 2010;23:83-91.
- Saleh F, Ara F, Hoque MA, Alam MS. Complementary feeding practices among mothers in selected slums of Dhaka city: A descriptive study. *J Health Popul Nutr.* 2014;32:89.
- Monte CM, Giugliani ER. Recommendations for the complementary feeding of the breastfed child. *J Pediatr.* 2004;80:s131-s41.
- Ethiopia Demographic and Health Survey. Central Statistical Agency Addis Ababa, Ethiopia. 2013.
- UNICEF/WHO/The World Bank Group joint child malnutrition estimates: Levels and trends in child malnutrition: key findings of the 2020 edition. 2020.
- WHO. Guiding Principles for Complementary Feeding of the Breastfed Child. World Health Organization. 2003.
- Garg A, Chadha R. Index for measuring the quality of complementary feeding practices in rural India. *J Health Popul Nutr.* 2009;27:763.
- Li L, Li S, Ali M, Ushijima H. Feeding practice of infants and their correlates in urban areas of Beijing, China. *Pediatr Int.* 2003;45:400-406.
- Ali SM. Poverty and Child Mortality in Pakistan. Pakistan Institute of Development Economics. 2008.
- Schiess S, Grote V, Scaglioni S, Luque V, Martin F, Stolarczyk A, et al. Introduction of complementary feeding in 5 European countries. *Journal of pediatric gastroenterology and nutrition.* 2010 1;50:92-98.
- Sapkota S, Shrestha S. Complementary feeding practices among the caretakers of the young children at Kathmandu. *J Chitwan Med Coll.* 2013;3:25-29.
- Kimani-Murage EW, Madise NJ, Fotso JC, Kyobutungi C, Mutua MK, Gitau TM, et al. Patterns and determinants of breastfeeding and complementary feeding practices in urban informal settlements, Nairobi Kenya. *BMC public health.* 2011;11:1-1.
- Semahegn A, Tesfaye G, Bogale A. Complementary feeding practice of mothers and associated factors in Hiwot Fana Specialized Hospital, Eastern Ethiopia. *The Pan African Medical Journal.* 2014;18.
- Tamiru D, Aragu D, Belachew T. Survey on the introduction of complementary foods to infants within the first six months and associated factors in rural communities of Jimma Arjo. *Int J Nutr Food Sci.* 2013;2:77-84.
- Rao S, Swathi P, Unnikrishnan B, Hegde A. Study of complementary feeding practices among mothers of children aged six months to two years-A study from coastal south India. *Australas Med J.* 2011; 4:252-257.
- Rebhan B, Kohlhuber M, Schwegler U, Koletzko BV, Fromme H. Infant feeding practices, and associated factors through the first 9 months of life in Bavaria, Germany. *J Pediatr Gastroenterol Nutr.* 2009;49:467-473.
- Inayati DA, Scherbaum V, Purwestri RC, Hormann E, Wirawan NN, Suryantan J, Hartono S, Bloem MA, Pangaribuan RV, Biesalski HK, Hoffmann V. Infant feeding practices among mildly wasted children: a retrospective study on Nias Island, Indonesia. *Int Breastfeed J.* 2012;7:1-9.
- Kaye S, Miles D, Antoine P, Burny W, Ojuola B, Kaye P, et al. Virological and immunological correlates of mother-to-child transmission of *Cytomegalovirus* in The Gambia. *J Infect Dis.* 2008;197:1307-1314.
- Mekbib E, Shumey A, Ferede S, Haile F. Magnitude and factors associated with appropriate complementary feeding among mothers having children 6-23 months-of-age in northern Ethiopia; a community-based cross-sectional study. *J Food Nutr Sci.* 2014;2:36-42.
- World Health Organization. 2011. Complementary feeding, report of the global consultation, Geneva. 2003.
- Global Nutrition Cluster. Strengthening Infant and Young Child Feeding Programming and Planning for Emergency Preparedness and Response. Proceedings of an international workshop. Save the Children UK in co-operation with UNICEF's IYCN and Emergencies Units and the IASC Global Nutrition Cluster. 1-50.
- Tarrant RC, Younger KM, Sheridan-Pereira M, White MJ, Kearney JM. Factors associated with weaning practices in term infants: a prospective observational study in Ireland. *British Journal of Nutrition.* 2010;104:1544-1554.
- Ethiopia FD. Ethiopia Demographic and Health Survey 2016. Addis Ababa Ethiopia, and Rockville. 2016.