

## Analysis of Human Androgen Receptor Polymorphism Using Fluorescent Loop-Hybrid Mobility Shift Technique

Tomokazu Ishikawa<sup>1</sup>, Shoichi Matsukuma<sup>2</sup>, Mitsuyo Yoshihara<sup>2</sup>, Tatsuo Kurosawa<sup>1</sup> and Yohei Miyagi<sup>2\*</sup>

<sup>1</sup>Diagnostics Research Laboratories, Wako Pure Chemical Industries, Ltd., 6-1 Takada-cho, Amagasaki 661-0963, Japan

<sup>2</sup>Molecular Pathology and Genetics Division, Kanagawa Cancer Center Research Institute, 2-3-2 Nakao, Asahi-ku, Yokohama 241-0815, Japan

### Abstract

A fluorescent loop-hybrid mobility shift (LH-MS) technique was introduced for the analysis of polymorphic sites in exon 1 of the human androgen receptor gene. (CAG)<sub>17-31</sub> or (CTG)<sub>17-31</sub> loops were assumed to protrude from the sense- or antisense strands of the PCR products following hybridization with reverse or forward LH probes, respectively. When an array of male DNA was analyzed using Cy5-labeled LH probes, a unique linear correlation was established between CAG repeat lengths and the fluorescent LH band positions on polyacrylamide gels. This linearly shifting band patterns were used to assemble LH ladder size markers of CAG repeat lengths. Analysis of female DNA revealed that 87% of females are heterozygous for CAG repeat polymorphisms, which could be informative for clonality analysis of tumors in female cancer patients. As a proof of principle, heterozygous female colorectal tumor DNA was examined with fluorescent LH-MS technique and loss of one allele after treatment with methylation-sensitive restriction enzyme *HpaII* was clearly exhibited.

**Keywords:** Loop-hybrid mobility shift; *HUMARA*; Genotyping; CAG repeats; Electrophoresis

### Introduction

Polymorphic CAG repeats in the human androgen receptor gene (*HUMARA*), located at Xq12, have been used as a highly informative genetic marker for human female tumor cells. Because of the random inactivation of one X chromosome, the heavily methylated state of one of the heterozygous alleles provides a useful tool, together with methylation-sensitive restriction enzymes, to examine clonality of tumor cells in female cancer patients [1-3]. In this respect, a simple method to determine the allelic status of CAG repeats using slab-gel electrophoresis would be of some value. Previously, we have developed LH-MS technology for detection of hot spot mutations at various oncogene loci for targeted therapy, such as *EGFR* codon 858, *BRAF* codon 600 and *KRAS* codons 12 and 13 [4,5]. The mutated alleles were differentiated from the wild type by means of mobility-shift of the loop-hybrids. Unique sequences in these genes produced simple PCR bands and the LH-bands for the mutant alleles were shifted from the wild type and detected unambiguously. On the other hand, when relatively short di-nucleotide repeat polymorphisms in *UGT1A1*, namely (TA)<sub>n</sub> vs. (TA)<sub>n-1</sub>, were genotyped using a loop-hybrid mobility shift (LH-MS) technique, PCR product of the segment containing these di-nucleotide repeats produced several confounding bands after polyacrylamide-gel electrophoresis (PAGE). Using a Cy5-labeled LH-probe, however, simple fluorescent LH bands were detected and they were uniquely associated to the variant alleles of di-nucleotide repeats [6]. In this study, we show that a fluorescent LH-MS technique facilitates the detection of far more complex genotypic variants of CAG repeats ranging from 17 to 31 in repeat numbers. The present technique is capable of detecting even a single repeat unit difference and reveals heterozygosity of this locus in 87% of human female study population.

### Materials and Method

#### DNA

Blood DNA samples from an anonymized healthy adult Japanese population (39 males and 46 females) obtained with informed consent [6] were used. Tumor DNA was previously obtained from frozen tumor tissue collected from 215 colorectal cancer patients with informed

consent [7], of which females were 91, and was used for clonality testing of the tumors. This study was approved by the Internal Review Board of the Kanagawa Cancer Center, Yokohama, Japan.

#### Restriction enzymes

*HpaII* (50000 U/μl, NEB, Ipswich, MA), a methylation-sensitive restriction enzyme, and *MspI* (20000 U/μl, NEB), a methylation-insensitive isoschizomer, both of which cleave CCGG, were used. Genomic DNA (50 ng/100 μl) was column-purified and concentrated (Zymo Research, Irvine, CA) to yield a purified DNA solution (8 μl). Purified DNA (6.2 μl) was treated with 1 μl of *HpaII* or *MspI* together with 0.8 μl of 10xbuffer for 24-48 h at 37°C. Two restriction sites were present in the amplicon used in this study. One microliter of the digestion mixture was used directly as template in a 10-μl PCR reaction mixture, and 0.8 μl of the eluted DNA was used as untreated control template DNA.

#### Primers and LH probes

Primer sequences and LH probe sequences (Fw- and Rv-types) are given in Table 1. LH probes were labeled with Cy5 at the 5'-end. The human androgen receptor gene, *HUMARA*, resides at Xq12. The genomic sequence used in this study was derived from nucleotides 67544032-67730619 of the human X chromosome sequence (GenBank accession number NC\_000023, [http://www.ncbi.nlm.nih.gov/nuccore/NC\\_000023.11](http://www.ncbi.nlm.nih.gov/nuccore/NC_000023.11)). The polymorphic CAG repeat is located in exon 1. An LH probe was designed to produce loop-hybrids such that the variable

**\*Corresponding author:** Yohei Miyagi, Molecular Pathology and Genetics Division, Kanagawa Cancer Center Research Institute, 2-3-2 Nakao, Asahi-ku, Yokohama 241-0815, Japan, Tel: 81455202222; Fax: 81455202216; E-mail: [miyagi@gancen.asahi.yokohama.jp](mailto:miyagi@gancen.asahi.yokohama.jp)

Received April 15, 2016; Accepted April 20, 2016; Published April 27, 2016

**Citation:** Ishikawa T, Matsukuma S, Yoshihara M, Kurosawa T, Miyagi Y (2016) Analysis of Human Androgen Receptor Polymorphism Using Fluorescent Loop-Hybrid Mobility Shift Technique. Adv Tech Biol Med 4: 174. doi: [10.4172/2379-1764.1000174](https://doi.org/10.4172/2379-1764.1000174)

**Copyright:** © 2016 Ishikawa T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Oligomer	N-mer	Sequence (5'-3')
Forward PCR primer	20	agcgtgctg aagtgatcca
Reverse PCR primer	20	atgggcttgg ggagaacct
Fw-type LH probe (LH-CTG)	99	agcgtgctg aagtgatcca gaaccgggc cccaggcacc cagaggccgc gagcgcagca cctcccggcg ccagttgct gctggagact agccccagg
Rv-type LH probe (LH-CAG)	64	atgggcttgg ggagaacct cctcacctt gctggggct agtctctgca gcagcaaac ggcg

**Table 1:** Nucleotide sequences of PCR primers and LH probes.

repeat of (CTG)<sub>n</sub> and six neighboring nucleotides would loop out from the antisense strand of the amplified polymerase-chain reaction (PCR) product following hybridization with Fw-type LH-CTG probe, or from the sense strand using Rv-type LH-CAG probe. The LH probes, which were shorter than the amplicon, would be filled by polymerase extension in the hybrid to yield the complete LH form. In Rv-type LH probe, both the polymorphic CAG repeats and the other stable repeat, (CAG)<sub>6</sub>, were deleted so that the LH generated by hybridization of the Rv-type LH probe with the sense-strand was assumed to produce two loops 15 bp apart.

### Fluorescent LH mobility shift technique

A previously described LH protocol [4] was modified as follows. PCR was performed using AccuPrime *Taq* polymerase (ThermoFisher, Waltham, MA) together with the primer pairs (Table 1) and the DNA template described above, under the following conditions: 94°C for 4 min, followed by 40 cycles of 94°C for 15 s, 55°C for 15 s, and 68°C for 45 s. LH probe (0.6 μl of 200 nM stock) was combined with 4.5 μl of PCR product to generate an LH by denaturation at 94°C for 4 min, followed by 55°C for 15 s and 68°C for 4 min. LH product (1.5 μl) was separated on a 10% pre-formed polyacrylamide gel (6 cm long, ATTO Co., Tokyo, Japan) by electrophoresis at 25 mA in Tris-glycine running buffer (37.5 mM Tris, 288 mM glycine) for 30 min. The gel was then stained with SYBR GreenI (ThermoFisher) for 8 min, and visualized using a laser scanner (STORM860, GE Healthcare, Little Chalfont, UK)

at 450 nm excitation together with Longpath (LP) filter 520 nm for SYBR GreenI, or at 635 nm/LP 650 nm for Cy5 fluorescence [6].


### Cloning and sequencing

To determine the actual repeat length, PCR products were cloned and sequenced. PCR products were ligated into vector pCR2.1 using a TOPO TA cloning kit (ThermoFisher) and then transformed into One Shot TOP10 electrocompetent *Escherichia coli* according to the manufacturer's instructions. Plasmid DNA was extracted from 20 colonies using CloneChecker (ThermoFisher) and the DNA was amplified using Phi29 DNA polymerase (TempliPhi, GE Healthcare) for sequencing using a capillary sequencer (3130 Genetic Analyzer, ThermoFisher). After the CAG repeat numbers had been determined, diluted plasmid DNA was used as a template and the segment containing CAG repeats was PCR-amplified with the specified primers (Table 1), and analyzed using the fluorescent LH-MS technique to assign these actual CAG repeat numbers to the relevant LH band positions.

## Results

### CAG length polymorphism in human male DNA

For simplicity, CAG repeat length polymorphism in *HUMARA* exon 1 was first analyzed in male DNA. PAGE analysis of the PCR products of the exon 1 region containing CAG repeats exhibited various extra bands besides the bands of the expected sizes, as usually observed for amplicons containing highly repeated sequences [8]. Fluorescent LH-MS technique can circumvent this difficulty of complex band patterns due to short repeats. After hybridization with the fluorescent LH-CTG probe (Fw-type), the polymorphic CTG repeats and three neighboring nucleotides (CAG, TTG) looped out from the antisense strand of the LH. The LH can be visualized specifically as a single Cy5-fluorescent band on the gel. Among 39 male DNA samples examined with the fluorescent LH-MS technique, an array of 15 male DNA was obtained, which showed their LH bands positioned in a consecutive order displaying a stepwise shift from one LH band to the other (Figure 1A). By cloning and sequencing PCR products from these male DNA,



CAG repeat numbers were determined. The CAG repeat number in the cloned plasmids was largely consistent but, in a small fraction, the repeat number smaller by one repeat unit were observed, probably owing to polymerase slippage during PCR [9]. Such slippage-derived shorter PCR products were considered to account for the weak-intensity bands associated with the LH bands of male DNAs (Figure 1B). These weak-intensity bands were regarded as irrelevant bands for *HUMARA* genotyping.

Each LH band of the arrayed male DNA samples were allocated with CAG repeat numbers from 17 to 31 and used as a standard size marker to estimate CAG repeat lengths in fluorescent LH-MS technique. In the standard array of LH bands, it was noted that the migration-shift of the LH bands from 2n to 2n+1 repeats was always larger than the shift from 2n-1 to 2n when Fw-type LH-CTG probe was used (Figures 1A and 1C). On the other hand, when Rv-type LH-CAG probe was used, the migration shift from 2n-1 to 2n was larger than the shift from 2n to 2n+1 (Figures 1B and 1C). It follows that, in genotyping female DNA with heterozygosity of one CAG repeat unit difference, the difference of 2n and 2n+1 can be determined by Fw-type LH-CTG probe, meanwhile, the difference of 2n-1 and 2n can be determined by Rv-type LH-CAG probe (Figure 1C). Standard LH size ladder markers were prepared for even numbered (18-30) repeats and odd numbered (17-31) repeats, separately, with either Fw-type LH-CTG probe or Rv-type LH-CAG probe.

### Genotyping *HUMARA* alleles in female DNA

In contrast to male, a large part of female samples revealed two LH bands, which indicated allelic difference of CAG repeats and were sized properly using the standard LH size ladder markers (Figure 2A). Heterozygous cases of one repeat unit difference were also rendered to reveal two distinct LH bands by either Fw-type LH-CTG or Rv-type LH-CAG probe depending on CAG repeat composition. Only when both of these probes produced a single LH band, the female DNA was considered to be homozygous (Figure 2B).


In a healthy female Japanese population, 87% (40/46) were heterozygous, of which 22% (9/40) showed an allelic difference of one repeat unit. From the allelic frequency distribution (Figure 2C) for the population examined, including males, the expected rate of homozygosity (10.8%) was close to the actual rate (13%) found in the 46 female DNA samples, indicating that nearly 90% of female cases would be informative.

### A clonality test of female tumor DNA

According to random inactivation of X chromosomes in female cells during early embryogenesis [10,11], methylated CpG sites near the CAG polymorphic site of *HUMARA* in the inactive X chromosome present in the incipient tumor cells may be maintained and passed to the replicating tumor cells in female patients. In contrast, infiltrating macrophages and lymphocytes may be randomly methylated at the same locus. Female tumor DNA with informative heterozygous CAG repeats in *HUMARA* was digested with the methylation-sensitive restriction enzyme *HpaII*, and examined for the undigested methylated allele, which would be amplified by PCR and analyzed for the undigested allele with fluorescent LH-MS technique. As shown in the representative female cases, one of the two LH bands was undigested by *HpaII*, indicating non-random methylation of these alleles and putative clonal origin of these tumor cells (Figures 3A and 3B).

In one male tumor DNA sample, a single LH band was lost following *HpaII* digestion, consistent with a single active unmethylated X chromosome. However, another male tumor DNA sample showed an LH band only partially digested following *HpaII* treatment (Figure 3B). A certain alteration at the restriction sites in a subpopulation of the tumor cells may be considered.

The female tumor DNA which exhibited *HpaII* digestion only in one allele was treated with the methylation-insensitive enzyme *MspI*. Unexpectedly, the same *HpaII*-insensitive allele remained also as *MspI*-insensitive (Figures 3B and 3C), though the LH band appeared less



**Figure 2:** Analysis of CAG repeat length polymorphism in human female DNA. **A.** Electropherogram of the LH for heterozygous female DNA formed by Cy5-labeled LH-CTG probe and detected by Cy5 signal. Heterozygous CAG repeat numbers are shown for each sample. Me and Mo are markers for even numbered (18-30) and odd numbered (17-31) CAG repeats. **B.** Electropherogram of LH for heterozygous female DNA with one repeat unit differences and homozygous female DNA. Either of LH-CTG or LH-CAG probes was able to detect the unit repeat difference distinctly but not in homozygous female samples. **C.** Frequency distribution of the alleles with varying CAG repeat numbers, ranging from 17-31, among 107 chromosomes. The expected probability of homozygosity calculated from the distribution (11%) was close to the observed value (13%).





## Acknowledgment

Patent pending PCT/JP201358211. This study was supported partly by Wako Pure Chemical Industries, Ltd. The sponsors played no role in study design, in the collection, analysis, or interpretation of data, in the writing of the report, or in the decision to submit the article for publication.

## References

1. Allen RC, Zoghbi HY, Moseley AB, Rosenblatt HM, Belmont JW (1992) Methylation of HpaII and HhaI sites near the polymorphic CAG repeat in the human androgen-receptor gene correlates with X chromosome inactivation. *Am J Hum Genet* 51: 1229-1239.
2. Mashal RD, Lester SC, Sklar J (1993) Clonal analysis by study of X chromosome inactivation in formalin-fixed paraffin-embedded tissue. *Cancer Res* 53: 4676-4679.
3. Kopp P, Jaggi R, Tobler A, Borisch B, Oestreich M, et al. (1997) Clonal X-inactivation analysis of human tumours using the human androgen receptor gene (HUMARA) polymorphism: a non-radioactive and semiquantitative strategy applicable to fresh and archival tissue. *Mol Cell Probes* 11: 217-228.
4. Matsukuma S, Yoshihara M, Kasai F, Kato A, Yoshida A, et al. (2006) Rapid and simple detection of hot spot point mutations of epidermal growth factor receptor, BRAF, and NRAS in cancers using the loop-hybrid mobility shift assay. *J Mol Diagn* 8: 504-512.
5. Matsukuma S, Yoshihara M, Suda T, Shiozawa M, Akaike M, et al. (2011) Differential detection of KRAS mutations in codons 12 and 13 with a modified loop-hybrid (LH) mobility shift assay using an insert-type LH-generator. *Clinica Chimica Acta* 412: 1874-1878.
6. Matsukuma S, Saito H, Yamada K, Okamoto N, Ishikawa T, et al. (2011) Simple and precise detection of UGT1A1 polymorphisms with a modified loop-hybrid mobility shift assay using Cy5-labeled loop probes. *Clin Chim Acta* 412: 1668-1672.
7. Suda T, Yoshihara M, Nakamura Y, Sekiguchi H, Godai TI, et al. (2011) Rare MDM4 gene amplification in colorectal cancer: The principle of a mutually exclusive relationship between MDM alteration and TP53 inactivation is not applicable. *Oncol Rep* 26: 49-54.
8. Bovo D, Rugge M, Shiao YH (1999) Origin of spurious multiple bands in the amplification of microsatellite sequences. *Mol Pathol* 52: 50-51.
9. Shinde D, Lai Y, Sun F, Arnheim N (2003) Taq DNA polymerase slippage mutation rates measured by PCR and quasi-likelihood analysis: (CA/GT)<sub>n</sub> and (A/T)<sub>n</sub> microsatellites. *Nucleic Acids Res* 31: 974-980.
10. Lyon MF (1989) X-chromosome inactivation as a system of gene dosage compensation to regulate gene expression. *Prog Nucleic Acid Res Mol Biol* 36: 119-130.
11. Garcia SB, Novelli M, Wright NA (2000) The clonal origin and clonal evolution of epithelial tumours. *Int J Exp Pathol* 81: 89-116.
12. Chi LM, Lam SL (2005) Structural roles of CTG repeats in slippage expansion during DNA replication. *Nucleic Acids Res* 33: 1604-1617.
13. Canceill D, Viguera E, Ehrlich SD (1999) Replication slippage of different DNA polymerases is inversely related to their strand displacement efficiency. *J Biol Chem* 274: 27481-27490.
14. Vilar E, Gruber SB (2010) Microsatellite instability in colorectal cancer—the stable evidence. *Nat Rev Clin Oncol* 7: 153-162.
15. Van der Ploeg LH, Groffen J, Flavell RA (1980) A novel type of secondary modification of two CCGG residues in the human gamma delta beta-globin gene locus. *Nucleic Acids Res* 8: 4563-4574.