

Development and Validation of a New Chromatographic Method for the Estimation of Vismodegib by RP-HPLC

Veera Shakar Pulusu^{1*} and Parasharamulu Kommarajula²

¹School of Chemical Sciences, Swami Ramanand Teerth Marathwada University, Nanded, Maharashtra, India

²University College of Pharmaceutical Sciences, Kakatiya University, Warangal, Telangana, India

Abstract

An analytical method using high performance liquid chromatography equipped with photodiode array detector (HPLC-PDA) was developed and validated for estimation of Vismodegib as per ICH and FDA guidelines. Mobile phase consisted 0.1% Orthophosphoric acid: Acetonitrile (50:50, v/v) pumped at flow rate of 1.0 mL/min, in isocratic mode and detector was set at 264 nm. The method was validated in terms of the precision, linearity, accuracy, degradation and robustness. The detector response of Vismodegib is directly proportional to concentration ranging from 0.012-0.120 mg/mL and the mean recovery was 99.6% (RSD=0.19%, n=9). In the intra- and inter assay, the percentage difference was found to be less than 2%. Robustness was proved performing variation in mobile phase, flow rate and column temperature. A forced degradation study of Vismodegib was conducted under the condition of acidic, basic, thermal, peroxide, photo and hydrolysis. Vismodegib was found to be degraded (water bath at 60°C for 2 hours) in peroxide stress. The result of the study showed that the proposed method is simple, fast, precise and accurate, which is useful for the routine determination of Vismodegib in bulk and dosage forms.

Keywords: Vismodegib; Method development; RP-HPLC; Validation; Forced degradation

Introduction

Vismodegib is used for the treatment of adults with metastatic basal cell carcinoma, or with locally advanced basal cell carcinoma that has recurred following surgery or who are not candidates for surgery, and who are not candidates for radiation [1]. Vismodegib is a crystalline free base with a pKa (pyridinium cation) of 3.8, appearing as a white powder. Its molecular weight is 421.30 g/mol and the partition coefficient (log P) is 2.7. Vismodegib exhibits pH dependent solubility and it has been classified as a Class 2 molecule under the Biopharmaceutics Classification System (BCS) [2]. Vismodegib is chemically known as 2-Chloro-N-(4-chloro-3-(pyridine-2-yl)phenyl)-4-(methylsulfonyl)benzamide and its molecular formula C₁₉H₁₄Cl₂N₂O₃S (Figure 1). The mechanism of action of Vismodegib binds to and inhibits smoothed, a transmembrane protein involved in Hedgehog signal transduction [3]. In this present work, a new sensitive and rugged RP-HPLC method was developed for the determination of Vismodegib in bulk, and this method was validated according to FDA and ICH guidelines.




Figure 1: Chemical structure of Vismodegib.

Materials and Methods

Chemicals and reagents

Vismodegib standard (Purity \geq 99.7 as is basis), Acetonitrile (HPLC grade), HPLC grade water (Millipore), 85% Orthophosphoric acid, Sodium hydroxide, Hydrochloric acid and Hydrogen peroxide were purchased from Merck.

Instrumentation

The instrument used in the study were electronic balance (Make: Mettler Toledo, Model: XP56), sonicator (Make: Elma, Model: S300H), hot air oven (Make: Serve well Instruments, Model: H02436), digital pH meter (Make: Mettler Toledo) and UV-Visible chamber (Make: Mack Equipment, Model: MK-2). HPLC (Waters, 2695 with PDA detector 2996) was monitored and integrated using empower software.

Chromatographic conditions

Chromatographic analysis was performed on the isocratic mode with a mobile phase consisting of 0.1% phosphoric acid and Acetonitrile in the proportion of 50:50 (v/v). Prior to use the mobile phase was filtered through 0.45 μ m membrane filter and degassed for 10 min. The analysis was carried out on Waters 2695 series HPLC system with empower software. The analytes were conducted on an analytical column C18, (150 \times 4.6) mm, 5 μ m Zodiac, with a detection wavelength of 264 nm. The operating temperature of the column was 30°C and run time was 6 minutes. The injection volume was 10 μ L, and the flow rate was maintained at 1.0 mL/min.

***Corresponding author:** Veera Shakar Pulusu, School of Chemical Sciences, Swami Ramanand Teerth Marathwada University, Latur Road, Nanded - 431606, Maharashtra, India, Tel: 7829274853; E-mail: veerashakar@gmail.com

Received May 12, 2019; Accepted May 29, 2019; Published June 03, 2019

Citation: Pulusu VS, Kommarajula P (2019) Development and Validation of a New Chromatographic Method for the Estimation of Vismodegib by RP-HPLC. J Chromatogr Sep Tech 10: 421.

Copyright: ©2019 Pulusu VS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Method development

Selection and preparation of mobile phase: In order to select a suitable mobile phase for the analysis of Vismodegib, various ratios of buffers and organic solvents were tried on the basis of pKa and some trials. The most suitable mobile phase was found to be 0.1% Orthophosphoric acid: Acetonitrile in the ratio of 50:50 while considering the system suitability parameters such as retention time, tailing factor, number of theoretical plates. The mobile phase was filtered through 0.45 µm membrane filter and degassed by sonication. The optimization of mobile phase buffers was captured in Table 1.

Standard preparation: The standard solution of Vismodegib prepared as 30 mg of standard was dissolved in 50 mL volumetric flask and diluted to volume with mobile phase (Stock concentration 0.6 mg/mL). Further 5.0 mL of stock was diluted to 50 mL with mobile phase and mixed. (working std. concentration 0.06 mg/mL). The

representative chromatogram of Vismodegib and peak purity were shown in Figure 2 for the proposed method.

Method validation: The developed method was validated as per FDA and ICH guidelines by evaluating Precision (Repeatability and Reproducibility) linearity, accuracy, degradation and robustness.

Specificity: The ICH guidance defines specificity is the ability to assess unequivocally the analyte in the presence of components which may be expected to be present” [4]. Forced degradation study was conducted by exposing drug substance sample to various stress conditions. Stressed samples were analyzed, active peak was checked for the retention time, peaks interference and purity.

Precision

Precision is defined as “the measure of how close the data values are to each other for a number of measurements under the same analytical conditions” [5]. In precision analysis, system precision,




Figure 2: Typical chromatogram of Vismodegib under optimised chromatographic conditions.




Figure 3: Typical overlay chromatogram of Vismodegib on Acidic, Basic, Oxidative, Thermal, Photolytic and Hydrolysis degradation.

method precision and intermediate precision have been carried out. The system precision was determined by analyzing standard solution in six replicates, % RSD of area counts of Vismodegib peak was calculated. In method precision, six preparations of 100% test concentration against standard solution were analyzed. Intermediate precision was performed by different analyst on different day using different column. Overall RSD for assay between the two precision sets of data was calculated.

Linearity

Linearity is defined as “the ability to obtain test results which are directly proportional to the concentration of analyte in the sample”. For the establishment of linearity, a minimum of 5 different concentrations are recommended [4]. From the standard stock solution, a series of solution were prepared at a concentration levels ranging 0.012 mg/mL to 0.120 mg/mL. The peak area response of solutions at all levels in triplicate were measured. The peak response verses concentration data was treated by linear regression analysis and the linearity of response for Vismodegib was determined by calculating correlation coefficient.

Accuracy

Accuracy is the measure of how close the experimental value is to the true value [6]. For the determination of accuracy, the standard addition method was applied. In this study, known amount of active substance was spiked in sample solvent at three different levels in triplicate. Accuracy has been performed at about 50%, 100% and 150% of sample target concentration. The samples were analyzed by the proposed method and the amount of Vismodegib recovery was calculated by this formula:

$$\% \text{ Recovery} = \frac{\text{mg found}}{\text{mg added}} \times 100$$

S. No	Mobile phase	Remarks
Trial-1	0.1% Orthophosphoric Acid: ACN	Symmetrical peak shape
Trial-2	10 mM Na ₂ HPO ₄ ·H ₂ O (pH 4.9): ACN	Good peak shape but salt used
Trial-3	10 mM Na ₂ HPO ₄ ·H ₂ O (pH 4.9): Methanol	Late elution
Trial-4	10 mM KH ₂ PO ₄ : Methanol	Broad peak and late elution
Trial-5	10 mM KH ₂ PO ₄ : Acetonitrile	Good peak shape but salt used
Trial-6	0.1% Orthophosphoric Acid: Methanol	Broad peak and late elution

Table 1: Optimization of mobile phase and impact on chromatography.

Robustness

Robustness of the method was investigated by varying the instrumental conditions such as flow rate (± 0.2 mL/min), mobile phase composition ($\pm 10\%$ absolute) and column temperature ($\pm 5^\circ\text{C}$). System suitability criteria of the standard solution was checked at each minor variable condition. The retention time (RT), USP tailing factor, theoretical plate counts and % RSD of area counts of Vismodegib from standard solution for each set of data was calculated.

Forced degradation

Force degradation or stress testing includes four main degradation mechanisms: thermal, acid/base hydrolysis, oxidative, and photolytic degradation. Selecting suitable reagents and length of exposure can achieve the preferred level of degradation. Over stressing a sample may lead to the formation of secondary degradants that would not be seen in formal shelf-life stability studies and under-stressing may not serve the purpose of stress testing [7]. Therefore, it is necessary to control the degradation to a desired level [7].

Thermal stress

In thermal stress, solid drug substances and drug products should be exposed to heat. It is recommended that the effect of temperature be studied in 10°C increment above that for routine accelerated testing, and humidity at 75% relative humidity or greater [8]. The heating time can be increased if there is no significant degradation observed in initial study. By increasing the temperature, the rate of reaction also tends to increase the production of degradation products. Thermal degradation was performed by treating the Vismodegib drug substance at $40^\circ\text{C}/75\%$ RH for 14 days in an open container. Sample was diluted as per required concentration with sample solvent and mixed. The obtained chromatogram was analyzed for any degradation occurred during the process. The results are given in Table 2.

Acid/base hydrolysis

In this stress study, the drug reacts with different pH conditions. In general, the drug substances are treated with different concentrations of Hydrochloric acid and Sodium hydroxide. If the reasonable degradation was not achieved, then higher concentration or longer duration time can be extended. After subjected to stress conditions,




Figure 4: Linearity curve of Vismodegib.

the samples should be neutralized with acid or base to avoid further degradation.

Acidic and basic degradations were performed using 0.1 M HCl and 0.1 M NaOH. Added 5.0 mL to each stock and refluxed at 60°C for 5 hours. After stressing sample stocks were neutralized with respective solutions and further diluted with sample solvent as per required concentration. The obtained chromatograms were analyzed for any degradation occurred during the process and the results are given in Table 2.

Oxidation stress

For oxidation stress, drug substances require free radical initiators for oxidation process. Oxidizing agents such as hydrogen peroxide, metal ions, oxygen and radical initiators can be used in oxidation stress. Different stress conditions may generate the same or different degradants [7]. The type and extent of degradation depends on the functional groups of the drug molecule and the stress conditions [7].

Peroxide degradation of Vismodegib was performed using 3% Hydrogen peroxide. Added 5.0 mL of 3% H₂O₂ and kept in water bath at 60°C for 2 hours. After attained room temperature, diluted to volume and further diluted as per required concentration. This solution was injected immediately to avoid excess degradation. The results are given in Table 2.

Photolytic degradation

In this study, the drug substances are exposed to light source. Some recommended conditions for photostability testing are described in ICH Q1B photostability Testing of New Drug Substances and Products [7]. Samples of drug substance, and solid/liquid drug product, should be exposed to a minimum of 1.2 million lux hours and 200-watt hours per square meter light. The samples should be exposed to both white and UV light. Temperature control may be necessary to minimize the effect of temperature changes during exposure [7]. The presence of the C=C, C=O, Aryl chloride, C₆H₄Cl₂, Nitroaromatic group, -C₆H₄NO₂, a weak C-H bond, Sulphides, alkanes, polyenes, and phenols chemical function groups in the drug molecules is usually necessary for the occurrence of photochemical reactions [9].

Vismodegib drug substance was transferred into a quartz container

and exposed to light about 1.2 × 10⁶ lux-hours. The exposed drug was diluted as per required concentration with sample solvent and analyzed. The results were captured in Table 2.

Hydrolysis degradation

Hydrolysis degradation of Vismodegib drug substance was performed using distilled water. Added 5.0 mL of distilled water and kept in water bath at 60°C for 1 hour. After attained room temperature, Sample was diluted as per required concentration with sample solvent and mixed. The results are given in Table 3 and Figure 3.

Results

Method development

Chromatographic separation: Method development was performed by screening organic modifiers (Acetonitrile and Methanol) and different strengths of buffers (Phosphate and Phosphoric acid). The final optimized mobile phase composition was found to be 0.1% phosphoric acid and Acetonitrile in the proportion of 50:50 (v/v). and run time was 6 min. The analysis was conducted at 264 nm with Zodiac C18 column (150 × 4.6) mm, 5 μm particle size and the column temperature was 30°C. The injection volume was 10 μL and the isocratic mobile phase flow rate was 1.0 mL/min.

Method validation

Linearity: The standard stock was used for linearity stock, serial dilutions were prepared and covered the range from 0.012 mg/mL to 0.120 mg/mL. Three replicate injections of each concentration were analyzed for this study and regression coefficient (r²) was found to be 0.9998 (Figure 4).

Precision: The HPLC is found to be precise at the % RSD of the area counts for six standard injections was 0.89. The method precision and intermediate precision samples were prepared at 100% of the sample target concentration and evaluated six samples as per method. The peak area was measured and % RSD was calculated. The developed method also precise as the % RSD of amount present for Vismodegib in the method precision sample was 0.22 and intermediate precision sample was 0.12. The details were given below Table 4.

Parameters	Modification	RT (min)	Area (AU)	USP Plate count	USP Tailing
Flow rate	0.8 mL/min	3.469	1886970	10855	1.07
	Optimized	2.989	2014527	10903	1.12
	1.2 mL/min	2.714	1780554	9710	1.08
Mobile phase composition	5% less	3.403	2074642	9449	1.05
	Optimized	2.989	2014527	10903	1.12
	5% more	2.783	1968163	10256	1.14
Column temperature	25°C	2.875	1916612	10064	1.11
	Optimized	2.989	2014527	10903	1.12
	35°C	3.227	2103336	9735	1.06

Note: RT (min), area counts, USP Plate count and USP Tailing values are in average.

Table 2: Robustness data at different variations.

S. No	Name of Degradation	RT (min)	Area (AU)	Purity Angle	Purity Threshold	% Degradation
1	Acid degradation	2.971	1897238	0.135	0.343	5.7
2	Base degradation	2.981	1931230	0.092	0.294	4.6
3	Peroxide degradation	3.046	1889376	0.111	0.294	6.5
4	Thermal degradation	3.029	1999243	0.168	0.303	0.7
5	UV degradation	3.028	2008919	0.121	0.295	0.3
6	Hydrolysis degradation	3.034	2012955	0.092	0.291	0.2

Table 3: Results of Vismodegib forced degradation.

System Precision		Method precision		Intermediate precision	
Inj/Sample No	Std. area (AU)	peak area (AU)	% Assay	peak area (AU)	% Assay
1	2028097	2025964	99.8	1983283	99.8
2	1993286	2057355	100.2	1994915	99.9
3	2018960	1999618	99.8	1987099	99.9
4	2011501	2034515	100.1	1999893	100.0
5	1996247	2010705	99.9	1979664	99.7
6	2039069	2013697	99.8	1982603	99.7
Average	2014527	--	99.9	--	99.9
S.D.	17891.9	--	0.219	--	0.121
% RSD	0.89	--	0.22	--	0.12

Table 4: System precision, method precision and intermediate precision.

S. No	Recovery level %	mg Added (spiked)	mg Found	% Recovery	Mean	SD	%RSD
1	50%-1	15.22	15.102	99.2	99.5	0.26458	0.27
2	50%-2	15.62	15.567	99.7			
3	50%-3	15.32	15.251	99.6			
4	100%-1	30.71	30.639	99.8	99.6	0.20817	0.21
5	100%-2	31.09	30.912	99.4			
6	100%-3	30.81	30.715	99.7			
7	150%-1	45.11	44.991	99.7	99.6	0.11547	0.12
8	150%-2	46.72	46.508	99.5			
9	150%-3	46.25	46.114	99.7			

Table 5: Recovery of Vismodegib drug substance.

Accuracy: The accuracy of the method was determined by analyzing samples at three different (50%, 100% and 150%) levels of concentrations by spiking Vismodegib active substance. The % recovery is found within the acceptable criteria, also individual and overall % RSD of recovery is found within the acceptable criteria; hence it shows that the method is accurate. The % recovery results are summarized in below Table 5.

Forced degradation studies: These stress conditions produce minute or no degradation due to the nature of a drug molecule. The nature of degradation depends on the functional groups of the drug molecule and the effect of stress conditions.

Robustness: Robustness of the method was studied by deliberate variation of the analytical parameters such as flow (1.0 ± 0.2 mL/min), column temperature (30 ± 5 °C) and mobile phase composition ($\pm 10\%$ absolute). The results are given in Table 4.

Discussion

No stability indicating method is available in the official compendia using HPLC for analyzing Vismodegib in bulk and formulation dosage forms until now. The present work has been reported with an intention to develop a new validated method for the estimation of Vismodegib by RP-HPLC method using a photo diode array detector. Different makes of C18 columns, buffers and organic solvents were used to develop a stability indicating method. To achieve symmetric peak shape, mobile phase prepared with 0.1% orthophosphoric acid (pH 2.50) and acetonitrile in the ratio of 1:1, column was selected Zodiac C18, (150 × 4.6) mm, 5 μm at a flow rate of 1.0 mL/min and column temperature at 30°C. Under the above optimized conditions, the retention time of Vismodegib was 3.0 min. The linearity of an analytical procedure was demonstrated by preparing at seven different concentrations from 0.012 mg/mL to 0.120 mg/mL and analyzed. The calibration curve constructed for Vismodegib by plotting the peak area versus concentration and the regression coefficient (R^2) is 0.9998. The mean recovery was 99.6 and this molecule was sensitive

to peroxide degradation. The degradation was achieved around 6.5% with 3% Hydrogen peroxide at 60°C for 1 hour. Robustness study was performed with variation of flow, mobile phase composition and column temperature but none of the above variations resulted in critical changes to the system analysis parameters. This method was developed and validated for precision, Linearity, accuracy, degradation and robustness as per ICH and FDA guidelines. Hence, the purpose of developing an accurate and stable method along with its validation was achieved for Vismodegib.

Conclusion

A robust and accurate RP-HPLC assay method was developed and validated for estimation of Vismodegib as per ICH/FDA guidelines. This method was validated by using various validation parameters such as system suitability, method and intermediate precision, linearity, accuracy, degradation and robustness. Degradation was performed under acidic, basic, thermal, photolytic, peroxide and hydrolysis conditions. The developed method was sensitive, precise and capable of determine the impurities in stress study and interference as well. Apart from these, this is low cost analytical methodology, shorter run time and it can be applied in various laboratories for estimation of Vismodegib in bulk and pharmaceutical dosage form.

Conflicts of Interest

There are no conflicts to declare.

References

- www.centerwatch.com/drug-information/fda-approved-drugs/drug/1183/erivedge-vismodegib
- www.accessdata.fda.gov/drugsatfda_docs/nda/2012/203388Orig1s000ChemR.pdf
- www.formulationdiary.com/Home/Details/VISMODEGIB
- ICH (2005) Validation of analytical procedures: Text and Methodology Q2(R1). Guidelines: ICH.
- Kim HB (2008) Handbook of stability testing in pharmaceutical development:

- Regulations, Methodologies, and Best Practices. (1st edn). Springer, New York, NY, USA.
6. US FDA (1994) Reviewer Guidance: Validation of Chromatographic Methods. Centre for Drug Evaluation and Research (CDER).
 7. Maheswaran R (2012) FDA perspectives-scientific considerations of forced degradation studies in ANDA submissions. *Pharmaceutical Technology* 36: 73.
 8. Namdeo GS, Bhaskar NB, Sunil MD, Suyog PS, Dipak PS (2013) Pharmaceutical forced degradation studies with regulatory consideration. *Asian J Res Pharm Sci* 3: 178-188.
 9. Ahmad I, Ahmed S, Anwar Z, Muhammad AS, Sikorski M (2016) Photostability and photostability of drug and drug products. *International Journal of Photoenergy* 16: 1-19.