

# Dust Mite Allergen Der f 8 Promotes Th2 Polarization via Modulating Chromatin Structure at the *TIM4* Gene Locus in Dendritic Cells

Mo LH<sup>1</sup>, Yang LT<sup>1,2,3</sup>, Zeng L<sup>1</sup>, Xu LZ<sup>1</sup>, Zhang HP<sup>3</sup>, Li LJ<sup>3</sup>, Liu JQ<sup>1,2,3</sup>, Xiao XJ<sup>1</sup>, Liu ZG<sup>1\*</sup> and Yang PC<sup>\*1,3</sup>

<sup>1</sup>Center of Allergy & Immunology, Shenzhen University School of Medicine, Shenzhen, 518060, PR China

<sup>2</sup>ENT Hospital, Longgang Central Hospital, Shenzhen, 518116, China

<sup>3</sup>Department of Pathology & Molecular Medicine, McMaster University, Hamilton, ON L8N 3Z5, Canada

\*Corresponding authors: Ping-Chang Yang, Center of Allergy & Immunology, Shenzhen University School of Medicine, Shenzhen, 518060, PR China, Tel: 86-755-26536108, E-mail: pcy2356@163.com

Zhi-Gang Liu, Center of Allergy & Immunology, Shenzhen University School of Medicine, Shenzhen, 518060, China, E-mail: lzg@szu.edu.cn

Received date: March 26, 2016; Accepted date: April 28, 2016; Published date: May 02, 2016

Copyright: © 2016 Mo LH, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

Dust mites are an important source of allergens. The mechanism by which dust mite allergens induce allergic diseases is to be further investigated. This study aims to elucidate the role of one of the components of dust mite allergens, the Dermatophagoides farinae-8 (Der f 8), in the induction of airway allergy. In this study, the expression of T cell immunoglobulin and mucin domain 4 (*TIM4*) in dendritic cells (DC) s was analyzed after stimulating with Der f 8 in the culture. The role of Der f 8 in the induction of T helper (Th) 2 inflammation was assessed with an airway allergy mouse model. We observed that, in 34 subtypes of dust mite allergens, Der f 8 uniquely and markedly induced high levels of the *TIM4* expression in DCs by modulating the chromatin at the *TIM4* promoter locus. Der f 8 played a critical role in the expansion of the Th2 response in the mouse airway via inducing DCs to produce *TIM4*. Administration with Der f 8-null dust mite extracts (DME) efficiently inhibited the allergic inflammation and induced regulatory T cells in the airway of mice. We conclude that Der f 8 plays an important role in the pathogenesis of dust mite allergy. Vaccination with the Der f 8-null DME is more efficient to inhibit the dust mite allergic inflammation in the airway than the wild DME.

## Key words:

Experimental allergy; Allergy mechanism; Specific immunotherapy

## Introduction

Allergic diseases have become a worldwide health problem [1,2]. The T helper (Th)2 polarization in the local tissue is one of the major pathologic features of allergy; over production of the antigen specific IgE is another. The IgE binds to the high affinity IgE receptors on mast cells to make mast cells sensitized [3]. Upon re-exposure to specific allergens, the sensitized mast cells release allergy-related mediators to initiate allergic attacks [4]. Although the pathologic process of allergy is somewhat clear, yet, the mechanisms about how antigen-specific Th2 cells get polarization and how they sustain a dominant status in allergic inflammation are still unknown; the therapeutic efficacy on allergic diseases is unsatisfactory currently.

Allergens derived from dust mites are the most common allergens of asthma, atopic dermatitis and perennial rhinitis [5]. Mite allergy affects about 50% of the population with allergic diseases in the world [6]. The dust mite contains a large number of proteins; some of them are allergens. There are more than 30 subtypes of mite allergens have been recognized; some of them have been well characterized; such as Dermatophagoides farina-1 (Der f 1) and Dermatophagoides pteronyssinus (Der p) 1 are important allergens in airway allergy [7,8]. The group 8 allergen (Der f 8) of *D. farina* was cloned and characterized by us; the DNA sequence is listed in the NCBI database (<http://www.ncbi.nlm.nih.gov/nucleotide/KM010011.1>), from which we found that the gene sequence of Der f 8 was quite similar to the gene

sequence of glutathione S-transferase; the latter has a close relation with the pathogenesis of allergic asthma [9]. Previous studies have shown that 40.9 % of mite allergic patients with asthma exhibited IgE reactivity to Der f 8 [10,11]. It reveals that Der f 8 is a significant allergen of mite allergy, but the role of Der f 8 leading to allergic disorders remains to be further investigated.

Besides being the major antigen presenting cells, dendritic cells (DCs) also produce a number of co-stimulatory molecules; one of them is the T cell immunoglobulin mucin domain (TIM)4. We and others previously reported that *TIM4* played a critical role in the expansion of Th2 response and allergic disorders [12-14]. Clinical data show that Der f 8-specific IgE can be detected in the patients with mite allergy [10]. Yet, the mechanism of the over expression of *TIM4* by DCs in allergic diseases is unclear. It is reported that the transcriptional coactivators p300 and STAT6 (Signal transducers and activators of transcription-6) are associated with the pathogenesis of allergic diseases [15,16]; whether these molecules are associated with *TIM4* expression has not been investigated. Based on the above information, we hypothesize that Der f 8 modulates the expression of *TIM4* in DCs, which plays an important role in the pathogenesis of dust mite allergy. In this study, we observed that Der f 8 induced the *TIM4* expression in DCs and played an important role in the dust mite allergen-induced airway Th2 polarization and allergic inflammation. The Der f 8-null dust mite extracts (DME) vaccine showed much better therapeutic efficacy on allergic airway inflammation than the Der f 8-containing DME vaccine in the specific immunotherapy (SIT).

## Materials and Methods

### Research ethic statement

The using mice in the present study was approved by the Research Ethic Committee at Shenzhen University (#SZU2015R032). The experiments were performed in accordance with the approved guidelines.

### Luciferase reporter assay

The luciferase reporter of *TIM4* promoter was constructed with the sequence between -500 and -1047 (or a non-specific sequence, using as a negative control) by the Sangon Biotech Limited Company (Shanghai, China) and transfected DCs following the manufacturer's instruction. In the reporter-gene analysis, the proteins were quantitated with the Bradford reagent (Bio-Rad), and the relevant light intensity was measured with a luminometer with a Luciferase kit (Biomart, China). All assays were done in triplicate and protein expression was evaluated by immunoblot analysis.

### Chromatin IP (ChIP)

The DCs were fixed with 1% formalin for 15 min and sonicated. The supernatant was collected by centrifugation and incubated with antibodies of interest and protein G overnight at 4°C. The beads were collected by centrifugation and washed for 3 times with PBS. The immune complexes on the beads were eluted by SDS buffer and collected by centrifugation. The DNA was purified from the immune complexes and subjected to qPCR with the *TIM4* promoter primers, aggaagagaggagagagca and agaatcgctgaaccagga. The results were calculated and presented as relevant changes against the DNA input.

### A murine model of asthma

Mice were subcutaneously injected on day 1, 3, and 7 with Der f 8 (100 µg/mouse), or DME (100 µg/mouse), or Der f 8-null DME, in 0.1 ml saline respectively. Saline was used as a negative control. From day-15 to day-21, the mice were challenged via nostril with Der f 8 (50 µg/mouse) or DME (50 µg/mouse) in 50 µl PBS once a day for a week. The mice were sacrificed on day 22 to analyze the allergic airway inflammation, or treated with specific Immunotherapy (SIT).

### Airway hypersensitivity response (AHR) measurement

Airway hyperresponsiveness to methacholine was measured with the unrestrained whole-body plethysmography. The mice were subjected to progressively increasing doses of methacholine (0, 6.25, 12.5, 25, 50, 100 µg/mL), and the Penh value was recorded.

### Lung histology

Lung tissues were fixed in 4% phosphate-buffered formalin and embedded in paraffin wax. Sections (3 µm) were cut and stained with hematoxylin and eosin, and observed under a light microscope.

### Specific immunotherapy (SIT)

From day 28, the asthma mice were treated with DME vaccine, or Der f 8-null vaccine, or saline (using as a control) for two weeks with the doses of 5 µg (days 1 and 2), 10 µg (days 3 and 4), 20 µg (days 5-7), 30 µg (days 8 and 9), and 50 µg (days 10-14), in 0.1 ml saline by intraperitoneal injection. Two days before the sacrifice, AHR of the

mice was assessed. One day before the sacrifice, the mice were intranasally challenged with 50 µg DME in 50 µl saline.

### Isolation of mononuclear cells from the lung tissue

The lung tissue was cut into small pieces (2 × 2 × 2 mm), incubated with collagen IV (0.5 mg/ml) at 37°C with mild agitation for 2 h. The cells were filtered with a cell strainer (70 µm), harvested by centrifugation, and cultured in RPMI1640 medium.

### Enzyme-linked immunosorbent assay (ELISA)

Blood samples were collected from each mouse at the sacrifice of decapitation. The sera were isolated to store at -80 °C until use. The serum levels of Th2 cytokines (IL-4, IL-5, and IL-13) were determined by ELISA with commercial reagent kits following the manufacturer's instruction. The levels of specific IgE were determined by an in-house ELISA. Briefly, the microtiter plates were coated with purified allergens at 100 ng/well in 100 µl in carbonate buffered solution (pH 9.5) overnight at 4°C. Then blocked with 100 µl 3% bovine serum albumin (BSA) in PBS for 120 min at 37°C. The serum (diluted to 5 times) or BSA (using as a negative control) was then added to each well (100 µL/well) and incubated for 60 min at room temperature. The plates were incubated with peroxidase-labeled goat anti-mouse IgE (1:2000) for 60 min at 37°C. Each incubating step was followed by 3 washes with phosphate-buffered saline+Tween 20 (PBST). The color was developed by adding tetramethylbenzidine (TMB; 100 µL/well), stayed for 20 min at room temperature and stopped by addition of 2M H<sub>2</sub>SO<sub>4</sub> (50 µL/well). The plates were read with an ELx808 absorbance microplate reader (BioTek, Shanghai, China) at 450 nm.

### Statistical analysis

Data were presented as mean ± SD. Differences between two groups were analyzed Student t-test or by ANOVA if more than two groups. The Newman-Keuls was used for post hoc analysis. A p<0.05 were set as statistically significant. Reagents, mice, the procedures of DC preparation and treatment, Western blotting, qRT-PCR, flow cytometry, purification of immune cells, T cell proliferation assay, preparation of mite proteins, bio-synthetization of mite allergen components and depletion of Der f 8 in DME are presented in supplemental materials.


## Results

### Der f 8 enforces the expression of *TIM4* in DCs

To test if Der f 8 had any effect on regulating *TIM4* expression, we exposed DCs to DME in the culture. The cells were analyzed 48 h later. The results showed that DCs expressed high levels of *TIM4*. Since there are a number of different components of DME allergens have been identified, to recognize which DME components induces *TIM4* expression in DCs is of significance. Thus, we exposed DCs to 34 DME components in the culture for 48 h. The cells were analyzed by RT-qPCR. The results showed that the exposure to Der f 1, 2, 8 and Der p 1, 2 markedly enhanced the expression of *TIM4* in DCs, in which the *TIM4* mRNA was uniquely higher in the DCs exposed to Der f 8 (Figure 1A).

To enforce the results, we exposed DCs to Der f 8 in the culture at gradient doses for 48 h. It resulted in the increases in *TIM4* expression in DCs in a Der f 8 dose-dependent manners (Figure 1B-C). The

results suggest that Der f 8 plays a critical role in the expression of TIM4 in DCs. To strengthen the results, we removed Der f 8 from DME. DCs were stimulated by the Der f 8-null DME or Der f 8-containing DME for 48 h. The results showed that although the wild type DME induced TIM4 expression in DCs, the Der f 8-null DME did not (Figure 1B-C). Since exposure to Der f 1, 2 and Der p 1, 2 also increased the expression of TIM4 mRNA in DCs somehow; we assessed the TIM4 protein levels in DCs after exposure to these 4 subtypes of mite allergen. The results showed that the TIM4 protein was not apparently increased in DCs (Figure 1D). Taken together, the results indicate that Der f 8 can increase the expression of TIM4 in DCs.



**Figure 1:** Der f 8 increases TIM4 expression in DCs. CD11c+ MHC II+ DCs were prepared from the bone marrow. The DCs were cultured in the presence of dust mite extract (DME), or the subtypes of Der f allergens or Der p allergens (denoted on the X axis of panel A; 100 ng/ml) for 48 h. The DCs were collected and analyzed. A, the bars indicate the mRNA levels of TIM4 (by qRT-PCR). B-D, DCs were stimulated in the culture as denoted in the Figure for 48 h. B, the bars indicate the TIM4 mRNA levels. C-D, the Western blots indicate the protein levels of TIM4 after stimulated with the indicated allergens (denoted below the blots; 100 ng/ml) for 48 h. The data of bars are presented as mean  $\pm$  SD. \*,  $p < 0.01$ , compared with the saline group. #,  $p < 0.01$ , compared with the Der f 8 group. The data are representatives of 3 independent experiments. The full length gel graphs are presented in supplemental materials.


### Der f 8 activates the TIM4 promoter

The data of Figure 1 implicate that Der f 8 modulates the expression of TIM4. To test the inference, we transfected DCs with a luciferase reporter of TIM4 promoter. The cells were exposed to Der f 8 (or BSA, using as a negative control) in the culture for 3 h and then subjected to the analysis of the luciferase activity at 24 h. The results showed that the exposure to Der f 8, but not to BSA, markedly increased the luciferase activity (Figure 2). The results indicate that Der f 8 can modulate the TIM4 promoter activity.


### Der f 8 modulates *TIM4* gene chromatin in DCs

We next assessed the phosphorylation status of p300 and STAT6 in DCs after exposure to Der f 8. The results showed that the exposure to Der f 8 increased the phosphorylation of p300 and STAT6 in DCs (Figure 3A-B). Further analysis by ChIP showed that the levels of acetylated H2A, H2B, H3 and H4 at the TIM4 promoter were increased (Figure 3C-H). Finally we observed that the exposure to Der f

8 increased the expression of TIM4 in DCs, which was abolished by the presence of either p300 inhibitor or STAT6 inhibitor (Figure 3I-J). The results indicate that exposure to Der f 8 triggers the expression of TIM4 in DCs.



**Figure 2:** Der f 8 activates *TIM4* promoter. A reporter gene with TIM4 promoter sequence (A) or a non-specific sequence (B) was constructed. C, BMDCs were transfected with the reporter gene. The cells were exposed to Der f 8 (100 ng/ml) in the culture medium for 3 h and analyzed 24 h later. The bars indicate the luciferase activity (mean  $\pm$  SD; \*,  $p < 0.01$ , compared with the medium group). The data are representatives of 3 independent experiments.




**Figure 3:** Der f 8 modulates chromatin of TIM4 gene in DCs; DCs were treated with Der f 8 (100 ng/ml) in the culture for 2 h. The nuclear extracts of DCs were analyzed. A-B, the Western blots indicate the levels of p300/pp300 and STAT6/pSTAT6. C-H, the nuclear extracts were analyzed by ChIP. The bars indicate the binding to the TIM4 promoter by the molecules denoted on the Y axis. I-J, DCs were treated with Der f 8 in the culture for 48 h. I, the bars indicate the mRNA levels of TIM4. J, the Western blots indicate the protein levels of TIM4. Der f 8a: In the presence of both Der f 8 and p300 inhibitor (garcinol: 3  $\mu$ M). Der f 8b: In the presence of Der f 8 and STAT6 inhibitor (A77-1726: 10 nM) The data of bars are presented as mean  $\pm$  SD. \*,  $p < 0.05$ , compared with the time point of “0” (C-H) or the saline group (I). The data are representatives of 3 independent experiments. The full length gel graphs are presented in supplemental materials.

### Der f 8 promotes antigen specific Th2 response


We also assessed the role of Der f 8 in the induction of the antigen specific Th2 responses. We sensitized mice with Der f 8, or DME, or Der f 8-null DME. As shown by Figure 4, the Der f 8 specific CD4+ T cell proliferation, serum levels of Th2 cytokines and Der f 8 specific IgE were higher in mice immunized with Der f 8 than control mice, while much higher serum levels of IL-4, DME specific IgE and DME specific CD4+ T cell proliferation were detected in mice immunized with DME, but not in those immunized with the Der f 8-null DME. The

results suggest that Der f 8 plays an important role in the development of the DME specific Th2 responses. To elucidate if TIM4 is the molecule by which Der f 8 facilitates the DME specific Th2 responses, a group of mice was treated with anti-TIM4 antibody in the course of immunization with DME. Indeed, the Th2 responses were abolished. We also stimulated the sensitized CD4+ T cells with an irrelevant antigen, OVA, in the culture; it did not induce the T cell proliferation. The results demonstrate that that Der f 8 induces TIM4 production by DCs plays a critical role in the development of DME allergy.



### Inhibition of airway allergic inflammation with Der f 8-null DME vaccine

The above data suggest that Der f 8 is a critical molecule in the induction of skewed dust mite-specific Th2 polarization. We hypothesized that the removal of Der f 8 from the DME vaccine might greatly improve the efficacy of the DME-specific immunotherapy (SIT). To test the hypothesis, we developed an airway allergy mouse model with DME as the specific antigen. The mice were treated with a two-week SIT with DME vaccine or Der f 8-null DME vaccine. As shown by Figure 5, the immunization with DME induced lung inflammation, airway hypersensitivity to methacholine challenge, high levels of serum IL-4 and specific IgE, and specific CD4+ T cell proliferation. After DME SIT, although the parameters of lung allergic inflammation were somewhat down regulated, yet none of them reached the significant levels; while those mice treated with the Der f 8-null DME SIT showed a significant attenuation in the lung allergic inflammation. Furthermore, treatment with the Der f 8-null DME SIT, but not the DME SIT, significantly up regulated the frequency of antigen specific CD4+ CD25+ Foxp3+ regulatory T cells (Treg) in the lung.



### Discussion

We previously found that DCs produced TIM4 upon the exposure to Staphylococcal enterotoxin B (SEB) [12] that played a critical role in the induction of intestinal allergy [17]. Rodriguez-Manzanet et al also reported that after activation, DCs produced TIM4 that induced T cell expansion and survival [14]. Li et al reported recently that hepatic ischemia reperfusion injury could increase TIM4 expression by DCs [18]. The present data have expanded the knowledge in the studies of TIM4. After exposing to Der f 1, Der f 2, Der p 1 and 2, the transcripts of TIM4 in DCs were markedly increased, which was much higher in DCs after exposing to Der f 8.

From the time course we found that, upon exposure to Der f 8, the phosphorylation of p300 occurred, followed by the acetylation of histone at the TIM4 promoter locus, and then the STAT6 phosphorylation was increased. These data suggest that Der f 8 can modulate the chromatin of the TIM4 promoter, and induces the *TIM4* gene transcription and expression in DCs. Both p300 and STAT6 play an important role in the Der f 8-induced TIM4 expression because the presence of either a p300 inhibitor or a STAT6 inhibitor abolished the TIM4 expression.

Structurally, Der f 8 is similar to the glutathione S-transferase (GST). One of the GST functions is to bind both the substrate at the enzyme's hydrophobic H-site and glutathione at the adjacent, hydrophilic G-site, which together form the active site of the enzyme; and subsequently to activate the thiol group of glutathione, enabling the nucleophilic attack upon the substrate [19]. Our data show that Der f 8 has a similar feature of GST that activates p300 in DCs and

induced p300 phosphorylation. This action of Der f 8 is of significance because it initiated *TIM4* gene transcription in DCs, induced an Der f 8-specific Th2 polarization and facilitated the DME-specific allergic inflammation in the lung, as shown by the present study. Supportive data have also published that GST is associated with the pathogenesis of asthma [9].

In line with previous studies [20,21], we created a DME-specific airway allergy mouse model without an extra adjuvant. A significant finding in the present study is that the capacity of inducing the specific Th2 polarization in the airway was significantly attenuated with the Der f 8-null DME, pinpointing the importance of Der f 8 in the allergy-induction capacity of dust mite. Such a feature of Der f 8 implicates that the removal of Der f 8 from DME may be useful in the SIT using DME as a vaccine. The results from vaccination with the Der f 8-null DME on inhibiting DME-specific allergic airway inflammation support the inference.

To induce Tregs in the allergic subjects is one of the therapeutic purposes of SIT [22]. The present data also show that using the Der f 8-null DME vaccine recovered the induction of Tregs in the mice with dust mite allergy, which did not occur in those allergic mice treated with the Der f 8-containing DME. This is in line with previous studies. Li et al reported that in a DC: CD4<sup>+</sup> T cell co-culture system, blockade of TIM4 on DCs significantly inhibited Th2 cell differentiation and facilitated the induced CD4<sup>+</sup> CD25<sup>+</sup> Foxp3<sup>+</sup> Treg expansion [18]. Yeung et al indicate that interruption of DC-mediated TIM4 signaling induces regulatory T cells and promotes skin allograft survival [23]. To induce Tregs in an allergic environment is generally refractory. The present data provide a possible strategy to enhance the efficacy in the induction of Tregs by suppressing the production of TIM4.

In summary, the present data have revealed a previously unknown feature of dust mite allergens, one of the components of the dust mite allergens, the Der f 8, is capable of inducing TIM4 expression in DCs, which plays a critical role in the initiation of skewed Th2 polarization and allergic inflammation. Removal of Der f 8 from DME vaccine significantly improves the inhibitory efficacy on the dust mite-specific allergic airway inflammation. Thus, the Der f 8-null DME may be a promising vaccine to obtain a better therapeutic effect on inhibition of the dust mite-specific allergic inflammation.

## Acknowledgement

This study was supported by grants from the Natural Science Foundation of SZU (No.000004), the innovation of science and Technology Commission of Shenzhen Municipality (No.JCYJ20140418095735611; ZDSYS201506050935272), the Natural Science Foundation of China (81373176, 31400856, 31570932, 81571790 and 81501573) and Canadian Institutes of Health Research (CIHR; #191063).

## References

1. Fiocchi A, Pawankar R, Cuello-Garcia C, Ahn K, Al-Hammadi S, et al. (2015) World Allergy Organization-McMaster University Guidelines for Allergic Disease Prevention (GLAD-P): Probiotics. *World Allergy Organ J* 8:4.
2. Nordlund B, Melén E, Schultz ES, Grönlund H, Hedlin G, et al. (2014) Prevalence of severe childhood asthma according to the WHO. *Respir Med* 108: 1234-1237.
3. An S, Shen C, Liu X, Chen L, Xu X, et al. (2013) Alpha-actinin is a new type of house dust mite allergen. *PLoS One* 8: e81377.
4. Taketomi EA, Silva DA, Sopelete MC, Gervasio AM, Alves R, et al (2006) Differential IgE reactivity to Der p 1 and Der p 2 allergens of *Dermatophagoides pteronyssinus* in mite-sensitized patients. *J Invest Allergol Clin Immunol* 16: 104-109.
5. Eifan AO, Calderon MA, Durham SR (2013) Allergen immunotherapy for house dust mite: clinical efficacy and immunological mechanisms in allergic rhinitis and asthma. *Expert Opin Biol Ther* 13: 1543-1556.
6. Valenta R, Linhart B, Swoboda I, Niederberger V (2011) Recombinant allergens for allergen-specific immunotherapy: 10 years anniversary of immunotherapy with recombinant allergens. *Allergy* 66: 775-783.
7. Segundo GR, Sopelete MC, Terra SA, Pereira FL, Justino CM, et al. (2009) Diversity of allergen exposure: implications for the efficacy of environmental control. *Braz J Otorhinolaryngol* 75: 311-316.
8. Lim FL, Hashim Z, Than LT, Md Said S, Hisham Hashim J, et al. (2015) Airway Symptoms and Rhinitis in Office Workers in Malaysia: Associations with House Dust Mite (HDM) Allergy, Cat Allergy and Levels of House Dust Mite Allergens in Office Dust. *PLoS One* 10: e0124905.
9. Polosukhin VV, Polosukhin IV, Hoskins A, Han W, Abdolrasulnia R, et al. (2014) Glutathione S-transferase M1 modulates allergen-induced NF-kappaB activation in asthmatic airway epithelium. *Allergy* 69: 1666-1672.
10. Cui YB, Zhou Y, Wang N, Teng FX, Yu LL, et al. (2014) Expression, cloning, and IgE-binding of the full-length dust mite allergen Der f 8. *Immunol Res* 60: 60-68.
11. Jeong KY, Lee JY, Son M, Yi MH, Yong TS, et al. (2015) Profiles of IgE Sensitization to Der f 1, Der f 2, Der f 6, Der f 8, Der f 10, and Der f 20 in Korean House Dust Mite Allergy Patients. *Allergy Asthma Immunol Res* 7: 483-488.
12. Yang PC, Xing Z, Berin CM, Soderholm JD, Feng BS, et al. (2007) TIM-4 expressed by mucosal dendritic cells plays a critical role in food antigen-specific Th2 differentiation and intestinal allergy. *Gastroenterology* 133: 1522-1533.
13. Meyers JH, Chakravarti S, Schlesinger D, Illes Z, Waldner H, et al. (2005) TIM-4 is the ligand for TIM-1, and the TIM-1-TIM-4 interaction regulates T cell proliferation. *Nat Immunol* 6: 455-464.
14. Rodriguez-Manzanet R, Meyers JH, Balasubramanian S, Slavik J, Kassam N, et al. (2008) TIM-4 expressed on APCs induces T cell expansion and survival. *J Immunol* 180: 4706-4713.
15. Gunawardhana LP, Gibson PG, Simpson JL, Powell H, Baines KJ (2014) Activity and expression of histone acetylases and deacetylases in inflammatory phenotypes of asthma. *Clin Exp Allergy* 44: 47-57.
16. Noval Rivas M, Burton OT, Wise P, Charbonnier LM, Georgiev P, et al. (2015) Regulatory T cell reprogramming toward a Th2-cell-like lineage impairs oral tolerance and promotes food allergy. *Immunity* 42: 512-523.
17. Feng BS, Chen X, He SH, Zheng PY, Foster J, et al. (2008) Disruption of T-cell immunoglobulin and mucin domain molecule (TIM)-1/TIM4 interaction as a therapeutic strategy in a dendritic cell-induced peanut allergy model. *J Allergy Clin Immunol* 122: 55-61.
18. Li J, Zhao X, Liu X, Liu H (2015) Disruption of TIM-4 in dendritic cell ameliorates hepatic warm IR injury through the induction of regulatory T cells. *Mol Immunol* 66: 117-125.
19. Eaton DL, Bammler TK (1999) Concise review of the glutathione S-transferases and their significance to toxicology. *Toxicol Sci* 49: 156-164.
20. Al-Garawi AA, Fattouh R, Walker TD, Jamula EB, Botelho F, et al. (2009) Acute, but Not Resolved, Influenza A Infection Enhances Susceptibility to House Dust Mite-Induced Allergic Disease. *J Immunol* 182: 3095-3104.
21. Cates EC, Fattouh R, Wattie J, Inman MD, Goncharova S, et al. (2004) Intranasal exposure of mice to house dust mite elicits allergic airway inflammation via a GM-CSF-mediated mechanism. *J Immunol* 173: 6384-6392.
22. Verhagen J, Wegner A, Wraith DC (2015) Extra-thymically induced T regulatory cell subsets: the optimal target for antigen-specific immunotherapy. *Immunology* 145: 171-181.
23. Yeung MY, McGrath MM, Nakayama M, Shimizu T, Boenisch O, et al. (2013) Interruption of dendritic cell-mediated TIM-4 signaling induces

regulatory T cells and promotes skin allograft survival. *J Immunol* 191: 4447-4455.