

## Future Perspectives for Cancer Therapy Using the CRISPR Genome Editing Technology

Noah Isakov\*

*The Shraga Segal Department of Microbiology, Immunology and Genetics, Faculty of Health Sciences and The Cancer Research Center, Ben Gurion University of the Negev, Beer Sheva, Israel*

\*Corresponding author: Noah Isakov, The Shraga Segal Department of Microbiology, Immunology and Genetics, Faculty of Health Sciences, Ben Gurion University of the Negev, P.O.B. 653, Beer Sheva 84105, Israel, Tel: 972-8-6477267; Fax: 972-8-6477626; E-mail: [noah@bgu.ac.il](mailto:noah@bgu.ac.il)

Received date: May 25, 2017; Accepted date: May 30, 2017; Published date: June 07, 2017

Copyright: © 2017 Noah Isakov, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Introduction

The term cancer relates to a large group of diseases characterized by abnormal growth of cells which are capable of invading nearby and even remote tissues. Cancer can occur by numerous types of disease-causing genetic alterations leading to activation of oncogenes or inactivation of suppressor genes in almost any cell type in our body.

Over the last two decades, substantial progress has been made in cancer diagnosis and treatment, resulting in a steady improvement in cancer survival. While cancers can be treated using several different protocols, matching a protocol to a patient depends on the type and stage of cancer, in addition to age, sex, race, and the overall health of the patient, as well as considerations relevant to the potential side effects. The most common types of cancer treatment include surgery, chemotherapy, radiation therapy, hormonal therapy, targeted therapy, and often, combinations of two or more types of the above therapies. However, the rapidly changing field of genome engineering and gene therapy promises a number of new and novel potential strategies for treatment of cancer.

One of the most recently developed strategies for cancer treatment is based on a technology utilizing the clustered regularly interspaced short palindromic repeats (CRISPR) and the CRISPR-associated genes (cas). The CRISPR-cas system is naturally occurring gene-editing machinery in prokaryotes which operates as an adaptive and heritable immune system enabling bacteria and archaea to defeat invading viruses. Understanding of the mode of operation of the CRISPR-cas-mediated gene editing machinery, followed by adaptation of this system for manipulation of eukaryotic cells, has revolutionized genomic engineering, allowing researchers to target specific genes, correct mutations at precise locations, and modify DNA in essentially any cell and organism. Thus, the CRISPR technology has the potential to revolutionize cancer gene therapy and enable the correction of defective genes which cause a variety of devastating diseases, including cystic fibrosis, muscular dystrophy, and hemophilia.

The remarkable accomplishments obtained using CRISPR-Cas9 across multiple biological disciplines has led the journal Science to select CRISPR-Cas9 as the '2015's Breakthrough of the Year' [1].

A first hint for the existence of a CRISPR loci was obtained two decades ago, when Ishino et al. sequenced the *iap* gene in *Escherichia coli* and noted the existence of repeated DNA sequences downstream of the gene, which at that time, were considered to have no apparent biological significance [2]. Further studies by Mojica et al. noted the presence in archaea of repetitive and palindromic sequences separated from each other by short variable DNA sequences (called 'spacers') and comprising a leader sequence at their start [3,4]. These sequences were

termed CRISPR, an acronym for "Clustered Regularly Interspaced Short Palindromic Repeats" [4].

The bacterial and archaeal CRISPR sequences were located near CRISPR-associated (cas) genes which encoded nucleases capable of cutting and degrading exogenous DNA at specific sequence sites [5]. In addition, many of the spacer sequences within CRISPR matched those of bacteriophages and plasmids, which could represent genetic signatures of past invaders [6-8]. The latter findings and the observations that CRISPR loci are transcribed [9], suggested that CRISPR-cas functions as a novel prokaryotic adaptive defense system that might utilize antisense RNA as memory signatures of past invasions [10].

Additional support for this assumption was obtained in studies of *Streptococcus thermophilus* where a direct correlation was found between the number of spacers in the CRISPR locus and the bacterial ability to resist infection by phages [6]. Analysis of the *Streptococcus thermophilus* genome before and after a phage challenge confirmed that the CRISPR-cas system is capable of cutting foreign DNA and integrate small fragments of this digested DNA into the host genome. Removal or addition of particular spacers modified the bacterial resistance to the phage [11], indicating that specificity of the CRISPR-cas-mediated bacterial resistance to phages is determined by the similarity of spacer-phage sequence. Thus, the CRISPR-cas system, which was found in approximately 90% of sequenced archaea and 50% of sequenced bacterial genomes [12,13], serves as a heritable adaptive immune defense system in a majority of prokaryotes [14,15]. Based on the signature of the Cas proteins and features of the architecture of cas loci, the current CRISPR-cas systems encompass two classes, five types and 16 subtypes [16,17]. However, the existence of currently unclassifiable rare variants and the potential future sequencing of additional types of bacterial and archaeal genomes may well increase these numbers.

The elucidation of the CRISPR-cas-based gene editing machinery and understanding its mode of operation, led to the development of a CRISPR-cas-based biotechnological tool which enables the manipulation of eukaryotic genomes [18-21] and is useful in numerous applications in basic science, biotechnology and medicine [22]. Among its many potential applications, the CRISPR-cas9 technology can help correct genetic defects, prevent the spread of a wide range of diseases, and restore and enhance the immune system's ability to fight cancer. Adaptation of the CRISPR technology to cancer therapy is currently at its very early stages, but is gaining momentum and is beginning to be implemented using different approaches.

A pioneering study by Chen et al. utilized the CRISPR-Cas9 technology to specifically target cancer-causing genes in human

prostate and liver cancers and replace them with a cell death-promoting gene [23]. The selected target genes, TMEM135-CCDC67 and MAN2A1-FER, were identified in human prostate cancer [24] and hepatocellular carcinoma [25], respectively, and are responsible for the rapid growth and invasiveness of the cancer cells. They represent abnormal hybrid genes formed by deletions and re-joining of DNA sequences of two previously independent genes. As a consequence, the newly formed fusion genes contain cancer cell-specific sequence breakpoints, which are not present in normal cells, and therefore provide unique targets for therapeutic intervention.

Chen et al. used two specifically designed adenoviruses to deliver the gene-editing tools into the cancer cells. One virus was designed to deliver a mutated Cas9 (Cas9D10A), which produces single-strand breaks at the target DNA, plus guide RNAs, which target the cancer cell-specific fusion gene breakpoint sequences, thus allowing the removal of the mutated DNA of the fusion gene. A second virus was designed to deliver into the cancer cells a DNA construct encoding the herpes simplex virus types 1 (HSV1) thymidine kinase (TK), which is flanked by sequences homologous to those surrounding the breakpoint, and thereby enabling the replacement of the fusion gene by HSV1-TK.

HSV1-TK is a commonly used suicide gene for different tumor cell types [26]. It is a phosphotransferase which can convert thymidine into a monophosphate thymidine, a building block for DNA synthesis. However, unlike its mammalian TK counterpart, HSV1-TK can also phosphorylate ganciclovir (prodrug), a synthetic nucleoside homolog, which can be further phosphorylated in the cells to triphosphate ganciclovir, a poor substrate for chain elongation which disrupts DNA synthesis and promotes cell death [27]. HSV1-TK-negative mammalian cells are immune to ganciclovir because of their inability to phosphorylate it, while the entire gene-editing approach used by Chen et al. is highly specific because the *HSV1-TK* gene is specifically targeted to fusion gene sequences which are present only in the cancer cells.

To test the applicability of their system to cancer therapy, Chen et al. transplanted human prostate and liver cancer cells carrying the TMEM135-CCDC67 and MAN2A1-FER fusion genes, respectively, to severe combined immunodeficiency (SCID) mice, and allowed the tumor cells to grow for 3 weeks. When tumors were visible, the mice were infected with the adenoviruses and treated with ganciclovir. This resulted in a very dramatic effect in which the treatment not only halted the exponential growth of the tumors, but also induced a ~30% shrinkage in their total mass and prevented metastases formation. This study demonstrates that the CRISPR-Cas-mediated defective gene replacement by a suicide-gene may represent a novel future strategy for cancer therapy.

A different CRISPR-Cas9-based cancer therapy approach was employed by Eyquem et al., who attempted to boost the immune response against cancer cells by engineering T lymphocytes to express a highly specific chimeric antigen receptor (CAR) [28]. Previous studies demonstrated that CARs, which are highly specific against cancer cell antigens, can be engineered in the laboratory, and upon their expression in T cells, they can direct efficient T cell-mediated killing of antigen-expressing cancer cells [29]. Clinical trials using CAR-modified T cells (CAR-T cell) directed against CD19 [30], CD20 [31], CD22 [32] and CD30 [33] demonstrated promising outcomes in patients with B cell malignancies. A most remarkable result was obtained using CD19-directed CAR-T cells in B cell acute lymphoblastic leukemia (B-ALL) patients, where a complete remission

was obtained in more than 80% of the patients [30,34]. The success of the CAR-T cell-mediated therapy in hematological malignancies was hampered by the therapy-mediated tumor antigen loss which led to tumor relapse. These data emphasize the need for multiple types of CAR-T cells directed against a battery of different tumor antigens which will remain effective even after a loss of one or more tumor antigens.

In the study by Eyquem et al., the authors employed the CRISPR-Cas9 genome editing system to insert a CD19-specific CAR to the T-cell receptor  $\alpha$  (TCR $\alpha$ ) constant locus. The authors showed that the newly produced T cells were potent and highly specific *in vitro* and were also able to kill human acute lymphoblastic leukemia cells *in vivo* in a mouse model of cancer.

Recently, Nature magazine published the first-ever CRISPR gene-editing application in human cancer immunotherapy [35,36]. This clinical trial was performed by researchers and clinicians from the Sichuan University in Chengdu and the West China Hospital in China, who utilized the CRISPR-Cas9 technology to switch off a gene encoding the programmed cell death protein 1 (PD-1; CD279) in T lymphocytes derived from a lung cancer patient. They then propagated the genetically engineered T lymphocytes *in vitro* and injected them back to the patients' blood system in anticipation of boosting the T cell-mediated immune response against the cancer cells.

The PD-1 cell surface receptor is an immune checkpoint protein that promotes self-tolerance and downregulates T cell-mediated responses [37,38]. Many types of tumor cells express the PD-1 ligand (PD-L1) on their surface and exploit this immune-checkpoint pathway as a mechanism to evade immune destruction. Upon interaction of tumor cells PD-L1 surface proteins with the PD-1 receptors on the surface of T cells, they deliver a signal which inhibits T cell-mediated cytotoxic responses against the tumor cells.

Some retrospective studies demonstrated that overexpression of PD-L1 on tumor cells is closely related to the poor prognosis and high invasiveness of non-small-cell lung carcinoma (NSCLC) [39] and hepatocellular carcinoma [40]. In addition, clinical trials in NSCLC patients demonstrated that administration of pembrolizumab (Keytruda) or nivolumab (Opdivo) anti-PD-1 blocking antibodies enhanced the endogenous anti-tumor immune response and extended the overall survival of a small fraction of the NSCLC patients [38,41]. While the major aim of the clinical studies that utilize CRISPR-based technology to switch off PD-1 is to test the CRISPR's safety in gene therapy application, it is expected that inactivation of the PD-1 gene will become a more efficient therapeutic strategy for the treatment of NSCLC and other types of cancer.

Recently, a new CRISPR system, termed CRISPR-Cpf1, was discovered in the gram-negative pathogenic bacteria, *Francisella novicida* [42]. This new system presents simpler and more precise genome engineering machinery and provides flexibility in the selection of target sites. Future utilization of alternative CRISPR-based systems will enable flexibility in choosing a system that is more appropriate for targeting the genes of choice.

Yin et al. recently revealed that the CRISPR-Cas9 technology can be used to excise the HIV-1 DNA from the genome of a "humanized" mouse that carries HIV-1-infected human lymphocytes [43]. The authors also demonstrated that the gene-editing apparatus can be engineered to carry a set of multiple guide RNAs, all designed to efficiently excise integrated HIV-1 DNA from the host cell genome, a strategy that might well overcome a potential escape of mutated HIV-1

viruses. This proof-of-concept study indicates that the CRISPR-Cas9 technology may be useful in the design of new treatments directed against virally-induced malignancies, such as the Epstein-Barr virus (EBV)-induced Burkitt lymphoma [44], human T-lymphotropic virus leukemia (HTLV) [45], and Kaposi sarcoma-associated herpes virus (KSHV or HHV8) [46], as well as other life-threatening diseases induced by viruses such as the Ebola [47], Marburg [47], and rabies [48].

Future design of new strategies for CRISPR-Cas9-based cancer therapy and improvements of the efficiency of the gene-editing strategies will open new and more promising avenues for cancer therapy.

## Acknowledgements

Work in our laboratory is funded in part by the USA-Israel Binational Science Foundation, the Israel Science Foundation administered by the Israel Academy of Science and Humanities, and donations by Mr. Martin Kolinsky and Ms. Linda Osofsky. N.I. holds the Joseph H. Krupp Chair in Cancer Immunobiology.

## References

1. Travis J (2015) Making the cut. *Science* 350: 1456-1457.
2. Ishino Y, Shinagawa H, Makino K, Amemura M, Nakata A (1987) Nucleotide sequence of the *iap* gene, responsible for alkaline phosphatase isozyme conversion in *Escherichia coli*, and identification of the gene product. *J Bacteriol* 169: 5429-5433.
3. Mojica FJ, Ferrer C, Juez G, Rodríguez-Valera F (1995) Long stretches of short tandem repeats are present in the largest replicons of the Archaea *Haloferax mediterranei* and *Haloferax volcanii* and could be involved in replicon partitioning. *Mol Microbiol* 17: 85-93.
4. Mojica FJ, Díez-Villaseñor C, Soria E, Juez G (2000) Biological significance of a family of regularly spaced repeats in the genomes of Archaea, Bacteria and mitochondria. *Mol Microbiol* 36: 244-246.
5. Jansen R, Embden JD, Gastra W, Schouls LM (2002) Identification of genes that are associated with DNA repeats in prokaryotes. *Mol Microbiol* 43: 1565-1575.
6. Bolotin A, Quinquis B, Sorokin A, Ehrlich SD (2005) Clustered regularly interspaced short palindrome repeats (CRISPRs) have spacers of extrachromosomal origin. *Microbiology* 151: 2551-2561.
7. Mojica FJ, Díez-Villaseñor C, García-Martínez J, Soria E (2005) Intervening sequences of regularly spaced prokaryotic repeats derive from foreign genetic elements. *J Mol Evol* 60: 174-182.
8. Pourcel C, Salvignol G, Vergnaud G (2005) CRISPR elements in *Yersinia pestis* acquire new repeats by preferential uptake of bacteriophage DNA, and provide additional tools for evolutionary studies. *Microbiology* 151: 653-663.
9. Tang TH, Bachelier JP, Rozhdestvensky T, Bortolin ML, Huber H, et al. (2002) Identification of 86 candidates for small non-messenger RNAs from the archaeon *Archaeoglobus fulgidus*. *Proc Natl Acad Sci U S A* 99: 7536-7541.
10. Makarova KS, Grishin NV, Shabalina SA, Wolf YI, Koonin EV (2006) A putative RNA-interference-based immune system in prokaryotes: computational analysis of the predicted enzymatic machinery, functional analogies with eukaryotic RNAi, and hypothetical mechanisms of action. *Biol Direct* 1: 7.
11. Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, et al. (2007) CRISPR provides acquired resistance against viruses in prokaryotes. *Science* 315: 1709-1712.
12. Grissa I, Vergnaud G, Pourcel C (2007) The CRISPRdb database and tools to display CRISPRs and to generate dictionaries of spacers and repeats. *BMC Bioinformatics* 8: 172.
13. Bhaya D, Davison M, Barrangou R (2011) CRISPR-Cas systems in bacteria and archaea: versatile small RNAs for adaptive defense and regulation. *Annu Rev Genet* 45: 273-297.
14. Horvath P, Barrangou R (2010) CRISPR/Cas, the immune system of bacteria and archaea. *Science* 327: 167-170.
15. Marraffini LA, Sontheimer EJ (2010) CRISPR interference: RNA-directed adaptive immunity in bacteria and archaea. *Nat Rev Genet* 11: 181-190.
16. Makarova KS, Haft DH, Barrangou R, Brouns SJ, Charpentier E, et al. (2011) Evolution and classification of the CRISPR-Cas systems. *Nat Rev Microbiol* 9: 467-477.
17. Makarova KS, Wolf YI, Alkhnbashi OS, Costa F, Shah SA, et al. (2015) An updated evolutionary classification of CRISPR-Cas systems. *Nat Rev Microbiol* 13: 722-736.
18. Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA (2012) A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337: 816-821.
19. Jinek M, East A, Cheng A, Lin S, Ma E, et al. (2013) RNA-programmed genome editing in human cells. *Elife* 2: e00471.
20. Cong L, Ran FA, Cox D, Lin S, Barretto R (2013) Multiplex genome engineering using CRISPR/Cas systems. *Science* 339: 819-823.
21. Mali P, Yang L, Esvelt KM, Aach J, Guell M, et al. (2013) RNA-guided human genome engineering via Cas9. *Science* 339: 823-826.
22. Komor AC, Badran AH, Liu DR (2017) CRISPR-Based Technologies for the Manipulation of Eukaryotic Genomes. *Cell* 168: 20-36.
23. Chen ZH, Yu YP, Zuo ZH, Nelson JB, Michalopoulos GK, et al. (2017) Targeting genomic rearrangements in tumor cells through Cas9-mediated insertion of a suicide gene. *Nat Biotechnol*.
24. Yu YP, Ding Y, Chen Z, Liu S, Michalopoulos A, et al. (2014) Novel fusion transcripts associate with progressive prostate cancer. *Am J Pathol* 184: 2840-2849.
25. Chen ZH, Yu YP, Tao J, Liu S, Tseng G, et al. (2017) MAN2A1-FER Fusion Gene Is Expressed by Human Liver and Other Tumor Types and Has Oncogenic Activity in Mice. *Gastroenterology*.
26. Caruso M, Panis Y, Gagandeep S, Houssin D, Salzmänn JL, et al. (1993) Regression of established macroscopic liver metastases after *in situ* transduction of a suicide gene. *Proc Natl Acad Sci USA* 90: 7024-7028.
27. St Clair MH, Lambe CU, Furman PA (1987) Inhibition by ganciclovir of cell growth and DNA synthesis of cells biochemically transformed with herpesvirus genetic information. *Antimicrob Agents Chemother* 31: 844-849.
28. Eyquem J, Mansilla-Soto J, Giavridis T, van der Stegen SJ, Hamieh M, et al. (2017) Targeting a CAR to the TRAC locus with CRISPR/Cas9 enhances tumour rejection. *Nature* 543: 113-117.
29. Wang Z, Wu Z, Liu Y, Han W (2017) New development in CAR-T cell therapy. *J Hematol Oncol* 10: 53.
30. Maude SL, Frey N, Shaw PA, Aplenc R, Barrett DM, et al. 2014. Chimeric antigen receptor T cells for sustained remissions in leukemia. *N Engl J Med* 371: 1507-1517.
31. Till BG, Jensen MC, Wang J, Chen EY, Wood BL (2008) Adoptive immunotherapy for indolent non-Hodgkin lymphoma and mantle cell lymphoma using genetically modified autologous CD20-specific T cells. *Blood* 112: 2261-2271.
32. Haso W, Lee DW, Shah NN, Stetler-Stevenson M, Yuan CM, et al. (2013) Anti-CD22-chimeric antigen receptors targeting B-cell precursor acute lymphoblastic leukemia. *Blood* 121: 1165-1174.
33. Wang CM, Wu ZQ, Wang Y, Guo YL, Dai HR, et al. (2017) Autologous T Cells Expressing CD30 Chimeric Antigen Receptors for Relapsed or Refractory Hodgkin Lymphoma: An Open-Label Phase I Trial. *Clin Cancer Res* 23: 1156-1166.
34. Lee DW, Kochenderfer JN, Stetler-Stevenson M, Cui YK, Delbrook, et al. CI 2015. T cells expressing CD19 chimeric antigen receptors for acute lymphoblastic leukaemia in children and young adults: a phase 1 dose-escalation trial. *Lancet* 385: 517-528.
35. Cyranoski D (2016) Chinese scientists to pioneer first human CRISPR trial. *Nature* 535: 476-477.

36. Cyranoski D (2016) CRISPR gene-editing tested in a person for the first time. *Nature* 539: 479.
37. Pardoll DM (2012) The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer* 12: 252-264.
38. Ying H, Kang Y, Zhang H, Zhao D, Xia J, et al. (2015) MiR-127 modulates macrophage polarization and promotes lung inflammation and injury by activating the JNK pathway. *J Immunol* 194: 1239-1251.
39. Mu CY, Huang JA, Chen Y, Chen C, Zhang XG (2011) High expression of PD-L1 in lung cancer may contribute to poor prognosis and tumor cells immune escape through suppressing tumor infiltrating dendritic cells maturation. *Med Oncol* 28: 682-688.
40. Gao Q, Wang XY, Qiu SJ, Yamato I, Sho M, et al. (2009) Overexpression of PD-L1 significantly associates with tumor aggressiveness and postoperative recurrence in human hepatocellular carcinoma. *Clin Cancer Res* 15: 971-979.
41. Killock D (2016) Lung cancer: Anti-PD-1 therapy in the frontline. *Nat Rev Clin Oncol* 13: 715.
42. Zetsche B, Gootenberg JS, Abudayyeh OO, Slaymaker IM, Makarova KS, et al. (2015) Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. *Cell* 163: 759-771.
43. Yin C, Zhang T, Qu X, Zhang Y, Putatunda R, et al. (2017) In Vivo Excision of HIV-1 Provirus by saCas9 and Multiplex Single-Guide RNAs in Animal Models. *Mol Ther* 25: 1168-1186.
44. Klein G (2009) Burkitt lymphoma--a stalking horse for cancer research? *Semin Cancer Biol* 19: 347-350.
45. Ciminale V, Rende F, Bertazzoni U, Romanelli MG (2014) HTLV-1 and HTLV-2: highly similar viruses with distinct oncogenic properties. *Front Microbiol* 5: 398.
46. Dittmer DP, Damania B (2016) Kaposi sarcoma-associated herpesvirus: immunobiology, oncogenesis, and therapy. *J Clin Invest* 126: 3165-3175.
47. Rougeron V, Feldmann H, Grard G, Becker S, Leroy EM (2015) Ebola and Marburg haemorrhagic fever. *J Clin Virol* 64: 111-119.
48. Velasco-Villa A, Mauldin MR, Shi M, Escobar LE, Gallardo-Romero NF, et al. (2017) The history of rabies in the Western Hemisphere. *Antiviral Res.*