

Length – Weight Relationships of Twelve Fishes from the River Padma near Rajshahi City, Bangladesh

Golam Mortuza M^{1,2*} and Fahad A. Al-Misned¹

¹Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

²Department of Zoology, Faculty of Life and Earth Science, Rajshahi University, Rajshahi 6205, Bangladesh

Abstract

The objective of this study was to evaluate the length-weight relationships (LWRs) of freshwater fish species, which serves as a baseline for comparison to other relatively altered tropical Bangladeshi rivers. The LWRs of 12 species belonging to four classes, seven families and 11 genera (*Gagata youssoufi*, *Cirrhinus reba*, *Clupisoma garua*, *Ompok bimaculatus*, *Pangasius pangsius*, *Securicula gora*, *Ailia coila*, *Chanda nama*, *Parambassis ranga*, *Botia lohachata*, *Rhinomugil corsula* and *Labeo boga*) captured from the River Padma near Rajshahi City, Bangladesh, were studied. The allometric coefficient (b) of the LWRs indicated positive allometric growth in *P. pangsius*, *A. coila*, *C. reba*, *B. lohachata* and *C. nama* ($b > 3.00$), but negative allometric growth in *G. youssoufi*, *C. garua*, *O. bimaculatus*, *S. gora*, *P. ranga*, *R. corsula* and ($b < 3.00$). The results would be useful for sustainable management and conservation of the limited stocks in the Padma River ecosystem.

Keywords: Length-weight relationship; Allometric; Isometric; Growth; Padma River

Introduction

Length-weight relationships have been used extensively for the conversion of growth-in-length equations to growth-in-weight for use in stock assessment models to estimate the stock assessment, compare the life histories of certain species and other aspects of fish population dynamics [1-6]. It also allows for the study of the ontogenetic allometric changes in fish growth [7] and possible effects from parasites [8].

This study describes the length-weight relationships (LWRs) of 12 least concern and near threaten species according to IUCN Red list from the Padma River, near Rajshahi city, Bangladesh [9]. These fishes species once abundant in rivers, streams, canals, reservoirs, lakes, ponds and beel, haor and baor swamplands of Bangladesh, India, Nepal and Sri-Lanka [10], the populations are in serious decline due to over-exploitation augmented by various ecological changes and degradation of their natural habitats. The Length-Weight Relationships (LWRs) of threatened fishes are the most important biological parameters to provide information on the growth and condition of fish species and the entire fish community, and are highly significant for management and conservation of natural populations [11,12]. Length-weight relationships have been reported for some commercially important fishes from the River Padma, but data for most of the endemic freshwater fish species are still missing. The present study was undertaken with the objective to estimate the length-weight relationships for 12 fish species collected from the River Padma near Rajshahi, Bangladesh.

Materials and Methods

The Padma is one of the largest rivers of Bangladesh. It is the main tributary of the Ganges which originates in the Gangotri glacier of the Himalayan. The part of the Ganga in Bangladesh is known as the Padma which enters Bangladesh from India (Murshidabad district) at Shibganj Upazila (Manakosha and Durlavpur unions) of Chapai Nawabganj district. Its length in Bangladesh is 366 kilometers.

This study was conducted from the year 2009-2011 on the Padma River near Rajshahi City, (24°21' N and 88°36'E) Bangladesh. Monthly samples were collected from different fishing spots and fish markets of Rajshahi City, June 2012 to May 2013. The main gear used by

commercial fishers included cast nets, gill nets, drag nets, mosquito nets, hand nets as well as traditional fishing traps. The fresh samples were immediately chilled in ice on site and fixed with 10% buffered formalin upon arrival at the laboratory then identified according to [13-15] Jayaram, Rahman, Talwar, Jhingran. All morphometric measurements were taken according to [16] Froese and Pauly. All fishes were individually measured to the nearest 0.01 mm and weighed on a digital electronic balance (Bosch EP 628) to the nearest 0.01 g. The LWR was estimated using the expression $W = aL^b$ (Ricker) [17], where W = total weight (g) and L = total length (mm). The parameters 'a' and 'b' were estimated by linear regression after logarithmic transformation of weight and length data ($\log W = \log a + b \log L$), Where,

W = Weight of fish (g)

L = Length of fish (cm)

a = y Intercept or the initial growth coefficient

b = Slope or the growth coefficient

The statistical significance levels of r^2 and SE for standard error of b ($P < 0.001$) were calculated for all 12 species.

Results

A total of 2015 specimens from 12 fish species belonging to seven families and 11 genera were used for the LWR calculations. The observed maximum total length among all individuals sampled during the study was 1177 mm, which was a specimen of *P. pangasius* having total weight 16 kg and minimum total length 17 mm with total weight

***Corresponding author:** Golam Mortuza M, Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia, Tel: +966-46-97797, Fax: +966-46-78514; E-mail: mmortuza@ksu.edu.sa; mortuzaksu@yahoo.com

Received November 22, 2014; **Accepted** December 21, 2014; **Published** December 24, 2014

Citation: Mortuza MG, Al-Misned FA (2015) Length-Weight Relationships of Twelve Fishes from the River Padma near Rajshahi City, Bangladesh. Fish Aquac J 6: 113. doi: 10.4172/2150-3508.1000113

Copyright: © 2015 Mortuza MG, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

0.13 g of specimen *G. Youssoufi*. The regression coefficients (b) values ranged from 2.813 for *G. youssoufi* to 3.178 for *C. reba*, whereas the 'a' values ranged from 0.1242 to 0.8315 for *L. boga* and *A. coila* respectively.

The calculated 'b' value of the LWR indicated positive allometric growth in *P pangasius*, *A. coila*, *C. nama* *B. lohachata* ($b > 3.00$), but the rest of the species shown negative allometric growth ($b < 3.00$). All LWRs were highly significant ($P < 0.001$), with coefficient of determination (r^2) ranged from 0.838 for *G. youssoufi* to 0.976 for *C. garua*.

Discussion

In the present study, value of 'b' (slope) close to 3 shows that the fish grow isometrically and other values show allometric growth [18]. Most of the estimates for 'b' values obtained in the present work show a similar trend with those of Sani et al. [19] in tributaries of the Yamuna and Ganga, by Sarkar et al. [11] in Ganga basin, Pet et al. [20] in Sri Lankan reservoirs, Ahmed and Saha [2] in Kapatil Lake, Bangladesh, and by Sivakami [21] and Ramakrishniah [22] in the Nagarjunasagar reservoirs. However, in the present study, the higher value of b (> 3) for some species may be due to the dominance of juveniles and an incomplete coverage of the known size range. Differences in the slopes of the length-weight estimates for the same species in different regions can be affected by environmental conditions or developmental state of the fish [23] or, according to Froese [10], by the range lengths used in the length-weight relationships.

The coefficient of determination (r^2) ranged from 0.853 (*L. boga*) to 0.9741 (*C. reba*), nine of 12 regressions presented r^2 values higher than 0.90. All linear regressions were statistically significant ($P < 0.001$). The calculated Standard Error of b (SE) ranged from 0.0068 to 0.0817, thus indicating a tendency towards positive allometry, which is in accordance with the majority of fish species [10].

In conclusion, this study provides an important baseline study on the LWRs of 12 least concern and near threatened fish species from the River Padma near Rajshahi City, Bangladesh [9,24]. These results can be an effective tool for fisheries management and conservation to initiate early management strategies and regulations for conservation of the remaining stocks of the endangered species in the Padma River. This study also provides valuable information for the online database, as well as providing an important baseline for future studies within the Ganga-Brahmaputra basin have almost all been subjected to perturbations of various origins.

Acknowledgements

The authors are thankful to the Deanship of Scientific Research at King Saud University for the provision of funding through the Research Center, College of science, King Saud University.

References

1. Binohlan C, Pauly D (1998) The pop growth table. –In: Foresee R, Pauly D (eds.), FishBase 98: Concepts and design data sources- Manila. ICLARM 124-129.
2. Ahmed KK, Saha SB (1996) Length–weight relationships of major carps in Kapatil Lake, Bangladesh. Naga ICLARM Q 19: 28.
3. Moutopoulos DK, Stergiou KI (2002) Length-weight and length-length relationships of fish species from the Aegean Sea, Greec. Journal of Applied Ichthyology 18: 200-203.
4. Zhulan N, Wu H, Wei J, Zhang X, Ma Z (2013) Length-weight relationship and morphological studies in the Kashgarian loach *Triplophysa yarkandensis* (Day, 1877) from the Tarim River, Tarim River Basin, North-West China. Indian J. Fish 60: 15-19.
5. Mortuza MG, Almisned FA (2013) Length-weight relationships, condition factor and sex-ratio of Nile tilapia, *Oreochromis niloticus* in Wadi Hanifah, Riyadh, Saudi Arabia. Continental Journal of Biological Sciences 6: 1-5.
6. Nile Z, Wu H, Wei J, Zhang X, Ma Z (2013) Length-weight relationship and morphological studies in the Kashgarian loach *riplophysa yarkandensis* (Day, 1877) from the Tarim River, Tarim River Basin, North-West China. Indian Journal of Fisheries 60: 15-19.
7. Teixeira-de Mello F, Iglesias C, Borthagaray AI, Mazzeo N, Vilches J, et al. (2006) Ontogenetic allometric coefficient changes. Implications of diet shift and morphometric attributes in *Hoplias malabaricus* (Bloch) (Characiforme, Erythrinidae). Journal of Fish biology 69: 1770-1778.
8. Teixeira-de Mello F, Eguren G (2008). Prevalence and intensity of black-spot disease in fish community from Canˆ ada del Dragoˆ n stream (Montevideo, Uruguay). Limnetica 27: 251-258.
9. IUCN Bangladesh (2000) Red book of threatened fishes of Bangladesh, IUCN. The world conservation union. xii+116
10. Froese R (2006) Cube law, condition factor and weight–length relationships: history, meta-analysis and recommendations. Journal of Applied Ichthyology 22: 241-253.
11. Sarkar UK, Deepak PK, Negi RS (2009) Length–weight relationship of clown knife fish *Chitala chitala* (Hamilton 1822) from the River Ganga basin India. Journal of Applied Ichthyology 25: 232-233.
12. Muchlisin ZA, Musman M, Siti Azizah MN (2010) Length– weight relationships and condition factors of two threatened fishes, *Rasbora tawarensis* and *Poropuntius tawarensis*, endemic to Lake Laut Tawar, Aceh Province, Indonesia. Journal of Applied Ichthyology 26: 949-953.
13. Jayaram KC (1981) Freshwater fishes of the Indian region. Narendra Pub. House, New Delhi. 551.
14. Rahman AKA (1989) Freshwater fisheries of Bangladesh. Zoological Society of Bangladesh. pp. 364.
15. Talwar PK, Jhingran AG (1991) Inland fishes of India and adjacent countries - Oxford and IBH Publishing Co, New Delhi.
16. Froese R, Pauly D (2011) FishBase 2011. World Wide Web electronic publication.
17. Ricker W E (1973) Linear regressions in fisheries research. Journal of the Fisheries Research Board of Canada 30: 409-434.
18. Andreu-Soler A, Oliva-Paterna FJ, Torralva M (2005) A review of length-weight relationships of fish from the Segura River basin (Siberian Peninsula). Journal of Applied Ichthyology 22: 295-296.
19. Sani R, Gupta BK, Sarkar UK, Pandey A, Dubey VK (2010) Length–weight relationships of 14 Indian freshwater fish species from the Betwa (Yamuna River tributary) and Gomti (Ganga River tributary) rivers. Journal of Applied Ichthyology 26: 456–459.
20. Pet JS, Gevers GJM, Van Densen WLT, Vihverberg J (1996) Management options for a more complete utilization of the biological fish production in Sri Lankan reservoirs. Ecology of Freshwater Fish 4: 1-14.
21. Sivakami S (1987) Length–weight relationship and relative condition factor in *Ompok bimaculatus* (Bloch) from Bhavanisagar reservoir (Tamil Nadu). Indian Journal of Fisheries 34: 202-207.
22. Ramakrishniah M (1988) Age, growth and fisheries of *Mystus aor* (Hamilton) from Nagarjunasagar reservoir. In: The first Indian fisheries forum proceedings, Mohan Joshep (eds.) - Asian Fisheries Society, India Branch, Mangalore: 185-198.
23. Le Cren ED (1951) The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). Journal of Animal Ecology 20: 201-209.
24. Mohsin ABM, Huque SM M, Galib SM, Fahad MFH, Chaki N, et al. (2013) Seasonal Abundance of Fin Fishes in the Padma River at Rajshahi District, Bangladesh. World Journal of Fish and Marine Sciences 5: 680-685.