

Role of Corticosterone and Expression of 11β -Hydroxysteroid Dehydrogenase Type 1 in Liver on Post-Exercise Hyperglycemia in the Obese Diabetic Mouse

Tom L Broderick*

Department of Physiology, Laboratory of Diabetes and Exercise Metabolism, Midwestern University, Glendale, Arizona, USA

Abstract

Exercise training is recommended for the treatment of diabetes mellitus for the benefits on blood glucose control, obesity, and cardiometabolic risk factors. Exercise decreases adipose tissue content and fasting blood glucose, and improves insulin sensitivity. The diet-induced or spontaneous insulin-resistant obese and diabetic rat models to study the effects of exercise training have been widely used because of the favorable outcome on glucose homeostasis. However, the effects of exercise on overall glucose control in the db/db mouse of diabetes remain unclear. The db/db mouse resembles the human condition of type 2 diabetes mellitus and is characterized by hyperleptinemia, hyperglycemia and obesity resulting from a mutation in the leptin receptor gene. The db/db mouse exhibits hypercorticosteronemia, which is also reflective of the human condition. Depending on the exercise training regimen, glucose metabolism is either slightly improved or is further compromised. In this communication, the effects of voluntary and forced treadmill exercise running on glucose control in the db/db mouse are mentioned. In addition, the effects of acute exercise on glucose homeostasis are discussed, along with potential mechanisms explaining the post-exercise hyperglycemia seen in the db/db mouse.

Keywords: Exercise; db/db mouse; 11β -hydroxysteroid; Dehydrogenases; Gluconeogenesis; Leptin; Corticosterone

Introduction

The prevalence of type 2 diabetes mellitus (T2DM) has reached pandemic levels in the last decade in developed and developing countries as a result of increased obesity rates. The association between T2DM and cardiovascular diseases is well-established, and diabetics are at increased risk of myocardial infarct, heart failure and sudden death compared to no diabetics [1]. Obese diabetics also present with dysregulation in white adipose tissue metabolism, such as aberrant production of leptin [2]. Leptin is an adipocyte-derived peptide that circulates in the plasma at levels proportional to amount of adipose tissue content. This peptide has a key role in the regulation of energy intake and satiety through central mechanisms as well as peripheral actions on energy expenditure. Excessive secretion of this adipokine from adipose tissue, as expected, can potentially lead to leptin resistance or hyperleptinemia in T2DM. Common sequelae of hyperleptinemia include failure to regulate food intake, aberrations in energy homeostasis and fat distribution, worsening of the diabetic state and the development of other cardiovascular risks factors such as lipid disorders and hypertension [3]. The risk of cardiovascular disease and associated cardiometabolic burdens can be largely prevented by implementation of effective non-pharmacological approaches, such as increasing the level of physical activity [4]. Advances in our understanding of the role of exercise, from both a mechanistic aspect and application to lifestyle modification in patients with T2DM is clearly dependent on the use of relevant animal models [5]. In this review, the use of the hyperleptinemic db/db mouse model of T2DM in exercise-based studies is discussed as well as the potential involvement of endogenous glucocorticoids and the cortisol-cortisone shunt in the regulation of glucose metabolism in this model.

Benefits of Exercise in T2DM

Regular physical activity is recommended for the treatment of T2DM, obesity, and related metabolic disorders. Exercise performed on a regular basis increases muscle content, improves blood glucose control,

lowers plasma lipids and blood pressure, and increases insulin sensitivity [6,7]. Impaired mitochondrial dysfunction and energy metabolism can be improved, leading to enhanced exercise capacity for the diabetic patient [8,9]. Deflections in postprandial hyperglycemia and impaired fasting glucose, both risk factors in the development of cardiometabolic disorders can be prevented with exercise training [10]. Controlling blood glucose levels by exercise delays the progression of long-term complications of symptomatic cardiovascular disease and protects against the development of type 2 diabetes independent of obesity, age, history of hypertension and parental history of diabetes [6].

The db/db Mouse Model

Rodents because of their size, easy handling, affordable costs, and rapid maturation rates remain the obvious choice as model to examine the role of exercise on the diabetic state. Further, the variety and availability of inbred strains make them ideal for study because of the genetic uniformity as well as the predictable phenotypes that gradually develop over a short period of time. Various models and strains of diabetic and obese rodents are available for exercise-based studies, including the diet-induced model (DIO), low-dose streptozotocin (STZ) model, Zucker diabetic fatty and spontaneous insulin resistant models of type 2 diabetes [5]. These commonly used models are responsive to the stimulus of exercise and have provided valuable information on

*Corresponding author: Tom L Broderick, Midwestern University, Department of Physiology, Laboratory of Diabetes and Exercise Metabolism, 19555 North 59th Avenue, Glendale, AZ, 85308, USA, Tel: 6235723664; Fax: 6235723673; E-mail: tbrode@midwestern.edu

Received January 04, 2017; Accepted January 18, 2017; Published January 25, 2017

Citation: Broderick TL (2017) Role of Corticosterone and Expression of 11β -Hydroxysteroid Dehydrogenase Type 1 in Liver on Post-Exercise Hyperglycemia in the Obese Diabetic Mouse. Adv Tech Biol Med 5: 200. doi: [10.4172/2379-1764.1000200](https://doi.org/10.4172/2379-1764.1000200)

Copyright: © 2017 Broderick TL. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

the consequences of exercise training on cardiovascular and metabolic outcome in the diabetic state [11-15]. For these reasons, the use of these particular models is justified in exercise studies and for translational-based application in the treatment of humans with this metabolic disorder. However, in recent years, the use of the db/db mouse model of diabetes to investigate the impact of exercise training has gained popularity. Yet, since the early studies by Coleman and colleagues in the mid-1960s and 1970s characterizing the metabolic and endocrine profile of the db/db mouse [16-18], this model is still underrepresented compared to the other rodent models used in exercise studies. The db/db mouse is a model of T2DM and is considered a close counterpart to the human condition in terms of presenting phenotype, metabolic and cardiovascular perturbations. Diabetic-related complications in the db/db mouse result from the long-lasting effects of a mutation in the leptin receptor gene (*LepR^{db/db}*) disrupting post-receptor signaling [19]. This mutation affects hypothalamic responses to food intake, leading to hyperphagia and the gradual development of obesity. By 2-3 weeks of age, db/db mice show signs of insulin resistance and from 4 weeks of age are hyperglycemic. The db/db mouse also exhibits hypercortisolemia (hypercortisolemia in humans) as a result of hypertrophy of the adrenal cortex [18], a well-established risk factor associated with the development of insulin resistance and hyperglycemia [20,21]. Another hallmark feature of the db/db mouse, as expected, is hyperleptinemia from defective leptin signaling. This rarely is the underlying cause of diabetes but nonetheless observed in the obese diabetic patient [22].

The db/db Mouse and Exercise Training

Treadmill running remains the preferred exercise paradigm for training in rodents. The duration and intensity of exercise can be easily manipulated keeping the overall training volume identical in experimental groups. Oxygen consumption of mice and hence intensity of effort can be estimated simply based on treadmill belt speed without the need for direct measurement of oxygen use in each mouse [23], another reason why this form of exercise is considered the paragon of excellence in exercise physiology. Treadmill running comes with its drawbacks however, including motivation, innate low ambulatory function for some mice strains, and increased stress. Constant monitoring is also needed to avoid injury of mice with low ambulatory activity.

Assessing the effects of exercise training on glucose control in the db/db mouse was first addressed using swim training as regimen [24]. Combination of low and moderate swim training intensity over a period of 4 weeks decreased blood glucose concentrations but db/db mice remained hyperglycemic, hyperinsulinemic, and insulin resistant to an oral glucose challenge. Diabetic mice exhibited the typical obesity of this model after swim training despite a slight decrease in weight gain. However, when db/db mice were treated with common antihyperglycemic agents, an improvement in blood glucose and insulin concentrations was observed. Regardless of these negligible effects of blood glucose regulation with exercise alone, more recent studies have shown that mild-to-moderate exercise training is of benefit on vascular function and lipid metabolism. In fact, 5 weeks of voluntary wheel running improved vascular endothelial function and decreased common lipid levels, but failed to reduce body weight and correct plasma glucose concentrations [25]. Could these negligible effects of training on glycemic control and body weight be the result of a low training stimulus or work volume and can the diabetic state be improved simply by increasing the intensity of the training?

To address this, we exposed db/db mice to voluntary wheel running and forced treadmill running for a period of 12 weeks [26]. The training regimen for mice assigned to the treadmill running group was designed

to elicit a greater stimulus based on intensity and training volume (3-fold higher than voluntary running), and with expectations that this would decrease both the obesity and hyperglycemia in db/db mice. After 12 weeks of voluntary wheel running, body weight dropped by ~5% while fasting blood glucose levels decreased by ~21% compared to sedentary db/db mice (Figures 1 and 2). Forced treadmill running however, did not overcome the disturbances associated with the db/db mouse. In fact, both body weight and blood glucose were increased by ~8-10% and mice remained insulin resistant (Figure 3). When the data was expressed as percent change from the onset of exercise training, the greatest changes in body weight and fasting blood glucose levels were observed in db/db mice subjected to treadmill running (Table 1). Based on these observations, increasing the training volume by altering intensity does not always correlate with greater weight loss and improved metabolic control in the db/db mouse. Results from earlier studies in obesity models have suggested that an exaggerated stress response associated with forced treadmill exercise could explain these unfavorable changes in the db/db mouse. Daily short-term exposure to stress can induce adrenal gland hypertrophy and disruption




Figure 1: Changes in body weight in lean and diabetic mice with exercise training. Exercise training was performed over a period of 12 weeks.

CN: Lean Control; DS: Diabetic Sedentary; DV: Diabetic Voluntary Running Mice; DT: Diabetic Treadmill Running Mice. Values are reported as mean \pm SEM for 12 to 18 mice per group. *: Compared to initial week 0; †: Compared to CN mice; ‡: Compared to DS mice





Figure 2: Changes in fasting blood glucose concentrations in lean and diabetic mice with exercise training. Exercise training was performed over a period of 12 weeks.

CN: Lean Control; DS: Diabetic Sedentary; DV: Diabetic Voluntary Running Mice; DT: Diabetic Treadmill Running Mice. Values are reported as mean \pm SEM for 12 to 18 mice per group. *: Compared to initial week 0; †: Compared to CN mice; ‡: Compared to DS mice



| Group | Percent change in body weight | Percent change in blood glucose |
|--------------------|-------------------------------|---------------------------------|
| Lean control | 40 \pm 1 | 19 \pm 1 |
| Diabetic sedentary | 45 \pm 2 | 187 \pm 6* |
| Diabetic voluntary | 61 \pm 4 [†] | 185 \pm 21* |
| Diabetic treadmill | 83 \pm 2 ^{††} | 327 \pm 13 ^{††} |

Values are reported as mean \pm SEM from 12-18 mice in each group and expressed as percent change from the week 1. C57BL/KsJ-lept^{ob}-lept^{ob} mice were used [26]. *: Compared to lean control. [†]: Compared to sedentary diabetic mice. ^{††}: Compared to diabetic voluntary runners

Table 1: Percent changes in body weight and blood glucose levels after 12 weeks of exercise training in control and diabetic mice.

of the hypothalamus pituitary-adrenal axis (HPA), further increasing the susceptibility of the obese diabetic mouse to metabolic aberrations [27-29].

The Corticosterone Response to Exercise in the db/db Mouse

Supporting the link between the stress imposed by daily forced running and metabolic perturbations are the findings that treadmill exposure is associated with increased corticosterone secretion, elevated activity of the renin-angiotensin system, and insulin resistance [20,27-29]. Hedonic feeding behavior is also reported after the cessation of exercise [30], disrupting central signaling mechanisms in the regulation of food intake which are already affected by leptin resistance. Enhanced endogenous glucocorticoid synthesis from hypertrophy of the adrenal gland is an inherent feature of the db/db mouse [16-18] and our earlier studies highlight the importance of this feature on dysregulated glucose control during and after acute exercise as well as in response to chronic training (Table 2). Acute treadmill exercise is associated with greater corticosterone synthesis and lipolysis in db/db mice when compared to heterozygote lean controls derived from the same strain, leading to post-exercise hyperglycemia [31]. The increase in corticosterone production in the db/db mouse after exercise was not caused by activation of the HPA axis because release of both corticotropin-releasing hormone (CRH) and adrenocorticotrophic hormone (ACTH) levels were reciprocally decreased, indicating no disruption in feedback regulation. Corticosterone synthesis and spillover in urine are elevated in db/db mice after treadmill running compared to voluntary running, suggesting that db/db mice exhibit a greater stress response to treadmill running [32] (Table 3). A sympathetic nervous system response may

further contribute to increased blood glucose levels in the db/db mouse as normetanephrine content was elevated in urine of db/db mice, suggesting involvement of hepatic glycogenolysis in blood glucose regulation after treadmill training [32]. Interestingly, patients with T2DM presenting with hyperglycemia prior to exercise exhibit post-exercise hyperglycemia and hyperinsulinemia, indicating a worsening of metabolic control. Excessive norepinephrine secretion is also reported in both hypertensive and non-hypertensive T2DM after exercise [33-36]. Clearly, studies examining the role of sympathetic tone and function on glucose homeostasis in the db/db mouse during exercise are warranted and intriguingly also because of recent evidence indicating a beneficial effect of leptin treatment in reversing the development of autonomic nervous system neuropathy in older db/db mice [37-39].

Exercise and Hepatic Gluconeogenesis in the db/db mouse

During acute exercise, normoglycemia is maintained by a precise balance between hepatic glucose production from gluconeogenesis and peripheral glucose utilization. When hepatic glucose production is greater than the peripheral use of glucose, this results in hyperglycemia. The hyperglycemia observed in the obese Zucker rat and Zucker diabetic fatty rat is related, in part, to elevated rates of hepatic gluconeogenesis, which can be reversed with chronic exercise training [40,41]. In these rat models, the expression of major hepatic glucose production enzymes phosphoenolpyruvate carboxykinase (PEPCK) and glucose-6-phosphatase (G6Pase) is increased relative to lean non-diabetic littermates, and treadmill exercise training reduces their expression, improving the hyperglycemic state. The effects of both acute and chronic exercise training on the expression of gluconeogenic enzymes in the db/db mouse have received little attention to date. However, recent evidence from our laboratory suggests that the post-exercise elevation in plasma glucose levels in the db/db mouse may be caused by increased expression of PEPCK (Table 4). From a mechanistic aspect, activation of the glucocorticoid receptor (GR) by corticosterone could stimulate the expression of PEPCK, resulting in

| Parameter | Lean mice | Lean+exercise | Db/db mice | Db/db+exercise |
|------------------------|---------------|----------------|----------------------------|-----------------------------|
| Glucose (mg/dl) | 165 \pm 20 | 187 \pm 20 | 420 \pm 19 ^{††} | 571 \pm 22 ^{†††} |
| NEFA (mmol/L) | 0.6 \pm 0.1 | 0.63 \pm 0.1 | 0.68 \pm 0.1 | 0.94 \pm 0.1* |
| Corticosterone (ng/ml) | 193 \pm 50 | 432 \pm 48* | 470 \pm 46* | 560 \pm 54* |

Blood was collected 30 minutes before exercise and immediately after exercise. 7-8 week old db/db mice were used and lean littermates served as controls. Exercise was performed on a treadmill for 30 min at 12 m/min. Values are reported as mean \pm SEM for 10 mice in each group. NEFA: Non-Esterified Fatty Acids; *: Compared to lean mice; [†]: Compared to lean exercise mice; ^{††}: Compared to db/db mice

Table 2: Effects of acute exercise on plasma glucose, non-esterified fatty acids and corticosterone levels in control and db/db mice.

| | Lean control | Lean runners | Diabetic control | Diabetic runners |
|-------------------------|-----------------|-----------------|-------------------------------|-------------------------------|
| Corticosterone (pg/ml) | | | | |
| -Voluntary running | 99 \pm 24 | 264 \pm 65 | 319 \pm 49* | 241 \pm 96 |
| -Treadmill running | 133 \pm 5 | 171 \pm 19 | 269 \pm 26 [†] | 820 \pm 226 ^{††} |
| Normetanephrine (ng/ml) | | | | |
| -Voluntary running | 5.00 \pm 0.09 | 5.28 \pm 0.05 | 2.96 \pm 1.50 ^{††} | 3.12 \pm 0.35 ^{††} |
| -Treadmill running | 5.19 \pm 0.07 | 5.07 \pm 0.15 | 2.45 \pm 0.80 ^{††} | 3.65 \pm 0.37 ^{††} |

Values are reported as mean \pm SEM for 6 mice in each group. Five-week-old db/db mice and lean littermates were trained either on a treadmill or in a wheel system for a period of 5 weeks [32]. Urine was collected using metabolic cages over a 24h period. *: Compared to lean control mice; [†]: Compared to lean runners; ^{††}: Compared to db/db control mice

Table 3: Effects of voluntary wheel and treadmill running on urine corticosterone and normetanephrine levels in lean and db/db mice.

an increase in glucose production. In liver of db/db mice, in which the GR is abundantly expressed and endogenous corticosterone synthesis is increased [18,42], upregulation of the GR and PEPCK contributing to post-exercise hyperglycemia is an attractive mechanism that merits further investigation (Table 2).

Hepatic 11 β -hydroxysteroid Dehydrogenase and Exercise in the db/db Mouse



It is proposed that altered expression of the central and contrasting

| Protein | Lean mice | Lean+exercise | Db/db mice | Db/db+exercise |
|---------|-----------|---------------|------------|----------------|
| GR | 1.0 | +1.77 | +2.89* | +3.21** |
| PEPCK | 1.0 | +1.33 | +1.45 | +1.75* |

To highlight the changes induced by exercise or diabetes, values are presented as delta change in protein expression relative to control. For these data [31], liver from lean control littermates were used, which have one mutant and one normal copy of the leptin gene (db/*). GR: Glucocorticoid Receptor; PEPCK: Phosphoenolpyruvate Carboxykinase. Protein expression was normalized to actin. Exercise was performed on a treadmill for 30 min at 12 m/min in 8 week old db/db mice. Mice were sacrificed 8 hours after exercise for protein expression. Values are reported using 6-10 mice in each group. *: Compared to lean mice; **: Compared to lean exercise mice

Table 4: Changes in protein expression of glucocorticoid receptor and phosphoenolpyruvate carboxykinase in liver of db/db mice.

11 β -hydroxysteroid dehydrogenase (HSD) isoenzymes of the corticosterone-11-dehydrocorticosterone shunt may be contributing to post-exercise hyperglycemia in the db/db mouse. Corticosterone can bind the mineralocorticoid receptor and cause inappropriate activation of mineralocorticoid pathways. To prevent these effects, 11 β -HSD type 2 converts active corticosterone into inert 11-keto forms (11-dehydrocorticosterone in rodents, cortisone in humans) in the kidney (Figure 4). The inert glucocorticoid is then transported to the liver where 11 β -HSD type 1 reconverts the inactive form into active corticosterone. 11 β -HSD type 1 is expressed in tissues such as skin, muscle, and adipose tissue, and dysregulation of 11 β -HSD type 1 in liver and adipose tissue in obesity increases the intracellular levels of corticosterone and contributes to the plasma glucocorticoid pool [43]. The effects of acute exercise on the expression of 11 β -HSD type 1 in liver has been examined as a possible mechanism for the post-exercise excursion in blood glucose levels seen in db/db mice [31]. Our latest work indicates that increased expression of 11 β -HSD type 1 stimulates corticosterone synthesis, leading to increased protein expression of gluconeogenic enzymes and hepatic glucose production after exercise in the db/db mouse (Figure 5). This mechanism is consistent with the observations that overexpression of 11 β -HSD1 is associated with visceral adipose tissue accumulation, hepatic gluconeogenesis, hyperglycemia,



and insulin resistance in obesity [43-45]. Considering that the db/db mouse is leptin-resistant, the role of leptin in the regulation of 11 β -HSD type 1 synthesis could lead to increased understanding of the role of hypercorticotestosterone on hepatic glucose production. Indeed, leptin treatment to both STZ diabetic rats and in the genetic model of spontaneous T1DM (Bio Breeding, BB) was effective in correcting ACTH and corticosterone levels and restoring glucose levels by normalizing hepatic gluconeogenesis [46]. Enhanced CRH and ACTH production with resulting hypercortisolemia has been reported in obese women with abdominal body fat distribution [47] as well as in the obese Zucker (fa/fa) rat [48]. Chronic leptin replacement in the obese ob/ob mouse [49], which is leptin deficient, corrects the hypercorticotestosterone by establishing negative feedback to the HPA axis highlighting the importance of adipose tissue in corticosterone secretion [50]. As expected, with leptin resistance, the HPA axis feedback loop is disrupted in the db/db mouse.

Conclusion and Future Directions

Inactivity contributes to the development of obesity and diabetes, and exercise is recommended as a non-pharmacological approach for the treatment of these metabolic disorders. Regular exercise can prevent the progression of diabetes-related complications and improve overall cardiovascular health. Based on the results of exercise-based studies glucose control is not achieved in the obese diabetic db/db mouse [25,26,51-54]. This should not raise any doubts regarding the suitability of this model for training studies, because benefits on other cardiovascular risk factors have been reported. When comparing voluntary exercise as a training regimen to forced treadmill running, however, a slight improvement in glucose control occurs when mice run on their own volition. Voluntary running, by decreasing the production stress-related hormones [55], could explain this effect on glucose control although the effects of chronic voluntary running on these markers are unclear. On the other hand, the db/db mouse exhibits enhanced endogenous glucocorticoid synthesis at rest, and further synthesis occurs during and after exercise, leading to post-exercise hyperglycemia by gluconeogenic mechanisms. Further contributing to hyperglycemia in the db/db mouse is the observation that 11 β -HSD type 1 expression of the corticosterone-11-dehydrocorticosterone shunt in liver is upregulated during exercise. It is important to note that the role of kidney and expression of inactivating isoenzymes present in other tissues of this shunt has not been demonstrated in exercise-based studies using the db/db. Lastly, the effects of exercise on peripheral glucose metabolism and hepatic glucose production are also complicated by the fact that the db/db mouse is hyperleptinemic, which may affect running behavior. Evidence for a role of leptin in ambulatory behavior is based on the observation that leptin treatment to leptin-deficient ob/ob mice stimulates energy expenditure and wheel running behavior [56,57], suggesting that leptin is an important regulator of activity in mouse models of defective leptin signaling.

Acknowledgement

The author is grateful to the Midwestern University Office of Research and Sponsored Programs and the Diabetes and Action Research and Education Foundation for their continued support.

References

1. Kannel WB, McGee DL (1979) Diabetes and cardiovascular disease. The Framingham study. *JAMA* 241: 2035-2038.
2. Farr OM, Gavrieli A, Mantzoros CS (2015) Leptin applications in 2015: What have we learned about leptin and obesity? *Curr Opin Endocrinol Diabetes Obes* 22: 353-359.
3. Enriori PJ, Evans AE, Sinnayah P, Cowley MA (2006) Leptin resistance and obesity. *Obesity (Silver Spring)* 5: 254S-258S.
4. van der Berg JD, Stehouwer CD, Bosma H, van der Velde JH, Willems PJ, et al. (2016) Associations of total amount and patterns of sedentary behaviour with type 2 diabetes and the metabolic syndrome: The Maastricht study. *Diabetologia* 59: 709-718.
5. Ghosh S, Golbidi S, Werner I, Verchere B, Laher I (2010) Selecting exercise regimens and strains to modify obesity and diabetes in rodents: An overview. *Clin Sci* 119: 57-74.
6. Boule NG, Haddad E, Kenny GP, Wells GA, Sigal RJ (2001) Effects of exercise on glycemic control and body mass in type 2 diabetes mellitus: A meta-analysis of controlled clinical trials. *JAMA* 286: 1218-1227.
7. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, et al. (2002) Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346: 393-403.
8. Bajpeyi S, Pasarica M, Moro C, Conley K, Jubrias S, et al. (2011) Skeletal muscle mitochondrial capacity and insulin resistance in type 2 diabetes. *J Clin Endocrinol Metab* 96: 1160-1168.
9. Yokota T, Kinugawa S, Okita K, Hirabayashi K, Suga T, et al. (2011) Lower aerobic capacity was associated with abnormal intramuscular energetics in patients with metabolic syndrome. *Hypertens Res* 34: 1029-1034.
10. Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, et al. (2001) Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 344: 1343-1350.
11. Cortez MY, Torgan CE, Brozinick JT Jr, Ivy JL (1991) Insulin resistance of obese Zucker rats exercise trained at two different intensities. *Am J Physiol* 261: E613-E619.
12. Banks EA, Brozinick JT Jr, Yaspelkis BB, Kang HY, Ivy JL (1992) Muscle glucose transport, GLUT-4 content and degree of exercise training in obese Zucker rats. *Am J Physiol* 263: E1010-1015.
13. Brozinick JT Jr, Etgen GJ Jr, Yaspelkis BB 3rd, Kang HY, Ivy JL (1993) Effects of exercise training on muscle GLUT-4 protein content and translocation in obese Zucker rats. *Am J Physiol* 265: E419-427.
14. Podolin DA, Wei Y, Pagliassotti MJ (1999) Effects of a high-fat diet and voluntary wheel running on gluconeogenesis and lipolysis in rats. *J Appl Physiol* 86: 1374-1380.
15. Davidoff AJ, Mason MM, Davidson MB, Carmody MW, Hintz KK, et al. (2004) Sucrose-induced cardiomyocyte dysfunction is both preventable and reversible with clinically relevant treatments. *Am J Physiol Endocrinol Metab* 286: E718-724.
16. Naeser P (1976) Adrenal function in the diabetic mutant mouse (gene symbol dbm). *Acta Physiol Scand* 98: 395-399.
17. Hummel KP, Dickie MM, Coleman DL (1966) Diabetes, a new mutation in the mouse. *Science* 153: 1127-1128.
18. Coleman DL, Burkart DL (1977) Plasma corticosterone concentrations in diabetic (db) mice. *Diabetologia* 13: 25-26.
19. Leibel RL, Chung WK, Chua SC Jr (1997) The molecular genetics of rodent single gene obesities. *J Biol Chem* 272: 31937-31940.
20. Rosmond R, Dallman MF, Bjorntorp P (1998) Stress-related cortisol secretion in men: Relationships with abdominal obesity and endocrine, metabolic and hemodynamic abnormalities. *J Clin Endocrinol Metab* 83: 1853-1859.
21. Anagnostis P, Athyros VG, Tziomalos K, Karagiannis A, Mikhailidis DP (2009) Clinical review: The pathogenetic role of cortisol in the metabolic syndrome: A hypothesis. *J Clin Endocrinol Metab* 94: 2692-2701.
22. Bray GA, York DA (1997) Clinical review 90: Leptin and clinical medicine: A new piece in the puzzle of obesity. *J Clin Endocrinol Metab* 82: 2771-2776.
23. Høydal MA, Wisløff U, Kemi OJ, Ellingsen O (2007) Running speed and maximal oxygen uptake in rats and mice: practical implications for exercise training. *Eur J Cardiovasc Prev Rehabil* 14: 753-760.
24. Tang T, Reed MJ (2001) Exercise adds to metformin and acarbose efficacy in db/db mice. *Metabolism* 50: 1049-1053.
25. Esser KA, Su W, Matveev S, Wong V, Zeng L, et al. (2007) Voluntary wheel running ameliorates vascular smooth muscle hyper-contractility in type 2 diabetic db/db mice. *Appl Physiol Nutr Metab* 32: 711-720.
26. Sennott J, Morrissey J, Standley PR, Broderick TL (2008) Treadmill exercise

- training fails to restore defects in glucose, insulin and muscle glut4 content in the db/db mouse model of diabetes. *Pathophysiology* 15: 173-179.
27. Moraska A, Deak T, Spencer RL, Roth D, Fleshner M (2000) Treadmill running produces both positive and negative physiological adaptations in Sprague-Dawley rats. *Am J Physiol Regul Integr Comp Physiol* 279: R1321-1329.
28. Tharp GD (1975) The role of glucocorticoids in exercise. *Med Sci Sports* 7: 6-11.
29. Rosmond R (2003) Stress induced disturbances of the HPA axis: A pathway to Type 2 diabetes? *Med Sci Monit* 9: RA35-39.
30. Applegate EA, Stern JS (1987) Exercise termination effects on food intake, plasma insulin and adipose lipoprotein lipase activity in the Osborne-Mendel Rat. *Metabolism* 36: 709-714.
31. Brust KB, Corbell KA, Al-Nakkash L, Babu JR, Broderick TL (2014) Expression of gluconeogenic enzymes and 11 β -hydroxysteroid dehydrogenase type 1 in liver of diabetic mice after acute exercise. *Diabetes Metab Syndr Obes* 7: 495-504.
32. Parrott CR, Ghosh P, Tedeschi J, Gunasekara G, Broderick TL (2011) Urinary corticosterone and normetanephrine levels after voluntary and forced treadmill running in the db/db mouse. *J Diabetes Mellitus* 40: 71-78.
33. Kjaer M, Hollenbeck CB, Frey-Hewitt B, Galbo H, Haskell W, et al. (1990) Glucoregulation and hormonal responses to maximal exercise in non-insulin-dependent diabetes. *J Appl Physiol* 68: 2067-2074.
34. Raber W, Raffenber W, Waldhäusl W, Gasic S, Roden M (2003) Exercise induces excessive normetanephrine responses in hypertensive diabetic patients. *Eur J Clin Invest* 33: 480-487.
35. Hübinger A, Franzen A, Gries FA (1987) Hormonal and metabolic response to physical exercise in hyperinsulinemic and non-hyperinsulinemic type 2 diabetics. *Diabetes Res* 4: 57-61.
36. Vettor R, Macor C, Rossi E, Piemonte G, Federspil G (1997) Impaired counter regulatory hormonal and metabolic response to exhaustive exercise in obese subjects. *Acta Diabetol* 34: 61-66.
37. Robertson DM, Sima AA (1980) Diabetic neuropathy in the mutant mouse [C57BL/ks(db/db)]: A morphometric study. *Diabetes* 29: 60-67.
38. Valerio A, Ghisi V, Dossena M, Tonello C, Giordano A, et al. (2006) Leptin increases axonal growth cone size in developing mouse cortical neurons by convergent signals inactivating glycogen synthase kinase-3 β . *Journal of Biological Chemistry* 281: 12950-12958.
39. Drel VR, Mashtalir N, Ilnytska O, Shin J, Li F, et al. (2006) The leptin-deficient (ob/ob) mouse: A new animal model of peripheral neuropathy of type 2 diabetes and obesity. *Diabetes* 55: 3335-3343.
40. Chang SP, Chen YH, Chang WC, Liu IM, Cheng JT (2006) Merit of physical exercise to reverse the higher gene expression of hepatic phosphoenolpyruvate carboxykinase in obese Zucker rats. *Life Sciences* 79: 240-246.
41. Campbell JE, Király MA, Atkinson DJ, D'souza AM, Vranic M, et al. (2010) Regular exercise prevents the development of hyperglucocorticoidemia via adaptations in the brain and adrenal glands in male Zucker diabetic fatty rats. *Am J Physiol Regul Integr Comp Physiol* 299: R168-R176.
42. Perelló M, Moreno G, Gaillard RC, Spinedi E (2004) Glucocorticoid-dependency of increased adiposity in a model of hypothalamic obesity. *Neurology and Endocrinology Letters* 25: 119-126.
43. Liu Y, Nakagawa Y, Wang Y, Sakurai R, Tripathi PV, et al. (2005) Increased glucocorticoid receptor and 11 β -hydroxysteroid dehydrogenase type 1 expression in hepatocytes may contribute to the phenotype of type 2 diabetes in db/db mice. *Diabetes* 54: 32-40.
44. Whorwood CB, Donovan SJ, Flanagan D, Phillips DI, Byrne CD (2002) Increased glucocorticoid receptor expression in human skeletal muscle cells may contribute to the pathogenesis of metabolic syndrome. *Diabetes* 51: 1066-1075.
45. Kargi AY, Iacobellis G (2014) Adipose tissue and adrenal glands: Novel pathophysiological mechanisms and clinical applications. *Int J Endocrinol* 2014: 614074.
46. Perry RJ, Zhang XM, Zhang D, Kumashiro N, Camporez JPG, et al. (2014) Leptin reverses diabetes by suppression of the hypothalamic-pituitary-adrenal axis. *Nat Med* 20: 759-763.
47. Pasquali R, Cantobelli S, Casimirri F, Capelli M, Bortoluzzi L, et al. (1993) The hypothalamic-pituitary-adrenal axis in obese women with different patterns of body fat distribution. *J Clin Endocrinol Metab* 77: 341-346.
48. Plotsky PM, Thiruvikraman KV, Watts AG, Hauger RL (1992) Hypothalamic-pituitary-adrenal axis function in the Zucker obese rat. *Endocrinology* 130: 1931-1941.
49. Stephens TW, Basinski M, Bristow PK, Bue-Valleskey JM, Burgett SG, et al. (1995) The role of neuropeptide Y in the anti-obesity action of the obese gene product. *Nature* 377: 530-532.
50. Heiman ML, Ahima RS, Craft LS, Schoner B, Stephens TW, et al. (1997) Leptin inhibition of the hypothalamic-pituitary-adrenal axis in response to stress. *Endocrinology* 138: 3859-3863.
51. Moien-Afsahri F, Ghosh S, Elmi S, Rahman MM, Sallam N, et al. (2008) Exercise restores endothelial function independently of weight loss or hyperglycemic status in db/db mice. *Diabetologia* 51: 1327-1337.
52. Moien-Afshari F, Ghosh S, Elmi S, Khazaei M, Rahman MM, et al. (2008) Exercise restores coronary vascular function independent of myogenic tone or hyperglycemic status in db/db mice. *Am J Physiol Heart Circ Physiol* 295: H1470-1480.
53. Broderick TL, Parrott CR, Wang D, Jankowski M, Gutkowska J (2012) Expression of cardiac GATA4 and downstream genes after exercise training in the db/db mouse. *Pathophysiology* 19: 193-203.
54. Gutkowska J, Broderick TL, Bogdan D, Wang D, Lavoie JM, et al. (2009) Down regulation of oxytocin and natriuretic peptides in diabetes: Possible implications in cardiomyopathy. *J Physiol* 587: 4725-4736.
55. Wong DL (2006) Epinephrine biosynthesis: Hormonal and neural control during stress. *Cell Mol Neurobiol* 26: 891-900.
56. Morton GJ, Kaiyala KJ, Fisher JD, Ogimoto K, Schwartz MW, et al. (2011) Identification of a physiological role for leptin in the regulation of ambulatory activity and wheel running in mice. *Am J Physiol Endocrinol Metab* 300: E392-E401.
57. Park S, Jang JS, Jun DW, Hong SM (2005) Exercise enhances insulin and leptin signaling in the cerebral cortex and hypothalamus during dexamethazone-induced stress in diabetic rats. *Neuroendocrinol* 82: 282-293.