

# The Effect of Mixed Plantation on the Stand Yield and Soil Attributes of *Eucalyptus globulus* and *Acacia decurrens* in North Shewa Zone, Ethiopia

Melese Bekele Nigussie<sup>1\*</sup>, Lemma Habeteyohannes<sup>1</sup>, Getabalew Teshome<sup>1</sup>, Damtew Ababu<sup>1</sup>, Abeje Tedila<sup>1</sup>, Reta Eshetu<sup>1</sup>, Mesafint Minale<sup>1</sup> Hailemariyam Fisha<sup>1</sup> and Haile Shiferaw<sup>2</sup>

<sup>1</sup>Debrbirhane Agricultural research Center, Debre Birhan, Ethiopia, <sup>2</sup>Environment and Forest Research Center, Bahir Dar, Ethiopia

## ABSTRACT

Eucalyptus plantation has been playing a vital role in narrowing the gap between the supply and demand for wood and wood products in the region. Though eucalyptus is fast-growing and has a better survival rate compared to other exotic species, there is a claim that Eucalyptus species over exploiting the nutrients and water. This consequently led to land degradation and soil fertility decline. Hence, it is important to develop mixed plantations to minimize the gap of monocultures. This study aimed to select *Acacia decurrens*-*Eucalyptus camaldulensis* mixed plantation to obtain better woody yield and improve site productivity than sole plantation. *A. decurrens* and *E. camaldulensis* seedlings were planted with the proportions of: 0%:100%, 100%:0%, 25%:75%, 75%:25% and 50%:50%. The experiment was arranged in a randomized complete block design with three replications. Results showed that the tree height and total volume not varied significantly between the mixed and pure stand plantations. Soil nitrogen varies significantly between the treatments. The plantation with 50% *E. camaldulensis* and 50% *A. decurrens* proportion had higher total nitrogen, organic carbon and organic matter in the soil. In line with the experiment, the farmers confirmed that the 50% *E. camaldulensis* with 50% *A. decurrens* plantations had relatively a better economic benefits and soil improvements. Further investigation is required to know the contribution of each species in soil nutrient addition. Investigation of the mixed plantation of exotic and indigenous is required to optimize economic and environmental benefits.

**Keywords:** *E. camaldulensis*; *A. decurrens*; Mixtures; Growth; Yield

## INTRODUCTION

Eucalyptus is a genus that has more than 500 species. This genus is native to Australia, the Malaysian region, and the Philippines but it is planted in a different part of the world since the 1890s [1]. Eucalyptus is adaptable to tropical, arid and temperate regions of the world. It is one of the most widely planted trees in the world which covers around 20 million hectares of land. It is planted for paper pulp, fuel wood, timber, amenity plantings and land rehabilitation [2]. Plantations of eucalypts (species of *Eucalyptus* and *Corymbia*), particularly in the tropics and Southern Hemisphere, have expanded dramatically during the last 100 years [3].

Since the introduction, 55 and more species of eucalyptus grown in Ethiopia, of which between five and ten are widely planted [1]. *E. globulus* performs well in terms of growth and survival rate in highlands and *E. camaldulensis* the second most common eucalyptus species of ten grown in lower altitudes [4].

Increasing benefit from eucalyptus is an incentive for farmers to

expand eucalyptus plantations at the expense of another land uses Engda TA, et al. [4], and also as a result of the availability of high demands for eucalyptus products the trend of expansion of eucalyptus plantation on crop and communal lands have been increasing. The market price for eucalyptus poles has grown to be 15 times greater over the last two decades [5-10].

Though fast-growing and have a better survival rate in degraded lands, there is a concern that mono-cultivation of eucalyptus has caused negative consequence in watershed management, soil and water conservation, wildlife habitat, and recreational values. Hence, to balance this, introducing an improved system could be an appropriate solution [11-14].

The mixed plantation is assumed to address more comprehensively the community needs of fuel and construction wood, nutrient cycle and the likes. Besides, different trees or shrub species are not equally affected by adverse environmental factors including drought, disease, and pests, and hence a mixed plantation avoids the risk of total failure or loss [6] studied that mixed stands of

**Correspondence to:** Nigussie MB, Debrbirhane Agricultural Research Center, Debrbirhane, Ethiopia. Tel: +251-0920718821; E-mail: mel21bekele@gmail.com

**Received:** August 04, 2021, **Accepted:** August 23, 2021, **Published:** August 30, 2021

**Citation:** Nigussie MB, Habeteyohannes L, Teshome G, Ababu D, Tedila A, Eshetu R, et al. (2021) The Effect of Mixed Plantation on the Stand Yield and Soil Attributes of *Eucalyptus globulus* and *Acacia decurrens* in North Shewa Zone, Ethiopia. J Forest Res 10:282.

**Copyright:** © 2021 Nigussie MB, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

species of similar light demand (*Eucalyptus globules* and *Acacia mearansi*) in Australia and at age of 6.5 found highest stem volume when compared with the mixture of 50% *Eucalyptus globules* and 50% *Acacia mearansii*. This mixture provided 20% more than pure *Acacia* stand, and more than 120% pure *Eucalyptus* stand. This showed the necessity of developing mixed plantation to maximize economic benefits and ecologic effects.

Hence, study the best performing mixture of *A. decurrens* and *E. camaldulensis* plantation and comparing with the sole plantation is required to optimize the benefits. The objective of this study is to evaluate effect *Eucalyptus* and *Acacia* mixed plantation on woody yield, soil nutrient changes and the economic benefits.

## MATERIALS AND METHODS

### Site description

The experiment was conducted in the high lands of the North shewa zone, Amhara Regional State of Ethiopia. Specifically, the study site is located in Merhabtie district, which is found in the south eastern part of Debere Birhan at a distance of 160 km south of Addis Abeba (Figure 1). The elevation of the study area reaches about 1931m.a.s.l, and the average annual rainfall ranges from 968 mm to 1486 mm.

### Experimental design

Seeds of *E. camaldulensis* and *A. decurrens* were purchased from Forestry Research Center (FRC) based at Addis Ababa, and raised in Alem Ketema nursery site. The treatments were arranged both a mixture and sole plantation. The proportions of *Acacia decurrens*: *Eucalyptus camaldulensis* were 0%:100%, 100%:0%, 25%:75%, 75%:25% and 50%:50%. Explicitly, the treatments were *Acacia decurrens* sole plantation, *Eucalyptus camaldulensis* sole plantation, *Acacia-Eucalyptus* mixed (one after the other on alternative rows), *Acacia-Eucalyptus* (three rows of *Eucalyptus* with one row of

*Acacia*), and *Acacia-Eucalyptus* (three rows of *Acacia* with one rows of *Eucalyptus*). The treatments were arranged in a randomized complete block design with three replications.

Each plot had a size of 6 m\*6 m size and the number of seedlings per plots was 16. The spacing between seedlings was 2 m\*2 m. The spacing between blocks and plots were 3 m and 2 m, respectively. At the beginning and end of the experiment, composite soil samples were collected from each treatment at 0-15 and 15. 1-30 cm depths. Soil physical properties (moisture content and bulk density) and chemical properties (organic carbon, total nitrogen and electric conductivity) were analyzed following the soil analysis procedures to investigate the difference before and after plantation. The economic effect of the mixed and sole plantation was assessed using the standard procedure. The data used for calculation were labor cost, estimated wood selling price, improvement/deterioration of soil fertility and social data.

### Data

Growth parameters (root collar diameter, diameter at breast height and height), survival count, pest and disease data were collected every six-month interval until the end of the trial. Soil samples were taken at the beginning and end of the experiment. Input cost for (labor, seed purchasing, and management) and estimated the final selling price of the wood.

### Data analysis

The volume of wood for standing trees was estimated by the following equation:  $V=0.44*B*H$ ; Where  $V$ =tree volume ( $m^3$ ),  $B$  is tree basal area at breast height ( $m^2$ ) and  $H$  is tree height. Sample data were analyzed with ANOVA to see the differences among species mean for each variable measured. Least Significance Difference test (LSD) was used for multiple comparisons when significant differences were found between treatments. Descriptive




Figure 1: Location map of the study area.

and other statistical analysis (ANOVA) were carried out using SPSS version 20 software.

## RESULTS AND DISCUSSION

### Growth performance

At the age of seven years, the overall mean height was 5.51 (  $\pm$  0.87 SD) m. The minimum observed height was 3.93 m and a maximum 7.27 m for Eucalyptus and 50% Eucalyptus proportions, respectively. The mean DBH was 6.33 (  $\pm$  1.44 SD) cm ranging from a minimum of 4.6 cm to a maximum of 9.5 cm (Table 1). The wood volume of the total stand was 21.12m<sup>3</sup>/ha with a standard deviation of 1.46 (Table 2).

A mix of 75% Acacia with 25% Eucalyptus and 50:50% of Acacia with Eucalyptus had shown a better survival rate compared to other treatments (Figure 2). Tree height was not significantly different among pure and mixed plantations ( $P > 0.05$ ) but at the age of 7 years, 50:50% *E. camaldulensis* : *A. decurrens* showed the highest mean height (6.1 m) growth than other treatments, while pure *E. camaldulensis* had the lowest height (4.6 m) growth (Figure 3).

The performance of the tree at the site might be ascribed to the soil conditions, especially poor nitrogen, at the trial site. The prime consideration for mixed plantation is the ecological requirements of species with climate, edaphic and topographic characteristics of the proposed planting site [15-17]. A careful selection of sites and species is, therefore, critical for successful establishment and improved growth performance of mixed plantations. Thus, a suitable mixtures should provide growth opportunities for each species to result in synergy among them. As the interaction of different species very divergent it is not easy to a generalization about how species will grow in a mixed plantation. However, height-age curves of each species in monoculture and knowledge of the relative's shade tolerances of each species should provide a useful basis for species selection.

### Tree volume

The study showed that tree volume production was not significantly different among a pure and mixed plantation of *E. camaldulensis* and *A. decurrens* species (Table 3) but 50:50% *E. camaldulensis* and

*A. decurrens* recorded slightly greater tree volume next to 100% *A. decurrens*. According to a study in Australia, a 6.5 year age of 50:50% *E. globules* and *A. mearnsii* mixed stand found to have the highest stem volume than other treatments, 20% and 120% more than a pure Acacia and Eucalyptus stand, respectively.

### Economic returns of pure and mixed plantation

The price of tree products at immediate harvest was estimated by evaluating the market prices each treatment by participating in the local community. Whereas the costs of tree management was calculated by considering the major management activities and it was similar for all treatments. According to this result, the 50:50% *E. camaldulensis*: *A. decurrens* generated more money than others. It




Figure 2: The Survival rate of pure and mixed *Acacia-Eucalyptus* plantation at different month after planting.




Figure 3: Bar graph of mean tree height of pure and mixed *Acacia* and *Eucalyptus* species at the age of 7 years.

Table 1: The mean value of the 7th year *Acacia-Eucalyptus* total plantation trial.

Parameters	Minimum	Maximum	Mean	Std. Error	Std. Deviation
Ht (m)	3.93	7.27	5.51	0.225	0.872
DBH (cm)	4.6	9.5	6.329	0.372	1.443
V (m <sup>3</sup> /ha)	7.18	48.31	21.12	3.31	1.46

Table 2: Mean volume ( $P=0.05$  and Std. Error=7.12) of a pure and mixed stand of *E. camaldulensis* and *A. decurrens* at age of 7 years.

The Proportion Of Tree Species In Each Treatment	Volume (m <sup>3</sup> /ha)
100%, <i>A. decurrens</i>	31.70
50:50%, <i>E. camaldulensis</i> : <i>A. decurrens</i>	29.92
75:25%, <i>E. camaldulensis</i> : <i>A. decurrens</i>	16.65
75:25%, <i>A. decurrens</i> : <i>E. camaldulensis</i>	17.49
100%, <i>E. camaldulensis</i>	9.873
Mean	21.12

Table 3: Mean values of soil pH, E.C, OC, TN under pure and mixed plantation (soil depth: 0-15 cm).

Treatments	PH	CEC	OC (%)	OM (%)	T.N (%)
100% <i>E. camaldulensis</i>	6.173	0.108	2.683	4.672	0.16
75: 25% <i>A. decurrens</i>	6.39	0.125	2.694	4.633	0.137
50: 50% <i>E. camaldulensis</i> : <i>A. decurrens</i>	6.333	0.157	3.24	5.572	0.2
25:75%, <i>E. camaldulensis</i> : <i>A. decurrens</i>	6.557	0.132	1.705	2.93	0.13
100% <i>A. decurrens</i>	6.507	0.125	2.659	4.57	0.157
Control	6.47	0.112	1.609	2.773	0.04
Mean	6.405	0.1265	2.437	4.191	0.1373
Std. Error	0.15	0.013	0.613	1.053	0.021
Sig(5%)	ns	ns	ns	ns	***

Where: Eu=*E. camaldulensis*, Ac=*Acacia decurrens*, CEC=Cation Exchange Capacity, OC=Organic Carbon, TN=Total Nitrogen and OM=organic matter

**Table 4:** Estimated cost-benefit analysis of each treatment at the final 7<sup>th</sup>-year harvest, note all measurements are in Birr/ha base.

Treatments	The Direct Benefit From The Trees	Indirect Benefit (Get From Grass)	Total Gross Benefit	Field Cost	Net Benefits	MNB%
100% Ec	99518.5	6944.4	106463	24305.5	82157.4	2.8
75:25%, Ec Ac	69333.3	6944.4	76277.7	24305.5	51972.2	
50:50% Eu :Ac	102111	6944.4	109055.5	24305.5	84749.9	2.9
75:25% Eu :Ac	70166.7	6944.4	77111.1	24305.5	52805.5	
100% Ac	46370.7	6944.4	53314.8	24305.5	29009.2	
Average	77500	6944.4	84444.4	24305.5		

has high marginal net benefit and was a combination profitable treatment (Table 4).

## CONCLUSION

In conclusion, pure stands of *E. camaldulensis* and *A. decurrens* plantations showed the lowest survival and growth performance than mixed plantation. The tree height and total volume did not show a significant difference between the mixed and pure stands. However, mixed plantation resulted in improvement of soil nutrients. A mix of 50:50% *E. camaldulensis*: *A. decurrens* plantations scored higher total nitrogen than other mixed plantations. In addition, 50:50% of *E. camaldulensis* and *A. decurrens* plantations provided better net benefit than other plantations. From this study, a 50:50% *E. camaldulensis* and *A. decurrens* is recommended for mixed plantation there by to obtain relatively better economic benefits and soil improvements.

## REFERENCES

- Dessie G, Erkossa AT. Eucalyptus in East Africa, Socio-economic and environmental issues. FAO, Planted Forests and Tress Working Papers, Working Paper FP 46/E, Forestry Department, FAO, Rome, Italy. 2011.
- Wingfield M, Slippers B, Hurley B, Coutinho T, Wing Field B, Roux J. Eucalypt pests and diseases: Growing threats to plantation productivity. Southern For J For Sci. 2008;70:139-144.
- Zerfu H. Ecological impact valuation of Eucalyptus plantations in comparison with agricultural and grazing land-use types in the highlands of Ethiopia Doctoral thesis. Institute of Forest Ecology, University of Natural Resources and Life Sciences, Vienna. J Forest Ecol. 2002.
- Kassahun HT, Collick AS, Adissu T, Ashagrie BB, Tessema ZK. Soil properties and fertility status dynamics of Northwestern Ethiopia as influenced by land use changes: Case of Dibanke watershed. 2008.
- Fisher RF, Binkley D. Ecology and management of forest soils (3<sup>rd</sup> edn): John Wiley and Sons, New York, USA, 2000.
- Alemie TC. The effect of Eucalyptus on crop productivity and soil properties in the Koga watershed Western Amhara Region, Ethiopia. Unpublished M.Sc. Thesis, Cornell University, USA, 2009.
- Baber S, Ahmad M, Bhatti A. The effect of *Eucalyptus camaldulensis* on soil properties and fertility. J Agric Biol Sci. 2006;1(3):47-50.
- Cao Y, Fu S, Zou X, Cao H, Shao Y, Zhou L. Soil microbial community composition under Eucalyptus plantations of different age in subtropical China. Eur J Soil Biol. 2010;46(2):128-135.
- Castro-Diez P, Fierro-Brunnenmeister N, Gonzalez-Munoz N, Gallardo A. Effects of exotic and native tree leaf litter on soil properties of two contrasting sites in the Iberian Peninsula. Plant Soil. 2011;350:179-191.
- Duguma LA, Hager H. Consumption and species preference for house construction wood in central high lands of Ethiopia-implications for enhancing tree growing. J For Res. 2010;21:104-110.
- Kaye JP, Resh SC, Kaye MW, Chimmer RA. Nutrient and carbon dynamics in a replacement series of Eucalyptus and Albizia trees. Ecol. 2000 81:3267-3273.
- Landon JR. Booker tropical soil manual: A hand book for soil survey and agricultural land evaluation in the tropics and sub-tropics. Longman Scientific and Technical, Essex, New York, USA. 1991;474.
- Leite FP, Silva IR, Ferreira R, De Barros NF, Neves LJC. Alterations of soil chemical properties by eucalyptus cultivation in five regions in the Rio Doce valley. For Ecol Manag. 2010;1:821-831.
- Mekonnen Z, Kassa H, Lemenh M, Camp Bell B. The role and management of eucalyptus in Lode Hetosa district, Central Ethiopia. For. Trees Livelihoods. 2007;17:309-323.
- Pohjonen V, Pukkala T. Eucalyptus globules in Ethiopian forestry. For Ecol and Manag. 1990;36:19-31.
- Tadesse T, Haque I, Aduayi EA. Working document: Soil, Plant, Fertilizer, Animal Manure and Compost analysis manual. International Livestock center for Africa, No.B13, and Addis Ababa, Ethiopia. 1991.
- Tererai F, Gaertner M, Jacobs SM, Richardson DM. *Eucalyptus camldulensis* invasion in riparian zones reveals few significant effects on soil physico-chemical properties. River Res Appl. 2014; 7(4):1-12.