

Journal of Clinical & Cellular Immunology

Review Article

The Role of Inflammatory Cytokines and the RANKL-RANK-OPG Molecular Triad in Periodontal Bone Loss-A Review

Prathiba Chichurakanahalli Srinivasan*

SV Dental Collge, Bangalore, Karnataka, India

Abstract

Periodontal diseases are chronic inflammatory disorders involving the supporting structures of the teeth. Connective tissue destruction and loss of bone support of the dentition are key features of this disease. Early studies have revealed that biofilm plaque accumulation in the dentogingival area is the primary etiological factor of periodontal diseases. Recent research has not only reinforced the bacterial etiology of periodontal disease, but has also emphasized the role of inflammation in this pathologic process. The host mounts an immune-inflammatory response to combat the bacterial attack. The host response is like a double-edged sword. It eliminates the offending pathogens, but overstimulation and amplification of the same leads to tissue destruction and bone loss. In the mid-1990s, extensive research in the field of bone biology led to the discovery of the RANKL-RANK-OPG molecular triad. This article explores the mechanisms by which inflammatory host response leads to alveolar bone loss-the role of cytokines, factors that stimulate osteoclastogenesis via the RANKL-RANK-OPG pathway and how inflammation interferes with the uncoupling of bone formation and bone resorption. In addition to the conventional therapeutic modalities aimed at eliminating the microbes, additional therapeutic strategies that interfere with the RANKL-RANK-OPG axis may have a protective effect on the bone loss.

Keywords: Inflammation; RANKL; OPG; RANK; Osteoclastogenesis; Cytokines; Periodontal disease; IL-1; TNF; Alveolar bone loss

Introduction

Periodontal diseases, which are chronic inflammatory disorders localized to the attachment structures of the teeth, are considered to be the major cause of tooth loss in adults and the most prevalent form of bone pathology in humans [1]. The term periodontal disease refers to both gingivitis and periodontitis. Gingivitis is an inflammatory condition of the soft tissues surrounding the teeth [2]. If unchecked, gingivitis progresses to periodontitis, an inflammation of the supporting structures of the teeth [3]. Specific gram-negative anaerobic bacteria species such as Porphyromonas gingivialis and Tannerella forsythia have been associated with periodontitis [4].

The initial host response to bacterial infection is a local inflammatory reaction that activates the innate immune system [5]. An imbalance between the plaque biofilm and the host immune system results in the overexpression of an array of proinflammatory cytokines, propagation of inflammation through the gingival tissues and the subsequent destruction of alveolar bone [6].

Thus the inflammatory process results in destruction of connective tissue and alveolar bone, the hallmarks of periodontal disease.

The Role of Inflammation in Alveolar Bone Loss

There are two critical factors, which dictate whether bone loss occurs in response to inflammatory reaction.

1. The concentration of pro-inflammatory cytokines and mediators present in the gingival tissue must be sufficient to activate the pathways leading to bone resorption.

2. The inflammatory mediators must penetrate the gingival tissue to reach within a critical distance to alveolar bone [5].

Page and Schroeder showed that bone resorption ceases when a 2.5 mm zone is created between the site of bacteria and bone [7].

Inflammatory Mediators

There are several classes of molecules that activate a host response which in turn stimulates osteoclastogenesis either directly or indirectly [8]. These include lipid-based molecules such as prostaglandins, leukotrienes [8], pro-inflammatory cytokines such as IL-1, IL-6, IL-11 and IL-17, tumor necrosis factor-alpha, leukemia inhibitory factor, and oncostatin M. The kinins such as bradykinin, kallidin and thrombin and various chemokines can also lead to bone resorption [9].

Chemokines are chemotactic cytokines that stimulate recruitment of inflammatory cells. Based on the structure of their ligand, they are divided into two major families-CC and CXC. Their receptors are referred to as CC chemokine receptor (CCR) and CXC chemokine receptor (CXCR). CXCR2 receptor binds several chemokines that are neutrophil chemoattractants [10]. Neutrophil recruitment is important to protect against bacterial attack. Yu JJ et al. conducted a study in which oral gavage of Porphyromonas gingivalis was administered to CXCR2-and IL-17-deficient mice. The results revealed increased periodontal bone loss in these mice compared to wild-type mice, suggesting that chemokines and IL-17 are important in protecting host against pathogen-initiated bone loss. IL-17 is a key cytokine produced by a newly identified subset of T helper cells (Th) called the "Th17" lymphocytes and it is important in stimulating chemokine production to recruit neutrophils. The enhanced bone loss observed in this study is the result of a defective ability to stimulate neutrophil migration to the inflamed bone, which requires IL-17 receptor dependant signals,

*Corresponding author: Prathiba Chichurakanahalli Srinivasan, SV Dental Collge, Bangalore, Karnataka, India, E-mail: csprati@yahoo.com

Received June 13, 2013; Accepted August 27, 2013; Published August 31, 2013

Citation: Srinivasan PC (2013) The Role of Inflammatory Cytokines and the RANKL-RANK-OPG Molecular Triad in Periodontal Bone Loss-A Review. J Clin Cell Immunol S13: 007. doi:10.4172/2155-9899.S13-007

Copyright: © 2013 Srinivasan PC. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Page 2 of 8

thus underscoring the importance of IL-17 in the defense against oral pathogens [11]. From their study Zhang et al. concluded that IL-17A produced by Th17 cells stimulate the development of osteoclasts (osteoclastogenesis) in the presence of osteoblasts [12] and Cardoso et al. observed expression of IL-17 in gingiva from patients with periodontitis [13].

The role of pro-inflammatory cytokines, interleukin-1 (IL-1) and tumor necrosis factor (TNF) in bacteria-induced bone loss was demonstrated in an animal model study by Assuma et al. Porphyromonas gingivalis (Pg)-soaked silk ligatures were applied to the posterior mandibular teeth of Macaca fascicularis primates to induce experimental periodontitis. The primates received local injections, over a period of 6 weeks, of antagonists to TNF-alpha and IL-1 (soluble TNF-alpha and IL-1 receptors) or vehicle control. The gingival connective tissue sections in close proximity to bone demonstrated significant inflammatory cell recruitment and osteoclast formation surrounding bone in the control primates. Thus, infection with Pg in these animals was associated with expansion of the inflammatory front to alveolar bone. The antagonists to cytokines reduced the appearance of inflammatory cells in this region and the formation of bone resorbing osteoclasts. Thus, inhibition of inflammatory mediators can prevent the "inflammatory front" from reaching the alveolar bone leading to a reduction in bone loss as observed in this animal model [14]. In a follow-up study with the same model Delima et al. in 2001, stated that IL-1 and TNF blockers inhibited the loss of connective tissue attachment [15].

In their study, Graves DT et al. concluded that P. gingivalis induced osteoclastogenesis was reduced in TNF receptor-deficient mice compared to wild-type controls, thus indicating that osteoclast formation resulted from stimulation of the host response rather that from the direct effect of bacterial products [16]. Gaspersic et al in rat ligature model study concluded that the administration of recombinant human TNF-alpha (rhTNF-alpha) accelerated the progression of periodontitis [17]. In a study by Garlet et al. TNF-alpha receptor-1knockout mice developed significantly less inflammation and alveolar bone loss in response to Aggregibacter actinomycetemcomitans oral gavage. The levels of the pathogen as quantified by real-time PCR was higher in the TNF receptor ablated mice, compared to the wild-type controls. The quantitative mRNA expression of inflammatory cytokines IL-1 beta, interferon-gamma, and receptor activator of nuclear factor-kappa B ligand (RANKL) was significantly lower in the TNF receptor ablated mice. Thus, the absence of TNFR-1 resulted in lower production of cytokines in response to the pathogen. Decrease in TNFalpha seemed to reduce the host response, thereby leading to higher levels of bacteria. However, there was reduced expression of cytokines that stimulate bone resorption, which resulted in less net bone loss [18]. In response to P. gingivalis oral gavage, mice with genetically deleted IL-6 had decreased bone loss compared to wild-type mice, thus indicating the pro-inflammatory effects of IL-6 leading to bone resorption. (Baker PJ et al.) [19]. Li et al. examined the role of B and CD4 T cells in adaptive immunity of rats infected with Aggregatibacter actinomycetemcomitans (Aa). Sprague-Dawley male rats were fed Aa-containing mash or control-mash for 2 weeks. B and CD4 T cells were obtained from draining lymph nodes at 2, 4 and 12 weeks, post inoculation. Quantitative polymerase chain reaction-based messenger RNA expression was conducted for 89 cytokine family genes. B and CD4 T cells of Aa-infected rats increased and were activated, resulting in enhanced isotype-switched serum immunoglobulin G by 2 weeks post inoculation. Bone resorption was evident 12 weeks after Aa-feeding. In B cells, interleukin-2 (IL-2), macrophage-inhibiting factor, IL-19, IL-

21, tumor necrosis factor (TNF), CD40 ligand (CD40L), CD70, bone morphogenetic protein 2 (BMP2), BMP3, and BMP10 were upregulated early; while IL-7, Fas ligand (FasL), small inducible cytokine subfamily E1, and growth differentiation factor 11 (GDF11; BMP11) were upregulated late (12 weeks). BMP10 was sustained throughout. In CD4 T cells, IL-10, IL-16, TNF, lymphotoxin-beta (LTβ), APRIL, CD40L, FasL, RANKL and osteoprotegerin were upregulated early, whereas IL-1β, IL-1RN, IL-1F8, IL-24, interferon-α1, GDF11 (BMP11), and GDF15 were upregulated late (12 weeks). There was upregulation in mRNA for a number of cytokines not normally ascribed to periodontal disease. IL-16 was upregulated in CD4 T cells in the early phase of the response [20]. IL-16 has been shown to be involved in the selective migration of CD4 T cells, and participates in inflammatory diseases [21]. IL-19, a novel cytokine of the IL-10 family, was also upregulated in CD4 T cells in response to Aa [20]. IL-19 produced by synovial cell in Rheumatoid arthritis (RA) patients promotes joint inflammation (Sakurai et al.) [22]. IL-21, which has recently been shown to induce receptor activator of nuclear factor kappaB ligand (RANKL) and was implicated in arthritis (Jang et al.) [23] was upregulated in B cells responding to Aa [20]. By 12 weeks, there was also an induction of IL-24 in CD4 T cells responding to Aa [20]. IL-24 was increased in the synovium of patients with rheumatoid arthritis, and this cytokine was implicated in recruitment of neutrophil granulocytes (Kragstrup et al.) [24]. B cellactivating-factor (BAFF, or TNFSF13B) and a proliferation-inducing ligand (APRIL), members of the TNF family, were upregulated in B cells and CD4 T cells, respectively, in response to Aa infection [20]. IL-23, a proinflammatory cytokine composed of IL-23p19 and IL-12/23p40 subunits, is known to promote the differentiation of Th17 cells. Studies showed that IL-23 and IL-12 were expressed at significantly higher levels in periodontal lesions than in control sites, suggesting that IL-23-induced Th17 pathway is stimulated in inflammatory periodontal lesions (Ohyama et al.) [25]. IL-33 is a new member of the IL-1 family, which plays a role in inflammatory response. Injecting IL-33 or IL-33R agonistic antibody into TNF transgenic mice overexpressing human TNF inhibited the development of spontaneous joint inflammation and cartilage destruction. In vitro, IL-33 directly inhibits mouse and human M-CSF/receptor activator for NF-kß ligand-driven osteoclast differentiation, suggesting an important role for IL-33 as a boneprotecting cytokine with potential for treating bone resorption (Zaiss et al.) [26]. The bone resorption protein RANKL and its soluble decoy receptor OPG was also induced in the CD4 T cells of Aa-fed rats [20].

The role of bone morphogenetic proteins (BMPs) and growth differentiation factors (GDFs) in periodontal disease

Bone morphogenic proteins (BMPs) and growth differentiation factors (GDFs) are members of the transforming growth factor- β (TGF- β) superfamily. They play important roles during development and organogenesis in delivering positional information in both vertebrates and invertebrates, and are involved in the development of hard as well as soft tissue (Herpin, Lelong and Favrel) [27]. BMP2 was induced in B cells early (week 4) of the inflammatory process, at the same time that RANKL was induced in CD4 T cells (Li et al.). This suggests that bone repair mechanisms were induced early, well ahead of impending bone resorption. However, by 12 weeks of infection by Aa, BMP2 was shut down, as bone resorption proceeded. BMP3 was also upregulated at week 4 in B cells responding to Aa [20]. BMP3 has been shown to be a negative regulator in the skeleton, as mice lacking BMP3 have increased bone mass. Transgenic mice over-expressing BMP3 had altered endochondral bone formation resulting in spontaneous rib fractures [28]. Both BMP2 and BMP3 were upregulated in B cells at the

same time (4 weeks post infection), and were shut down at 12 weeks, at which time bone resorption was evident [20]. B cells responding to *Aa* upregulated BMP10 at all time points [20]. BMP10 has been shown to regulate myocardial hypertrophic growth (Chen et al. 2006) [29]. BMP-10 may function as a tumor suppressor and apoptosis regulator for prostate cancer [30]. Thus the expression of BMP 10 in the Aa rat model at all points suggests it's role in inflammation and bone resorption [20]. Also the involvement of BMP10 in cardiac hypertrophy [29] and in prostate cancer [30] suggests a possible link between periodontal disease and systemic diseases.

Growth differentiation factor 11 (GDF11) or BMP11, plays an important role in establishing embryonic axial skeletal patterns [31]. In the GDF11 was upregulated at 12 weeks post infection, in both B and CD4 T cells, at the time of bone resorption. This suggests that GDF11 may have a novel role in bone resorption [20].

The Aa-induced rat model, the Pg-induced mouse model and several other animal studies discussed above strongly support the role of pro-inflammatory cytokines in bone loss. The role of novel cytokines not previously ascribed to periodontal disease has been extensively discussed and these studies also suggest a possible link between periodontal disease and systemic diseases.

Osteoimmunology

Extensive research in the fields of immunology and osteology has revealed an intimate relationship between inflammation/the immune system and the skeletal system [32].The immunoskeletal interface involves centralization of the immune and skeletal functions around common cells types and cytokine effectors that control the physiological bone mass. Prolonged immune activation can lead to skeletal deterioration. This interconnection of the skeletal and the immune systems has spawned the emergence of a new field of science termed "osteoimmunology" [33]. The field of osteoimmunology offers better insights into the understanding of pathogenesis of periodontal disease by explaining the role of inflammation in alveolar bone loss.

Osteoclasts and the Receptor Activator of NF-Kappa B (RANK)/RANK Ligand (RANKL)/Osteoprotegerin (OPG) System

Osteoclasts are unique cells of the human body tasked with resorbing bone. These multinucleated giant cells bind down onto the bone surfaces creating a sealing zone and a ruffled-membrane border into which apical proton pumps accumulate hydrogen ions that combine with chloride ions to form hydrochloric acid, which degrades bone mineral. The exposed collagen matrix is attacked by acid resistant endosomal and lysosomal enzymes that cleave collagen fibers effectively removing small quantities of bone [34].

How osteoclasts form, what regulates their differentiation and activity was an enigma. Research has revealed that osteoclast precursors circulate amid the monocyte/macrophage population and differentiate into preosteoclasts that fuse and form giant bone resorbing mature osteoclasts [35]. Inflammatory processes such as rheumatoid arthritis and periodontal infection predispose to osteoclastic bone resorption by production of pro-inflammatory cytokines such as IL-1 and TNFalpha [36]. The osteoclast however, does not possess receptors for the cytokines or hormones. So, for years the biologists were perplexed as to how they resorbed bone. This conundrum was solved with the discovery of the osteoprotegerin (OPG), receptor activator of nuclear factor-kappa B ligand (RANKL) and receptor activator of nuclear factor kappa-B (RANK).

Discovery of OPG/RANKL/RANK Molecular Triad -The Solving of the Puzzle

In 1981, Rodan and Martin proposed a novel hypothesis wherein the osteoblast played a central role in mediating the hormonal control of osteoclastogenesis and bone resorption [37]. But, the factors expressed by the osteoblast/stromal cells remained undetermined until they were discovered independently by four groups using different approaches. Boyle and coworkers at the Amgen Inc (Thousand Oaks, CA, USA) discovered OPG serendipitously when they were attempting to identify tumor necrosis factor receptor-related molecules with possible therapeutic utility by generating transgenic mice that overexpress various TNF receptor related cDNAs. Mice over-expressing one particular cDNA developed marked osteopetrosis because they did not have osteoclasts in their bones. The protein encoded by the gene was termed osteoprotegerin-OPG (the bone protector) [38].

Using the standard approach to test the Rodan-Martin hypothesis of purifying a factor from human embryonic fibroblasts that inhibited osteoclastogenesis, researchers at Snow Brand Milk Products Co. (Sapporo, Hokkaido, Japan) reported their discovery of an identical molecule. They obtained a partial protein sequence and subsequently cloned the cDNA for OPG [39].

The discovery of OPG became the key to identify the long-sought osteoclast differentiation factor expressed on osteoblastic/stromal cell that was essential for osteoclast development. Using expression cloning and OPG as a probe, both the groups quickly identified its ligand, which they termed OPG ligand [40] and osteoclast differentiation factor [41] respectively. This ligand turned out to be identical to a member of the TNF ligand family, which had been identified in the preceding year as RANKL [42] and TNF-related activation induced cytokine (TRANCE) [43]. The cellular receptor was identified as being identical to previously identified RANK, which Anderson and coworkers at Immunex (Seattle, WA, USA) had discovered while they were sequencing cDNAs from a human bone marrow derived myeloid dendritic cell cDNA library. [42]. Both the groups found that RANK had partial homology to a portion of the extracellular domain of human CD40, a member of the TNF receptor superfamily, and it was involved in the activation of T cells in the immune system. The groups then isolated RANKL by direct expression screening and found, like Wong and coworkers [43] did, that it increased dendritic cell stimulated naïve T cell proliferation and survival of RANK-expressing T cells. These discoveries led to the conclusion that RANKL is involved in osteoclastogenesis and T cell activation.

Osteoclast Formation/Osteoclastogenesis

Osteoclastogenesis is coordintated by the interaction of the three members of the TNF superfamily-OPG/RANKL/RANK [44] RANKL is expressed predominantly as a membrane-bound ligand on osteoblasts, fibroblasts, and activated T and B cells and its osteoclastogenic action can be blocked by its soluble decoy receptor osteoprotegerin (OPG) [9,35]. Together with Macrophage-colony stimulating factor (mCSF), RANKL is a key cytokine in the induction of osteoclastogeneis both *in vitro* and *in vivo* [45]. It binds directly to RANK on the surface of preosteoclasts and osteoclasts, stimulating both the differentiation of osteoclast progenitors and the activity of mature osteoclasts [40,46,47].

Regulation of Bone Remodeling - The Role of the RANKL-RANK-OPG System

Systemic factors and local stimuli in the bone microenvironment

Page 4 of 8

target the molecular interplay of RANKL-RANK-OPG cytokine system, thus regulating the remodeling of alveolar bone. (Lerner 2; Boyle et al. Mundy et al. Horowitz et al.) [48-51]. Systemic modulators include calcitonin, parathyroid hormone, thyroid hormone, gonadal steroids, vitamin D, glucocorticoids and growth hormone. Local mediators include pro-inflammatory cytokines as well as prostanoids (Lerner) [48]. Bacterial products may interfere with bone remodeling balance by regulating RANKL expression (Lerner, Henderson and Nair) [48,52]. Gram-negative bacteria such as *A. actinomycetemcomitans*, (Teng et al. Kikuchi et al. Teng et al.) [53-55], *P. gingivalis* (Okahashi et al.) [56], Treponema denticola (Choi et al.) [57] and *Streptococcus pyogenes* (Okahashi et al. Sakurai et al.) [58,59] have been shown to upregulate RANKL expression in a variety of cells.

RANKL-RANK-OPG in Periodontal Disease

Studies have revealed that during inflammatory response, cytokines, chemokines and other mediators stimulate periosteal osteoblasts, altering the expression levels of receptor activator of nuclear factorkappa B ligand (RANKL) on the osteoblast surface [9,49]. RANKL is expressed by osteoblasts in the form of a membrane-bound protein (mRANKL) or cleaved into a soluble form (sRANKL) [60,61]. IL-1 stimulates osteoclastogenesis and bone resorption, largely through up-regulation of RANKL, while TNF can stimulate osteoclastogenesis directly or indirectly through RANKL [62]. Inhibition of RANK ligand caused a decrease in alveolar bone loss in several models of periodontal disease (Teng et al. Han et al. Jin et al.) [55,63,64].

Teng et al. in their study demonstrated the role of $CD4^+$ T cells in mediating periodontal bone loss. Also, they found that administration of RANK ligand inhibitor; osteoprotegerin reduced osteoclastogenesis and inhibited alveolar bone destruction [55]. Han et al. in 2006 demonstrated that B cells contribute to periodontal bone loss in the absence of T cells through production of RANKL [63].

Bone resorption and formation are regulated by the relative concentrations of RANKL, RANKL receptor RANK on osteoclast

precursor cells and the soluble decoy receptor OPG. When RANKL expression is enhanced relative to OPG, RANKL is available to bind to RANK on osteoclast precursors, tipping the balance to favor activation of osteoclast formation and bone resorption [49]. The binding of RANKL to osteoclast precursors occurs at a stage when hematopoietic stem cells have differentiated from the colony forming unit for granulocytes and macrophages to the colony forming unit for macrophages (CFU-M). Binding of RANKL to RANK on CFU-M in the presence of macrophage colony stimulating factor induces differentiation of preosteoclast into a multinucleated cell that becomes a mature osteoclast, which then resorbs bone [65]. When OPG concentrations are high relative to RANKL expression, OPG binds RANKL, inhibiting it to bind to RANK. Preventing the binding of RANKL to RANK leads to reduced formation of osteoclasts and apoptosis of preexisting osteoclasts [49]. Jin et al. in 2007 conducted a study in which rats were administered subcutaneously an OPG-Fc fusion protein to bloc RANK ligand. Compared to the controls, reduced osteoclastogenesis and significant preservation of the alveolar bone volume was observed in among the OPG-Fc treated rats [64]. Leiwiecki in 2009 demonstrated that treatment with Denosumab, a human monoclonal antibody specific for RANKL reduces the risk for fractures and increases the bone mineral density in post-menopausal women. It also inhibits osteoclast-mediated damage caused by rheumatoid arthritis [66].

The studies discussed above have provided valuable insights into the role of inflammation or host response in the pathogenesis of periodontal disease. The studies have unequivocally demonstrated that over amplification of the immune system releases a host of pro-inflammatory cytokines and mediators that can lead to osteoclastogenesis. Also, these inflammatory mediators can stimulate the release of RANK ligand and thereby lead to bone resorption.

The Assessment of RANKL/OPG Ratio in the Severity of Periodontitis

A number of clinical studies have investigated the concentrations

Study	Healthy subjects (n)	Gingivitis subjects (n)	Mild perio subjects (n)	Moderate perio subjects (n)	Chronic perio subjects (n)	Generalized aggressive severe subjects (n)	Chronic with immunosuppressant subjects (n)	Conclusion
Bostanci et al. (GCF) [68]	21	22			28	25	11	GCF RANKL &OPG were oppositely regulated in periodontitis groups.
Bostanci et al. Gingival tissue [69]	9	8			11	12	10	RANKL/OPG ratio increased in all periodontitis groups.
Lu et al. (GCF and gingiva) [6]	4				20			GCF RANKL, but not OPG was elevated in periodontitis groups.
Mogi et al. (GCF) [70]	28		27	58		47		RANKL/OPG ratio was significantly elevated in periodontitis groups.
Liu et al. (Gingival tissue) [71]	6			27		25		RANKL/OPG was elevated in periodontitis groups.
Kawai et al. (gingiva and blood) [72]	12				32			sRANKL, but not OPG was significantly higher in periodontitis gropus.
Wara-aswapati et al. (gingiva and plaque) [73]	15				15			RANKL/OPG ratio was significantly higher in perioidontitis groups
Garlet et al. (gingival tissue) [74]	10	7			20	16		RANKL/OPG and MMP/TIMP expression determined disease progression/severity
Nagasawa et al. (gingival tissue) [75]	2				30			OPG is induced by LPS- stimulated gingival fibroblast.

sRANKL: Soluble Receptor Activator of Nuclear Factor-Kappa B Ligand; TIMP: Tissue Inhibitors of Metalloproteinases; LPS: Lipopolysaccharides; perio: Periodontitis **Table 1:** Summary of human studies evaluating the RANKL and OPG in Periodontal Disease. of RANKL and OPG in the gingival crevicular fluid of individuals with periodontitis to determine the RANKL/OPG ratio. Table 1 summarizes the human studies evaluating the RANKL and OPG levels in periodontal disease [67].

With a net increase in the ratio of RANKL/OPG in gingiva and crevicular fluid associated with bone loss and with the severity of periodontal disease, the possibility of interference with the RANK/RANKL/OPG axis may offer novel therapeutic outcomes. The desired outcome would be an increase in OPG or a decrease in RANKL that brings the OPG/RANKL ratio to a balance where bone formation is equal to bone resorption. A summary of animal studies in which osteoprotegerin fusion protein (OPG-Fc) and other inhibitors of RANK-mediated osteoclastogenesis are presented in Table 2 [67].

Thus the inference from the above mentioned animal studies is, therapeutic modalities which interfere with the RANK-RANKL-OPG axis will have a protective effect on osteoclastogenesis and periodontal bone loss.

Role of Cytokines in Bone Uncoupling

Bone is resorbed by osteoclasts, following which new bone is laid down by osteoblasts in the resorption lacunae. Under physiologic conditions, the two activities are coupled, i.e., the amount of bone formed by osteoblasts is equal to that resorbed by osteoclasts. In pathologic processes like periodontal disease and osteoporosis the two processes are uncoupled i.e., there is deficient bone formation following resorption [80].

The inflammatory process that leads to osteoclastogenesis and bone resorption may also be responsible for the failure to form adequate amount of new bone i.e., inflammation causes uncoupling of bone formation following bone resorption [8]. Al-Masahat et al. in 2006 and Liu et al. in 2006 stated that prolonged inflammation in diabetic animals interferes with bone formation in the periodontium following an episode of bone resorption. The prolonged inflammation may stimulate the death of the osteoblasts [81,82]. Application of cytokines like IL-1 beta and TNF-alpha in vivo not only stimulates bone resorption but limits bone formation by inhibiting the coupling process (Bertolini et al. Nyugen et al.) [83,84]. Osteoblast survival is a key factor in bone formation. A study by Diarra and colleagues reported that TNF-alpha stimulates production of Dickkopf-1 (DKK-1), which suppresses bone formation by inhibiting the WNT (wingless WNT/ beta catenin) pathway. DKK-1, a negative regulator of WNT pathway is up-regulated by TNF stimulation through TNF-1 receptor and P38 MAPK (mitogen-activating protein kinase) signaling. The up-regulated DKK-1 not only promotes bone resorption but also blocks bone

Study Method	Study Method				
Jin et al. [64]	Systemic delivery of rhOPG-Fc				
Teng et al. [55]	Intraperitoneal injection of srOPG-Fc				
Valverde et al. [76]	Subcutaneous kaliotoxin, K+-channel blocker T cells.				
Kawai et al. [77]	Intraperitoneal injection of OPG-Fc				
Mahamed et al. [78]	Intraperitoneal injection of hu-OPG-Fc				
Rogers et al. [79]	Oral gavage of SD282, a p38 MAPK inhibitor				
Assuma et al. [14]	Intrapapillary injection of TNF-alpha and IL-1 antagonists.				

rhOPG-Fc: Human recombinant osteoprotegerin fusion protein; MAPK: Mitogen-Activated Protein Kinase; SFRP1: Secreted Frizzled-related Protein 1; hOPG-Fc: Human Osteoprotegerin fusion protein; srOPG-Fc: soluble recombinant Osteoprotegerin fusion protein; huOPG-Fc: human osteoprotegerin fusion protein; SD282: an indole 5-carboxamide selective p38* MAPK inhibitor (Scois, Fremont, CA). (David L Cochran)

 Table 2: Summary of animal interventional trials of periodontal disease.



formation and repair in the diseased joint [85]. Thus, inflammatory cytokines like TNF-alpha can limit bone formation by inhibiting osteoblast differentiation. In addition, the pro-inflammatory cytokines may directly stimulate osteoblast or osteoblast precursor apoptosis or indirectly affect it by stimulating expression of Fas, a potent apoptotic mediator [86]. Periodontal ligament cells are an important source of osteoblast precursors. TNF-alpha induced apoptosis of periodontal ligament cells may affect the pool of osteoblast precursors [87]. Another mechanism for uncoupling is the reduced function of osteoblasts mediated by diminished production of bone matrix proteins. A study by Centrella et al. in 1988 showed that TNF-alpha reduces collagen production and alkaline phosphatase activity in cells obtained from fetal rat parietal bone [88]. TNF-alpha and TNF-beta induce a twofold to three-fold reduction in synthesis of non-collagen bone matrix proteins like osteocalcin by osteoblasts [89]. Eguchi et al. stated that TNF specific inhibitor, etanercept, promotes BMP-2 induced ectopic bone formation when applied systemically or locally in vivo, thereby improving the coupling process [90].

Thus evidence from the literature has unequivocally established the negative impact of the inflammatory process on bone coupling. The net bone loss observed in periodontitis is due to the cumulative effect of inflammation-induced bone loss and the uncoupling of bone resorption and bone formation (Figure 1).

Page 5 of 8

Citation: Srinivasan PC (2013) The Role of Inflammatory Cytokines and the RANKL-RANK-OPG Molecular Triad in Periodontal Bone Loss-A Review. J Clin Cell Immunol S13: 007. doi:10.4172/2155-9899.S13-007

Conclusion

Periodontal diseases are chronic inflammatory disorders that afflict the supporting structures of the teeth. It is widely recognized that bacteria initiate periodontal diseases. The host responds to this bacterial attack by mounting an immuno-inflammatory response. The host response is essential to ward off the bacterial attack and prevent the systemic dissemination of infection. But in the process, the inflammatory response can also cause tissue destruction and alveolar bone loss. Prolonged inflammation and amplification of the inflammatory response releases a wide spectrum of pro-inflammatory cytokines and mediators. The cytokines and pro-inflammatory mediators propagate the inflammatory process and the "inflammatory front" advances to the alveolar bone. Alveolar bone loss is the cardinal sign of periodontitis. Experimental periodontitis studies have shown that cytokines like IL-1 and TNF-alpha can lead to alveolar bone loss. These cytokines can also induce osteoclastogenesis via the RANKL-RANK-OPG pathway. IL-1 and TNF-alpha also have a negative impact on bone coupling. The net bone loss seen in periodontitis is due to the cumulative effect of inflammation-induced bone loss and the uncoupling of bone formation and bone resorption. This intricate relationship between inflammation and bone metabolism has led to a new field of science termed "osteoimmunology." Extensive research in field of bone biology has provided valuable insights into pathogenesis of periodontal disease. Inflammation is now recognized as the central component of periodontal disease. Better understanding of the disease process will pave the way for adopting novel treatment approaches and thereby improve clinical outcomes. Therapeutic agents antagonistic to the inflammatory mediators or agents which block RANK ligand may be useful adjuncts to the conventional periodontal treatment procedures. We are still at the tip of the iceberg and longitudinal human clinical trials are required to prove the benefits of these therapeutic procedures.

References

- 1. Papapanou PN (1999) Epidemiology of periodontal diseases: an update. J Int Acad Periodontol 1: 110-116.
- Haimov-Kochman R, Kochman T, Stabholz A, Hochner-Celinkier D (2004) Bisphosphonate and estrogen replacement therapy for postmenopausal periodontitis. Isr Med Assoc J 6: 173-177.
- Buencamino MC, Palomo L, Thacker HL (2009) How menopause affects oral health, and what we can do about it. Cleve Clin J Med 76: 467-475.
- Brennan RM, Genco RJ, Wilding GE, Hovey KM, Trevisan M, et al. (2007) Bacterial species in subgingival plaque and oral bone loss in postmenopausal women. J Periodontol 78: 1051-1061.
- Graves DT, Cochran D (2003) The contribution of interleukin-1 and tumor necrosis factor to periodontal tissue destruction. J Periodontol 74: 391-401.
- Lu HK, Chen YL, Chang HC, Li CL, Kuo MY (2006) Identification of the osteoprotegerin/receptor activator of nuclear factor-kappa B ligand system in gingival crevicular fluid and tissue of patients with chronic periodontitis. J Periodontal Res 41: 354-360.
- Page RC, Schroeder HE (1981) Current status of the host response in chronic marginal periodontitis. J Periodontol 52: 477-491.
- Graves DT, Li J, Cochran DL (2011) Inflammation and uncoupling as mechanisms of periodontal bone loss. J Dent Res 90: 143-153.
- Lerner UH (2006) Inflammation-induced bone remodeling in periodontal disease and the influence of post-menopausal osteoporosis. J Dent Res 85: 596-607.
- Graves D (2008) Cytokines that promote periodontal tissue destruction. J Periodontol 79: 1585-1591.
- Yu JJ, Ruddy MJ, Wong GC, Sfintescu C, Baker PJ, et al. (2007) An essential role for IL-17 in preventing pathogen-initiated bone destruction: recruitment of neutrophils to inflamed bone requires IL-17 receptor-dependent signals. Blood 109: 3794-3802.

- Zhang F, Tanaka H, Kawato T, Kitami S, Nakai K, et al. (2011) Interleukin-17A induces cathepsin K and MMP-9 expression in osteoclasts via celecoxibblocked prostaglandin E2 in osteoblasts. Biochimie 93: 296-305.
- Cardoso CR, Garlet GP, Crippa GE, Rosa AL, Júnior WM, et al. (2009) Evidence of the presence of T helper type 17 cells in chronic lesions of human periodontal disease. Oral Microbiol Immunol 24: 1-6.
- Assuma R, Oates T, Cochran D, Amar S, Graves DT (1998) IL-1 and TNF antagonists inhibit the inflammatory response and bone loss in experimental periodontitis. J Immunol 160: 403-409.
- Delima AJ, Oates T, Assuma R, Schwartz Z, Cochran D, et al. (2001) Soluble antagonists to interleukin-1 (IL-1) and tumor necrosis factor (TNF) inhibits loss of tissue attachment in experimental periodontitis. J Clin Periodontol 28: 233-240.
- Graves DT, Oskoui M, Volejnikova S, Naguib G, Cai S, et al. (2001) Tumor necrosis factor modulates fibroblast apoptosis, PMN recruitment, and osteoclast formation in response to P. gingivalis infection. J Dent Res 80: 1875-1879.
- Gaspersic R, Stiblar-Martincic D, Osredkar J, Skaleric U (2003) Influence of subcutaneous administration of recombinant TNF-alpha on ligature-induced periodontitis in rats. J Periodontal Res 38: 198-203.
- Garlet GP, Cardoso CR, Campanelli AP, Ferreira BR, Avila-Campos MJ, et al. (2007) The dual role of p55 tumour necrosis factor-alpha receptor in Actinobacillus actinomycetemcomitans-induced experimental periodontitis: host protection and tissue destruction. Clin Exp Immunol 147: 128-138.
- Baker PJ, Dixon M, Evans RT, Dufour L, Johnson E, et al. (1999) CD4(+) T cells and the proinflammatory cytokines gamma interferon and interleukin-6 contribute to alveolar bone loss in mice. Infect Immun 67: 2804-2809.
- Li Y, Messina C, Bendaoud M, Fine DH, Schreiner H, et al. (2010) Adaptive immune response in osteoclastic bone resorption induced by orally administered Aggregatibacter actinomycetemcomitans in a rat model of periodontal disease. Mol Oral Microbiol 25: 275-292.
- Akiyama K, Karaki M, Kobayshi R, Dobashi H, Ishida T, et al. (2009) IL-16 variability and modulation by antiallergic drugs in a murine experimental allergic rhinitis model. Int Arch Allergy Immunol 149: 315-322.
- Sakurai N, Kuroiwa T, Ikeuchi H, Hiramatsu N, Maeshima A, et al. (2008) Expression of IL-19 and its receptors in RA: potential role for synovial hyperplasia formation. Rheumatology (Oxford) 47: 815-820.
- Jang E, Cho SH, Park H, Paik DJ, Kim JM, et al. (2009) A positive feedback loop of IL-21 signaling provoked by homeostatic CD4+CD25- T cell expansion is essential for the development of arthritis in autoimmune K/BxN mice. J Immunol 182: 4649-4656.
- Kragstrup TW, Otkjaer K, Holm C, Jørgensen A, Hokland M, et al. (2008) The expression of IL-20 and IL-24 and their shared receptors are increased in rheumatoid arthritis and spondyloarthropathy. Cytokine 41: 16-23.
- Ohyama H, Kato-Kogoe N, Kuhara A, Nishimura F, Nakasho K, et al. (2009) The involvement of IL-23 and the Th17 pathway in periodontitis. J Dent Res 88: 633-638.
- Zaiss MM, Kurowska-Stolarska M, Böhm C, Gary R, Scholtysek C, et al. (2011) IL-33 shifts the balance from osteoclast to alternatively activated macrophage differentiation and protects from TNF-alpha-mediated bone loss. J Immunol 186: 6097-6105.
- Herpin A, Lelong C, Favrel P (2004) Transforming growth factor-beta-related proteins: an ancestral and widespread superfamily of cytokines in metazoans. Dev Comp Immunol 28: 461-485.
- Gamer LW, Cox K, Carlo JM, Rosen V (2009) Overexpression of BMP3 in the developing skeleton alters endochondral bone formation resulting in spontaneous rib fractures. Dev Dyn 238: 2374-2381.
- Chen H, Yong W, Ren S, Shen W, He Y, et al. (2006) Overexpression of bone morphogenetic protein 10 in myocardium disrupts cardiac postnatal hypertrophic growth. J Biol Chem 281: 27481-27491.
- Ye L, Kynaston H, Jiang WG (2009) Bone morphogenetic protein-10 suppresses the growth and aggressiveness of prostate cancer cells through a Smad independent pathway. J Urol 181: 2749-2759.
- 31. McPherron AC, Lawler AM, Lee SJ (1999) Regulation of anterior/posterior

patterning of the axial skeleton by growth/differentiation factor 11. Nat Genet 22: 260-264.

- 32. Takayanagi H (2007) Osteoimmunology: shared mechanisms and crosstalk between the immune and bone systems. Nat Rev Immunol 7: 292-304.
- Weitzmann MN (2013) The role of inflammatory cytokines, the RANKL/OPG axis, and the immunoskeletal interface in physiological bone turnover and osteoporosis. Scienctifica 2013: 125705.
- Baron R, Neff L, Brown W, Louvard D, and Courtoy PJ (1990) Selective internalization of the apical plasma membrane and rapid redistribution of lysosomal enzymes and mannose 6-phosphate receptors during osteoclast inactivation by calcitonin. J Cell Sci 97: 439-447.
- 35. Teitelbaum SL (2000) Bone resorption by osteoclasts. Science 289: 1504-1508.
- Pacifici R, Brown C, Puscheck E, Friedrich E, Slatopolsky E, et al. (1991) Effect of surgical menopause and estrogen replacement on cytokine release from human blood mononuclear cells. Proc Natl Acad Sci U S A 88: 5134-5138.
- Rodan GA, Martin TJ (1981) Role of osteoblasts in hormonal control of bone resorption-a hypothesis. Calcif Tissue Int 33: 349-351.
- Simonet WS, Lacey DL, Dunstan CR, Kelley M, Chang MS, et al. (1997) Osteoprotegerin: a novel secreted protein involved in the regulation of bone density. Cell 89: 309-319.
- Yasuda H, Shima N, Nakagawa N, Mochizuki SI, Yano K, et al. (1998) Identity of osteoclastogenesis inhibitory factor (OCIF) and osteoprotegerin (OPG): a mechanism by which OPG/OCIF inhibits osteoclastogenesis in vitro. Endocrinology 139: 1329-1337.
- Yasuda H, Shima N, Nakagawa N, Yamaguchi K, Kinosaki M, et al. (1998) Osteoclast differentiation factor is a ligand for osteoprotegerin/ osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. Proc Natl Acad Sci USA 95:3597-3602.
- Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, et al. (1998) Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. Cell 93: 165-176.
- Anderson DM, Maraskovsky E, Billingsley WL, Dougall WC, Tometsko ME, et al. (1997) A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. Nature 390: 175-179.
- Wong BR, Rho J, Arron J, Robinson E, Orlinick J, et al. (1997) TRANCE is a novel ligand of the tumor necrosis factor receptor family that activates c-Jun N-terminal kinase in T cells. J Biol Chem 272: 25190-25194.
- 44. Suda T, Takahashi N, Udagawa N, Jimi E, Gillespie MT, et al. (1999) Modulation of osteoclast differentiation and function by the new members of the tumor necrosis factor receptor and ligand families. Endocr Rev 20: 345-357.
- Haynes DR, Crotti TN, Zreiqat H (2004) Regulation of osteoclast activity in peri-implant tissues. Biomaterials 25: 4877-4885.
- 46. Takayanagi H, Iizuka H, Juji T, Nakagawa T, Yamamoto A, et al. (2000) Involvement of receptor activator of nuclear factor kappaB ligand/osteoclast differentiation factor in osteoclastogenesis from synoviocytes in rheumatoid arthritis. Arthritis Rheum 43: 259-269.
- 47. Kotake S, Udagawa N, Hakoda M, Mogi M, Yano K, et al. (2001) Activated human T cells directly induce osteoclastogenesis from human monocytes: possible role of T cells in bone destruction in rheumatoid arthritis patients. Arthritis Rheum 44: 1003-1012.
- Lerner UH (2004) New Molecules in the Tumor Necrosis Factor Ligand and Receptor Superfamilies with Importance for Physiological and Pathological Bone Resorption. Crit Rev Oral Biol Med 15: 64-81.
- Boyle WJ, Simonet WS, Lacey DL (2003) Osteoclast differentiation and activation. Nature 423: 337-342.
- Mundy GR, Chen D, Oyajobi BO (2006) Bone remodeling. In: Favus MJ (ed) Primer on the metoabolic diseases and disorders of mineral metabolism. American society of bone mineral and research, Washington DC, pp: 46-58.
- Horowitz MC, Xi Y, Wilson K, Kacena MA (2001) Control of osteoclastogenesis and bone resorption by members of the TNF family of receptors and ligands. Cytokine Growth Factor Rev 12: 9-18.

- 52. Henderson B, Nair SP (2003) Hard labour: bacterial infection of the skeleton. Trends Microbiol 11: 570-577.
- Teng YT (2002) Mixed periodontal Th1-Th2 cytokine profile in Actinobacillus actinomycetemcomitans-specific osteoprotegerin ligand (or RANK-L)mediated alveolar bone destruction in vivo. Infect Immun 70: 5269-5273.
- Kikuchi T, Matsuguchi T, Tsuboi N, Mitani A, Tanaka S, et al. (2001) Gene expression of osteoclast differentiation factor is induced by lipopolysaccharide in mouse osteoblasts via Toll-like receptors. J Immunol 166: 3574-3579.
- Teng YT, Nguyen H, Gao X, Kong YY, Gorczynski RM, et al. (2000) Functional human T-cell immunity and osteoprotegerin ligand control alveolar bone destruction in periodontal infection. J Clin Invest 106: R59-67.
- Okahashi N, Inaba H, Nakagawa I, Yamamura T, Kuboniwa M, et al. (2004) Porphyromonas gingivalis induces receptor activator of NF-kappaB ligand expression in osteoblasts through the activator protein 1 pathway. Infect Immun 72: 1706-1714.
- Choi BK, Lee HJ, Kang JH, Jeong GJ, Min CK, et al. (2003) Induction of osteoclastogenesis and matrix metalloproteinase expression by the lipooligosaccharide of Treponema denticola. Infect Immun 71: 226-233.
- Okahashi N, Sakurai A, Nakagawa I, Fujiwara T, Kawabata S, et al. (2003) Infection by Streptococcus pyogenes induces the receptor activator of NFkappaB ligand expression in mouse osteoblastic cells. Infect Immun 71:948-955.
- Sakurai A, Okahashi N, Nakagawa I, Kawabata S, Amano A, et al. (2003) Streptococcus pyogenes infection induces septic arthritis with increased production of the receptor activator of the NF-kappaB ligand. Infect Immun 71: 6019-6026.
- Mizuno A, Kanno T, Hoshi M, Shibata O, Yano K, et al. (2002) Transgenic mice overexpressing soluble osteoclast differentiation factor (sODF) exhibit severe osteoporosis. J Bone Miner Metab 20: 337-344.
- Nakashima T, Kobayashi Y, Yamasaki S, Kawakami A, Eguchi K, et al. (2000) Protein expression and functional difference of membrane-bound and soluble receptor activator of NF-kappaB ligand: modulation of the expression by osteotropic factors and cytokines. Biochem Biophys Res Commun 275: 768-775.
- Wei S, Kitaura H, Zhou P, Ross FP, Teitelbaum SL (2005) IL-1 mediates TNFinduced osteoclastogenesis. J Clin Invest 115: 282-290.
- Han X, Kawai T, Eastcott JW, Taubman MA (2006) Bacterial-responsive B lymphocytes induce periodontal bone resorption. J Immunol 176: 625-631.
- Jin Q, Cirelli JA, Park CH, Sugai JV, Taba M Jr, et al. (2007) RANKL inhibition through osteoprotegerin blocks bone loss in experimental periodontitis. J Periodontol 78: 1300-1308.
- Bar-Shavit Z (2007) The osteoclast: a multinucleated, hematopoietic-origin, bone-resorbing osteoimmune cell. J Cell Biochem 102: 1130-1139.
- 66. Lewiecki EM (2009) Denosumab update. Curr Opin Rheumatol 21: 369-373.
- 67. Cochran DL (2008) Inflammation and bone loss in periodontal disease. J Periodontol 79: 1569-1576.
- Bostanci N, Ilgenli T, Emingil G, Afacan B, Han B, et al. (2007) Gingival crevicular fluid levels of RANKL and OPG in periodontal diseases: implications of their relative ratio. J Clin Periodontol 34: 370-376.
- Bostanci N, Ilgenli T, Emingil G, Afacan B, Han B, et al. (2007) Differential expression of receptor activator of nuclear factor-kappaB ligand and osteoprotegerin mRNA in periodontal diseases. J Periodontal Res 42: 287-293.
- Mogi M, Otogoto J, Ota N, Togari A (2004) Differential expression of RANKL and osteoprotegerin in gingival crevicular fluid of patients with periodontitis. J Dent Res 83: 166-169.
- Liu D, Xu JK, Figliomeni L, Huang L, Pavlos NJ, et al. (2003) Expression of RANKL and OPG mRNA in periodontal disease: possible involvement in bone destruction. Int J Mol Med 11: 17-21.
- Kawai T, Matsuyama T, Hosokawa Y, Makihira S, Seki M, et al. (2006) B and T lymphocytes are the primary sources of RANKL in the bone resorptive lesion of periodontal disease. Am J Pathol 169: 987-998.
- 73. Wara-aswapati N, Surarit R, Chayasadom A, Boch JA, Pitiphat W (2007)

RANKL upregulation associated with periodontitis and Porphyromonas gingivalis. J Periodontol 78: 1062-1069.

- Garlet GP, Martins W Jr., Fonseca BA, Ferreira BR, Silva JS (2004) Matrix metalloproteinases, their physiological inhibitors and osteoclast factors are differentially regulated by the cytokine profile in human periodontal disease. J Clin Periodontol 31:671-679.
- Nagasawa T, Kobayashi H, Kiji M, Aramaki M, Mahanonda R, et al. (2002) LPS-stimulated human gingival fibroblasts inhibit the differentiation of monocytes into osteoclasts through the production of osteoprotegerin. Clin Exp Immunol 130: 338-344.
- Valverde P, Kawai T, Taubman MA (2004) Selective blockade of voltage-gated potassium channels reduces inflammatory bone resorption in experimental periodontal disease. J Bone Miner Res 19: 155-164.
- 77. Kawai T, Paster BJ, Komatsuzawa H, Ernst CW, Goncalves RB, et al. (2007) Crossreactive adaptive immune response to oral commensal bacteria results in an induction of receptor activator of nuclear factor-kappaB ligand (RANKL)dependent periodontal bone resorption in a mouse model. Oral Microbiol Immunol 22: 208-215.
- Mahamed DA, Marleau A, Alnaeeli M, Singh B, Zhang X, et al. (2005) G(-) anaerobes-reactive CD4+ T-cells trigger RANKL-mediated enhanced alveolar bone loss in diabetic NOD mice. Diabetes 54: 1477-1486.
- Rogers JE, Li F, Coatney DD, Otremba J, Kriegl JM, et al. (2007) A p38 mitogen-activated protein kinase inhibitor arrests active alveolar bone loss in a rat periodontitis model. J Periodontol 78: 1992-1998.
- Parfitt AM (1982) The coupling of bone formation to bone resorption: a critical analysis of the concept and of its relevance to the pathogenesis of osteoporosis. Metab Bone Dis Relat Res 4: 1-6.
- 81. Al-Mashat HA, Kandru S, Liu R, Behl Y, Desta T, et al. (2006) Diabetes

enhances mRNA levels of proapoptotic genes and caspase activity, which contribute to impaired healing. Diabetes 55: 487-495.

- Liu R, Bal HS, Desta T, Krothapalli N, Alyassi M, et al. (2006) Diabetes enhances periodontal bone loss through enhanced resorption and diminished bone formation. J Dent Res 85: 510-514.
- Bertolini DR, Nedwin GE, Bringman TS, Smith DD, Mundy GR (1986) Stimulation of bone resorption and inhibition of bone formation in vitro by human tumour necrosis factors. Nature 319: 516-518.
- Nguyen L, Dewhirst FE, Hauschka PV, Stashenko P (1991) Interleukin-1 beta stimulates bone resorption and inhibits bone formation in vivo. Lymphokine Cytokine Res 10: 15-21.
- Diarra D, Stolina M, Polzer K, Zwerina J, Ominsky MS, et al. (2007) Dickkopf-1 is a master regulator of joint remodeling. Nat Med 13: 156-163.
- Tsuboi M, Kawakami A, Nakashima T, Matsuoka N, Urayama S, et al. (1999) Tumor necrosis factor-alpha and interleukin-1beta increase the Fas-mediated apoptosis of human osteoblasts. J Lab Clin Med 134: 222-231.
- Thammasitboon K, Goldring SR, Boch JA (2006) Role of macrophages in LPS-induced osteoblast and PDL cell apoptosis. Bone 38: 845-852.
- Centrella M, McCarthy TL, Canalis E (1988) Tumor necrosis factor-alpha inhibits collagen synthesis and alkaline phosphatase activity independently of its effect on deoxyribonucleic acid synthesis in osteoblast-enriched bone cell cultures. Endocrinology 123: 1442-1448.
- Taichman RS, Hauschka PV (1992) Effects of interleukin-1 beta and tumor necrosis factor-alpha on osteoblastic expression of osteocalcin and mineralized extracellular matrix in vitro. Inflammation 16: 587-601.
- Eguchi Y, Wakitani S, Imai Y, Naka Y, Hashimoto Y, et al. (2010) Antitumor necrotic factor agent promotes BMP-2-induced ectopic bone formation. J Bone Miner Metab 28: 157-164.

This article was originally published in a special issue, entitled: "Innate Response to Infectious Diseases", Edited by Dr. Anshu Agrawal, University of California Irvine, USA Page 8 of 8