Short Commentary Open Access

Transcriptome Analysis in Banana

Kalibulla Syed Ibrahim, Nachimuthu Senthil Kumar and Robert Thangjam*

Department of Biotechnology, School of Life Sciences, Mizoram University, Aizawl - 796004, Mizoram, India

Short Commentary

DNA sequencing has become an inevitable platform for molecular biological studies. As a rapid and inexpensive method, the next-generation sequencing (NGS), offer high throughput through transcriptome profiling system called RNA-Seq [1]. As next-generation sequencing technologies provides deep coverage and at single base-pair resolution, RNA sequencing is preferred as an alternative to wholegenome sequencing as it analyzes only the transcribed portions of the genome [2].

NGS based transcriptome analysis have also been successfully applied to gene expression profiling by sequencing different mRNA species that measure the activity of thousands of genes in parallel and also quantify transcripts [3]. This helps in understanding the differential expression of genes at different biological conditions of cells and or tissues [4]. Besides that pathways, stress and environmental factors can be analyzed on a genome-wide scale using NGS technologies.

Banana (Musa spp.) belonging to the family Musaceae of the order Zingiberales is one of the most important food crops worldwide. It is derived from intra and inter specific crosses between M. acuminata colla (A genome) and M. balbisiana colla (B genome) which resulted to many genome groups such as AA, AB, AAA, AAB, ABB, AABB, AAAB and ABBB [5]. South-East Asia is considered as centre of origin of Musa species, where numerous wild species occur in an area stretching from Papua New Guinea to India. The major issues in the banana production are pest and diseases. Some pest and diseases are highly aggressive, easily spread and difficult to eradicate which dramatically reduced the yield affecting the economical balance in many banana producing states and countries [6]. The described draft sequence of 523-megabase genome of M. acuminata doubled-haploid genotype, providing a crucial stepping stone for genetic improvement of banana has been recently published [7]. Thus, banana is a good candidate for functional genomics because of its relatively small genome size and important characteristics such as climacteric fruit, parthenocarpy or vegetative propagation, which are absent in model organisms [8]. Recent advances in genetics and genomics including the use of molecular markers have enhanced our understanding of structural and functional aspects of plant genomes that can enhance our ability to improve banana [9]. The development of efficient and cost effective in vitro regeneration methods also has provided valuable options for improved production of superior planting

Musa based Transcriptome analysis has been performed on root [2] and leaf [11]. Recently, a whole transcriptome from different tissues like leaf, root, sheath, flower bud, flower bract, pulp, and rhizome was performed by Backiyarani [12]. The identification of rapid activation and selective induction of cold tolerance genes ICE1 and MYBS3 and their pathways in plantain and banana, along with expression of other cold-specific genes provided useful information on the expression patterns using comparative transcriptomics [13]. Sequencing of the genomes and transcriptomes of two Fusarium oxysporum f. sp. cubense (Foc) isolates, the pathogens of banana that causes vascular wilt disease lead to the identification of genes putatively involved in root attachment, cell degradation, detoxification of toxin, transport, secondary metabolites biosynthesis and signal transductions and more importantly transporters and transcription factors for toxins and nutrients transport that may facilitate its ability to adapt to host

environments and contribute to pathogenicity to banana [14]. The transcriptome analysis of the banana fruit during the ripening process revealed the underlying mechanism of ripening and also to the identification of a large number of novel differentially regulated genes associated with cell wall degradation and synthesis of aromatic volatiles [15].

Transcriptome analysis helps in interpreting the functional elements of the genome [16,17] and along with the recent introduction of next-generation deep sequencing could provide a platform for genetic improvement of economically viable crops such as banana or potato [18]. Its application in Musa could improve our understanding of the molecular mechanisms involved in defence [14], biosynthesis, and other biochemical processes [12,13,15] providing a potential resource for future genetic or genomic studies on this important food crop. It can also be expected to bridge a critical gap that exists in the Musa comparative genomics and could ultimately contribute to the evolutionary and functional studies of its genes and genomes.

Acknowledgements

The authors wish to acknowledge the Department of Biotechnology (DBT), Govt. of India, New Delhi for the financial support in the form of Bioinformatics Infrastructure Facility (BT/BI/12/060/2012) and the State Biotech Hub (BT/04/NE/2009).

References

- Mardis ER (2008) Next-generation DNA sequencing methods. Annu Rev Genomics Hum Genet 9: 387-402.
- Li CY, Deng GM, Yang J, Viljoen A, Jin Y, et al. (2012) Transcriptome profiling
 of resistant and susceptible Cavendish banana roots following inoculation with
 Fusarium oxysporum f. sp. cubense tropical race 4. BMC Genomics 13: 374.
- Voelkerding KV, Dames SA, Durtschi JD (2009) Next-generation sequencing: from basic research to diagnostics. Clin Chem 55: 641-658.
- Cullum R, Alder O, Hoodless PA (2011) The next generation: using new sequencing technologies to analyse gene regulation. Respirology 16: 210-222.
- Simmonds NW and Shepherd K (1955) The taxonomy and origins of the cultivated bananas. Linn Soc Bot 55: 302-312.
- Heslop-Harrison JS, Schwarzacher T (2007) Domestication, genomics and the future for banana. Ann Bot 100: 1073-1084.
- D'Hont A, Denoeud F, Aury JM, Baurens FC, Carreel F, et al. (2012) The banana (Musa acuminata) genome and the evolution of monocotyledonous plants. Nature 488: 213-217.
- Sági L, Remy S, Coemans B, Thiry E, Santos E, et al. (2005) Functional analysis of the banana genome by gene tagging and SAGE. Plant and Animal Genomes XIII Conference, San Diego, CA, USA.
- 9. Pachuau L, Atom AD, Thangjam R (2014) Genome classification of Musa

*Corresponding author: Robert Thangjam, Department of Biotechnology, School of Life Sciences, Mizoram University, Aizawl-796004, Mizoram, India, Tel: +91 9436352386; E-mail: robertthangjam@gmail.com

Received October 07, 2015; Accepted November 11, 2015; Published November 16, 2015

Citation: Ibrahim KS, Kumar NS, Thangjam R (2015) Transcriptome Analysis in Banana. Transcriptomics 3: 117. doi:10.4172/2329-8936.1000117

Copyright: © 2015 Ibrahim KS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

- cultivars from northeast India as revealed by ITS and IRAP markers. Appl Biochem Biotechnol 172: 3939-3948.
- Hrahsel L, Basu A, Sahoo L, Thangjam R (2014) In vitro propagation and assessment of the genetic fidelity of Musa acuminata (AAA) cv. Vaibalhla derived from immature male flowers. Appl Biochem Biotechnol 172: 1530-1530
- 11. Passos MA, de Oliveira Cruz V, Emediato FL, de Camargo Teixeira C, Souza MT Jr, et al. (2012) Development of expressed sequence tag and expressed sequence tag-simple sequence repeat marker resources for Musa acuminata. AoB Plants 2012: pls030.
- Backiyarani S, Uma S, Saraswathi MS, Saravanakumar AS, Chandrasekar A (2015) Transcriptome analysis of banana (Musa balbisiana) based on nextgeneration sequencing technology. Turk J Agric.
- 13. Yang QS, Gao J, et al. (2015) Comparative transcriptomics analysis reveals difference of key gene expression between banana and plantain in response to

- cold stress. BMC Genomics 16: 446.
- 14. Guo L, Han L, Yang L, Zeng H, Fan D, et al. (2014) Genome and transcriptome analysis of the fungal pathogen Fusarium oxysporum f. sp. cubense causing banana vascular wilt disease. PLoS One 9: e95543.
- Asif MH, Lakhwani D, Pathak S, Gupta P, Bag SK, et al. (2014) Transcriptome analysis of ripe and unripe fruit tissue of banana identifies major metabolic networks involved in fruit ripening process. BMC Plant Biol 14: 316.
- Wang Y, Chung SJ, Song WO, Chun OK (2011) Estimation of daily proanthocyanidin intake and major food sources in the U.S. diet. J Nutr 141: 447-452
- Wang Z, Gerstein M, Snyder M (2009) RNA-Seq: a revolutionary tool for transcriptomics. Nat Rev Genet 10: 57-63.
- 18. Hobert O (2010) The impact of whole genome sequencing on model system genetics: get ready for the ride.