

Understanding How Researchers could Generate Ovarian Follicles from Mouse Stem Cells

Anshika Raj*

Department of Nursing, Chandigarh University, India

INTRODUCTION

Researchers from Kyushu University have succeeded in reconstructing components in the ovarian environment that are necessary for the production of mature egg cells using mouse stem cells. They were also able to use their discoveries to grow healthy mice and manufacture functional egg cells.

The researchers hope that by knowing the underlying reasons of infertility, they will be able to treat it and help save critically endangered animals by producing egg cells [1].

Understanding how a single cell with the potential to become anything may specialize into cells that make up an organ or tissue, a process known as 'differentiation,' is a key component of stem cell research.

A researcher at Kyushu University has been studying the mechanisms that govern the formation and development of the oocyte, which is the most basic cell in the human body.

The oocyte is the immature stage of the ovum, commonly known as an egg cell, according to Hayashi, the study's lead author. "Many significant advances have been made in the field of oocyte development, including the generation of mouse and human oocytes from their respective stem cells and the breeding of healthy mice."

However, creating the correct support system to assist differentiate stem cells into oocytes is a key stumbling block in the process. Primary oocytes begin to form in the body during adolescence, with the help of other cephalic cells. The ovarian follicle is the collective aggregate that eventually generates a mature egg cell [2,3].

"While we were able to create mouse oocytes from stem cells, we needed to unite them with cells called gonadal somatic cells that were removed from mice in order for them to mature," says first author Takashi Yoshino, an assistant professor at the Faculty of Medical Sciences.

"We must be able to build the support system from stem cells if we wish to better understand oocyte development and apply this technology to endangered animals and even people." The team's initial steps was to research mice and determine which genes and when they are produced are responsible for gonadal somatic cells. Following the discovery, they investigated mouse stem cells under various growing settings until they were able to develop cells with similar characteristics [4].

"Because these cells have the same genetic expression profiles as gonadal somatic cells, we term them foetal ovarian somatic cell-like cells, or FOSLCs. The next step was to combine the FOSLCs with the stem-cell-derived primordial germ cells—oocyte precursors—that we'd been producing in previous research to see if they matured into oocytes" Yoshino explains.

They were able to see the cells begin to form follicular structures around the oocytes after several days of growth.

After confirming that these cells were fully formed oocytes, the eggs were fertilised and put into a surrogate mouse, where they successfully produced pups.

The new approach is expected to make it possible to produce a large number of oocytes without having to take cells from the host animals.

"This will greatly aid our understanding of ovarian follicle formation and reproductive biology in general," Hayashi says.

"While the application on humans is theoretically viable, we must carefully consider and evaluate the ethics and technology of its application due to the genetic and epigenetic instability of in vitro generated gametes [5]." We'll keep working to understand the mechanisms behind this basic and one-of-a-kind extension of life."

REFERENCES

1. Yoshino T, Suzuki T, Nagamatsu, G, Yabukami H, Ikegaya M, Kishima M, et al. Generation of ovarian follicles from mouse pluripotent stem cells. *Science*. 2021; 373: eabe0237.
2. Novak I, Lightfoot DA, Wang H, Eriksson A, Mahdy E, Höög C. Mouse embryonic stem cells form follicle like ovarian structures but do not progress through meiosis. *Stem Cells*. 2006; 24: 1931-1936.
3. Qing T, Shi Y, Qin H, Ye X, Wei W, Liu H, et al. Induction of oocyte-like cells from mouse embryonic stem cells by co-culture with ovarian granulosa cells. *Differentiation*. 2007; 75: 902-911.
4. Pacchiarotti J, Maki C, Ramos T, Marh J, Howerton K, Wong J, et al. Differentiation potential of germ line stem cells derived from the postnatal mouse ovary. *Differentiation*. 2010; 79: 159-170.
5. <https://www.technologynetworks.com/cell-science/news/ovarian-follicles-generated-from-mouse-stem-cells-353647>.

*Correspondence to: Anshika Raj, Department of Nursing, Chandigarh University, India, E-mail: rajanshika@gmail.com

Received: September 03, 2021; Accepted: September 17, 2021; Published: September 24, 2021

Citation: Raj A (2021) Understanding How Researchers could Generate Ovarian Follicles from Mouse Stem Cells. *J Fertil In vitro IVF Worldw Reprod Med Genet Stem Cell Biol* 9:5. doi: 10.35248/2375-4508.21.9.246.

Copyright: © 2021 Raj A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.